

Status of anthracnose disease cause by *Colletotrichum lindemuthianum* in major mungbean growing area of Rajasthan

ABSTRACT

Anthracnose caused by *Colletotrichum lindemuthianum* is one of the most important diseases in the mungbean. A roving survey to assess the intensity of anthracnose of mungbean was carried out at 60 locations comprising five districts of Rajasthan state during *Kharif* season of year 2022-23. The most extreme disease frequency was seen in plants at a flowering and podding stage. The overall mean disease intensity was observed to be 22.60%. Among the five districts lowest anthracnose intensity was noticed in Sikar (17.88%) followed by Nagaur (18.90%), while highest anthracnose intensity was recorded in Jaipur district (28.93%) followed by Ajmer (24.64%) and Tonk (22.65%).

Key-Words: Survey, Mungbean, Anthracnose, Disease intensity

INTRODUCTION

Mungbean [*Vigna radiata* (L.) Wilczek] is a short-duration food grain legume cultivated over 7 million hectares, predominantly across Asia and rapidly spreading to other parts of the world. The major mungbean growing states are Orissa, Maharashtra, Andhra Pradesh, Rajasthan, Karnataka and Gujarat (Khairi *et al.*, 2024). It ranks third among all pulses grown in India after chickpea and pigeonpea. Mungbean seeds are rich in proteins (24%), fiber, antioxidants, and phytonutrients (Itoh *et al.*, 2006). In general, the low productivity of mungbean is due to its cultivation in marginal lands, low rainfall areas, high rate of flower and fruit drop, non-uniform maturity, pod shattering and susceptibility to abiotic and biotic constraints, poor crop management practices and non availability of quality seeds of improved varieties to farmers are responsible for low productivity (Chauhan *et al.*, 2010 and Pratap *et al.*, 2019). Among biotic stresses, fungal, viral and bacterial diseases are the major factor in the reducing yield and quality in most of the regions of India can reduce the yield up to 40-60 per cent in mungbean crop (Kaur *et al.*, 2011; Khairi and Hake, 2018).

Anthracnose of mungbean caused by *Colletotrichum lindemuthianum* (Sacc. and Magn.) is one of the most important seed borne disease (Parthiban and Kavitha, 2014; Misal *et al.*, 2019). Mungbean anthracnose is an economically important disease results in a yield losses upto 30 to 70 per cent (Kulkarni, 2009 & Shukla *et al.*, 2014). The early signs of infection usually appear on the lower leaf surface along the veins, which show brick red to purplish red discoloration. Later, such discoloration also appears on the upper leaf surface. At the same time, brown lesions of various sizes, with black, brown or purplish red margins, develop around small veins (Alien *et al.*, 1996). Very little work has been done on systematic survey of this disease in Rajasthan. Hence, present investigation was initiated on survey for anthracnose in major mungbean growing

Commented [PK1]: Add the introduction and methods use for the experiment of the crop in 2 lines before starting the results part in the abstract

Commented [PK2]: Use alternate keywords which is not included in the title

Formatted: Font: Italic

Formatted: Font: Italic

districts of Rajasthan, to identify the intensity of the disease over time and geographical locations.

MATERIALS AND METHODS

A roving survey was conducted in the five major mungbean growing districts of Rajasthan viz., Jaipur, Tonk, Ajmer, Nagaur and Sikar during *Kharif* season 2022-23 to find out the intensity of anthracnose disease. Two tehsil selected under each district were surveyed. Under each tehsil two villages were selected and under each village three farmers' fields were assessed. The sample were collected, isolated, purified, identified and were subjected to pathogenic test.

The anthracnose severity was recorded on ten randomly selected mungbean plant per field by using 0-9 scale given by Mayee and Dattar, 1986 (Table 1). On the basis of numerical rating per cent disease intensity (PDI) was calculate applying the formula given by Mckinney (1923)

$$PDI = \frac{\text{Sum of all numerical rating}}{\text{No. of leaves/plant observed}} \times \frac{100}{\text{Maximum disease rating}}$$

Table 1. Anthracnose disease rating scale of mungbean given by Mayee and Dattar 1986
(This scale is used for the screening the varietal disease lines to check the reaction of the disease on the particular variety, not for the survey purpose)

(Use this paper to correct the mayee and datar scale for the survey purpose: DOI: <https://doi.org/10.60151/envec/NHEM2312>)

Grade	Disease-reaction	Disease-intensity (%)	Reaction-group
0	Highly-resistant	0-1	HR
1	Resistant	1.1-10	R
3	Moderately resistant	10.1-20	MR
5	Moderately susceptible	20.1-30	MS
7	Susceptible	30.1-50	S
9	Highly susceptible	≥50	HS

RESULTS AND DISCUSSION

Survey for the occurrence and distribution of Anthracnose of mungbean at farmer's field of Rajasthan district

Data pertaining to survey conducted during *Kharif* season 2022-23 as presented in Table 1 revealed that, Amongamong all five districts maximum mean per cent disease intensity (PDI) of anthracnose disease of mungbean was recorded in Jaipur district (28.93%) followed by Ajmer (24.64%) Tonk (22.65%) and Nagaur (18.90%) whereas minimum disease intensity was recorded in Sikar district having PDI 17.88 per cent. There was significant difference in mean PDI were observed in among the five districts

Formatted: Font: Not Italic

The anthracnose severity in Jaipur district ranged from 23.07 per cent (Nimera) to 35.26 per cent (Sewa). The disease index ranged between 20.56 (Juniya) to 27.90 per cent (Bhamolav) in Ajmer district. In Nagaur district, the disease ranged between 16.30 per cent (Bherunda) to 22.40 per cent (Nawa). In Tonk, the disease index varied from 11.06 (Deoli) to 27.88 per cent (Nagar). In Sikar district, the disease index ranged between 9.07 per cent (Shyamgarh) to 22.65 per cent (Mau). Similar result reported by Kulkarni and Benagi (2013) conducted a roving survey to assess the intensity of mungbean anthracnose during 2007 and 2008 in eleven major mungbean growing districts of northern Karnataka. The results revealed that during 2007 anthracnose of mungbean was noticed in the range of 21.36 to 58.97 per cent and during 2008, the disease severity noticed in the range of 24.67 to 60.07 per cent. Roopadevi *et al.* (2015) found the disease incidence and severity of anthracnose in the major mungbean regions of northern Karnataka. The disease severity was recorded from 28.1 to 62.7 per cent. The highest disease severity 59.8 per cent was obtained in Bidar district followed by Gulbarga (53.8%) and Bijapur (38.4%). Doganya *et al.* (2021) observed disease intensity between 20.0 to 29.60 per cent with an average intensity of 26.06 per cent of mungbean anthracnose at Khandwa district of Madhya Pradesh. While Patil (2022) noted 16.22 to 28.0 per cent disease intensity of mungbean anthracnose at Sehore district of Madhya Pradesh.




Fig. 1. Disease intensity of anthracnose disease in major mungbean growing district of Rajasthan

Table 2: Survey on severity of anthracnose of mungbean (*Colletotrichum lindemuthianum*) in different villages of the districts in Rajasthan during -Kharif, 2022-23

Districts	Tehsils	Villages	No. of fields	Per cent disease intensity in surveyed field			Avg. disease intensity (avg. of three fields)	Avg. disease intensity (Tehsil)	District Mean
Jaipur	Dudu	Sewa	3	32	35.20	38.50	35.26	32.89	28.93
		Mangalwara	3	28.66	32.8	30.11	30.52		
	Phagi	Kansya	3	27.71	29.12	23.69	26.84	24.96	
		Nimera	3	19.21	22.43	27.57	23.07		
Ajmer	Kishangarh	Bhamolav	3	32.89	26.90	23.90	27.90	26.62	24.64
		Chota lamba	3	23.78	24.34	27.91	25.34		
	Kekri	Meodakalan	3	22.10	28	24.21	24.77	22.67	
		Juniya	3	17.8	23.11	20.78	20.56		
Nagaur	Nawa	Moondgasoi	3	20.74	23.62	22.86	22.40	23.28	18.90
		Panchota	3	24.66	25.69	22.13	24.16		
	Riyanbadi	Morikalan	3	13.86	5.67	18.71	12.74	14.52	
		Bherunda	3	19	17.18	12.72	16.30		
Tonk	Malpura	Nagar	3	30	25.86	27.78	27.88	27.16	22.65
		Pachewar	3	28.22	26.21	24.92	26.45		
	Deoli	Deoli	3	8.08	10.12	14.98	11.06	18.15	
		Dooni	3	25.37	26.24	24.1	25.23		
Sikar	Khandela	Shyamgarh	3	11.04	7.21	8.96	9.07	11.7	17.88
		Barsinghpura	3	13.83	19.21	9.96	14.33		
	Srimadhapur	Arniya	3	20.14	19	26.2	21.78	22.22	
		Mau	3	24.60	19.56	23.8	22.65		
Overall mean							22.60		

Conclusions

Total sixty places visited under ten tehsils of five district in Rajasthan viz., Jaipur, Ajmer, Tonk, Nagaur and Sikar. During survey, anthracnose was found at all the sites and the intensity ranged from 11.7 to 32.89 per cent. Overall mean disease intensity of anthracnose of mungbean was observed 22.60 per cent under surveyed districts of Rajasthan. Lowest anthracnose intensity was noticed in Khandela (11.70 %), while highest anthracnose intensity was recorded in Dudu (32.89 %) Among the district, minimum anthracnose intensity was recorded in Sikar, while maximum anthracnose intensity was recorded in Jaipur (28.93%).

REFERENCES

- Alien DJ, Ampofo JKO, Wortman CS. Pest, disease, and nutritional disorders of the common bean in Africa: A Field Guide. *Centro Internacional de Agricultura Tropical*, Cali Colombia. 1996
- Chauhan YS, Douglas C, Rachaputi RCN, Agius PM and King K. "Physiology of mungbean and development of the mungbean crop model," in Proceedings of the 1st Australian Summer Grains Conference Australia, Gold Coast, QL. 2010; 21–24.
- Datar VV. "Phytopathometry". Technical Bulletin-I, Marathwada Agricultural University, Parbhani, India, 1986;146.
- Doganya D, Bobade A, Jain VK, Arsia SK, Vani DK. Survey of anthracnose of green gram in East Nimar region of Madhya Pradesh. *Prog. Res.: An Int. J.* 2021;16 (1): 8-13.
- Kaur L, Singh P, Sirari A. (). Biplot analysis for locating multiple disease resistant diversity in disease resistant diversity in mungbean germplasm. *Disease Res.* 2011; 26: 55–60.
- Kulkarni AS, Epidemiology and integrated management of anthracnose of green gram, Ph. D. (Agri.) Thesis, Univ. Agric. Sci., Dharwad (India).2009
- Kulkarni S, Benagi VI. Survey for the status of anthracnose of greengram in Northern Karnataka . *Int. J. Agri. Sci.*, 2013; 9(1): 1-13.
- Mayee CD, Itoh T, Garcia RN, Adachi M, Maruyama Y, Tecson-Mendoza EM, Mikami B. Structure of 8S α globulin, the major seed storage protein of mung bean. *Acta Crystallogr. D. Biol. Crystallogr.* 2006; 62, 824–832. doi: 10.1107/S090744490601804X
- McKinney GM. A new system of grading plant discase. *J. Agric. Res.* 1923; 26 : 195- 218.
- Parthiban VK, Kavitha R. In vitro screening of effective biocontrol agents against bean anthracnose pathogen, *Colletotrichum lindemuthianum*. *Int. J. of Pharm. Scr. Meth.* 2014; 4(1): 32-35.

Patil V. Management of *Colletotrichum lindemuthianum* (Sacc. and Magnus) Briosi and Cav. causing anthracnose of Green gram, M.Sc. (Agri.) Thesis submitted to RVSKVV, Gwalior, Madhya Pradesh. 2022.

Pratap A, Gupta S, Basu S, Tomar R, Dubey S, Rathore M. "Towards Development of Climate-Smart Mungbean: Challenges and Opportunities," in Genomic Designing of Climate Smart Pulse Crops. Ed. C. Kole (New York: Springer Nature). 2019; (In press). doi: 10.1007/978-3-319-96932-9_5

Roopadevi B., Jamadar MM. and Anusha BG. (2015). Survey for incidence and severity of greengram [*Vigna radiata* (L.) Wilczek] anthracnose caused by *Colletotrichum truncatum* (Schw.) Andrus and Moore. Trends in Biosci. 7(23): 3941-3943.

Shukla, V., Baghel, S., Maravi, K. and Singh, SK. (2014). Yield loss assessment in mungbean [*Vigna radiata* (L.) Wilczek] caused by anthracnose [*colletotrichum truncatum* (schw.)]. Journal Research of Punjab Agriculture University, 6:345-348.

Wilczek R. *Vigna radiata* (L.) R. Wilczek Flore du Congo Belge et du Ruanda-Urundi. 1954;6:343-393.

[Khaire PB, Hake LG \(2018\) Disease Management of kharif Green Gram and Black Gram. Popular Kheti 6 \(2\): 96-103.](#)

<https://www.researchgate.net/publication/333203658> Disease Management of Kharif Green Gram and Black Gram

[Pravin Babasaheb Khaire, Dilipkumar Gangadhar Hingole, Someshree Sampatrao Mane, Vaibhav Shelar \(2024\) Exploring the Role of Environmental Factors on the Emergence of Leaf Blight of Mungbean in Maharashtra State, India. Environment and Ecology 42 \(2B\) : 846–850, April–June 2024 Article DOI: https://doi.org/10.60151/envec/NHEM2312](#)

[Misal DM, Khaire PB, Misal MR, Hingole DG \(2019\) Integrated evaluation of fungicides, botanicals and bioagents against anthracnose of mungbean on natural field condition. International Journal of Chemical Studies 7\(2\): 1975-1978](#)

UNDER PEER REVIEW

