

Original Research Article

Evaluation *in vitro* potential for plant growth promotion detected in nodule endophyte bacteria of Bambara groundnut (*Vigna Subterranea* L.) from agricultural soils in Cote d'Ivoire

ABSTRACT

Legumes-endophytes association is an integral part in improving agricultural productivity and soil fertility. Some endophytes which are considered as PGPR can promote plant growth through mobilization of nutrients in soils, production of plant growth regulators, protection of plants from phytopathogens by controlling or inhibiting them, and improvement of soil structure.

In order to select the most effective endophytic bacteria to be used as biofertilizers, the objective of this study is to evaluate the PGPR potential of bacteria isolated from Bambara groundnut nodules *in vitro*. To accomplish this, 34 endophytic bacteria isolated from the nodules of 5 varieties of Bambara groundnut in soils from 5 localities of Daloa, which had been the subject of molecular characterization, were tested to evaluate their capacity to produce IAA and siderophores, to solubilize phosphate.

Generally, all strains produced a detectable amount of IAA which ranged from 16.3 to 275.5 µg/ml, and the four (4) major producers of IAA were RFK12, RFK34, RFK10 and RFK2. The range of phosphate solubilization was 16 to 453.45 µg/ml, meanwhile four strains (RFK27, RFK11, RFK8 and RFK12) were the most efficient P-solubilizers. The majority (94 % of strains) were able to produce siderophore. The halo diameter varies from 7.06 to 40 mm and siderophore production index ranged from 1.08 to 5.36 and the isolate that showed the best ability in siderophore production was RFK27, followed by RFK26 and RFK22.

In total, 19 strains showed multi-PGP traits and could be promising biofertilizers.

In perspective, it would be interesting to test the 19 strains with multi-PGPR traits under field conditions.

Keywords: Endophyte PGPR Biofertilizer Bambara groundnut... *In vitro*

1. Introduction

The high productivity of intensive agriculture is largely dependent on synthetic chemical inputs, particularly fertilizers and pesticides, with projections predicting the use of 120 million tons by 2040 (Vance, 2001; Soussou, 2013). However, improper use of these synthetic chemical inputs has upset the ecosystem's equilibrium and is the basis of several human, economic and environmental problems such as the contamination of agricultural soils (Soussou, 2013). Therefore, in order to restore soil quality for agricultural production, appropriate soil management and conservation

Commented [AB1]:

Commented [AB2]: Rewrite

Commented [AB3]: Full form ? correct it

Commented [AB4]: Were they identified as strain ?

Commented [AB5]: Please clarify isolate or strain

techniques are imperative, specifically bio-fertilization based on soil microorganisms. (Hussain et al., 2021).

Microorganisms serve as a reservoir of nutrients for plants and also contribute to soil structure. They also provide other benefits. (Imran et al., 2019a). Bacteria are the most numerous soil organisms and play an important role in plant growth and development (Imran et al., 2019b ; Khan, 2018). Hence some bacteria are named PGPR (Gupta et al., 2015). PGPR (Plant Growth Promoting Rhizobacteria) are bacteria able to promote plant growth and serve as biofertilizers and biopesticides in agricultural crops (Ferchichi et al., 2019). This bacteria may promote the plant growth through different direct and indirect mechanisms such as: fixation of nitrogen, solubilization of soil phosphates, production of hormones (auxin and gibberellin), production of siderophores (iron chelator), increasing the uptake of water and minerals, increasing the enzymatic activity of the plant. The PGPR may promote the plant growth also by suppressing plant pathogens (Pérez-Montaña et al., 2014; Zaatri & Tagzirt, 2023).

Thus, to promote sustainable agriculture, it is necessary to produce effective bacterial inocula capable of improving plant productivity. This work aims to evaluate the PGPR potential of endophytic bacteria isolated from Bambara pea nodules *in vitro* in order to select the most efficient ones and use them as biofertilizers.

2. MATERIALS AND METHODS

2.1. Bacterial strains

The bacteria were isolated *in vitro* at the Jean Lorougnon Guédé University in Daloa, Côte d'Ivoire. These isolates were tested for their genetic diversity and Plant Growth Promoting (PGP) activities in a laboratory at Mohammed V's Faculty of Science in Rabat, Morocco. The 34 bacterial strains examined in this study had previously undergone morphological and molecular characterization and their partial sequences of the 16S rRNA were deposited in the GenBank® database under the accession numbers MT661489 to MT661522. (GUE et al., 2024)

2.2. Reactivation and verification of strain purity

The strains stored in 20% (v/v) of glycerol at -80°C were removed from the freezer and their purity was checked by streaking technique according to the Jordan method (Gue et al., 2020).

2.3 Phosphate solubilization

The qualitative test of phosphate solubilization was conducted as previously described and a sample of 10 µL from each culture was spotted on Pikovskaya's (PVK) agar medium containing tricalcium phosphate ($\text{Ca}_3[\text{PO}_4]_2$) (Vyas et al., 2007). All cultures were incubated at 28 °C for 3 days and the presence of clear halo indicates the phosphate solubilization.

Quantitative estimation of inorganic phosphate solubilization was tested in medium Pikovskaya phosphate liquid medium, according to the methodology described by Pikovskaya (1948). The amount of soluble phosphate released was determined according to the colorimetric method with

Commented [AB6]: ?? correct it

Vanadomolybdo-phosphoric acid as follows: a 1 ml of each sample was added to a 1 ml of the Vanadate-molybdate reagent, incubated for 1 hour and finally measured from OD to 405 nm; it was determined from a standard curve which represents the OD at 405 nm as a function of the known concentrations of solubrious phosphate (KH_2PO_4).

The total soluble phosphorus was calculated from the regression equation of standard curve;

$$P(\mu\text{g/ml}) = (y-0,1175)/0,0058. \quad \text{Where } y = \text{sample OD. (1)}$$

2.4. IAA production

The qualitative test of the root growth-promoting hormone auxin (indole-3-acetic acid) was realized on the medium YEM agar supplemented with 0,5 mg/ml of L-tryptophan. After, an incubation period of 3 days at 28 °C for, sterile filter paper washers impregnated with Salkowski reagent had deposited onto the Petri dish containing the culture. The appearance of colonies with a pink-red hue after a duration of 30-60 minutes in darkness indicates the production of IAA.

The levels of the hormone auxin produced by bacteria in culture filtrate was estimated as described by Sarwar *et al.* (1992). Approximately 2 mL of culture was centrifuged at 9,000 rpm for 10 min, and a 1 mL aliquot of the supernatant was mixed with 2 mL of Salkowski's reagent (150 mL concentrated H_2SO_4 , 250 mL distilled water, 7,5 mL 0.5M $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$) and was incubated for 30-60min in darkness at room temperature. IAA production was observed as the development of a pink-red color, and the absorbances of each sample was read at 540 nm using a spectrophotometer (ThermoScientific, USA) and the level of IAA ($\mu\text{g.ml}^{-1}$) calculated using the formula below and information generated by a standard curve (Gordon and Weber, 1951) made from a series of solutions with known auxin concentrations ranging from 0 to 120 ppm;

$$\text{IAA } (\mu \text{ g/ml}) = (y-0,16)/0,004; \text{ where } Y = \text{sample OD. (2)}$$

2.5. Siderophore production

All glass ware was soaked overnight in 6M HCl and rinsed with distilled water several times to remove any traces of iron. Siderophore production was tested qualitatively using Chrome Azurol S medium (CAS-medium) as described by Schwyn and Neilands. 10 μl of each isolate culture was spotted on the surface of CAS agar medium and incubated at 28 °C for 72 h. Siderophore production was assessed on the basis of change in color of the medium from green to yellow after incubation, three replicates were performed. Measuring the diameters of the halo and the Bacteria colony makes it possible to establish a report for each strain reflecting the quantity of siderophores secreted;

$$\text{SI} = \text{HD/CD}; \text{ where SI} = \text{siderophore production index}; \text{ HD} = \text{halo diameter}; \text{ CD} = \text{colony diameter.}$$

2.6. Statistical Analysis

All data obtained were analyzed using SAS 9.4 software. This data reported were means of at least three replicates. One-way analysis of variance and Student-Newman-Keuls test were used to compare the amount of soluble phosphate, the quantity of IAA and siderophore production index of

endophytes isolated from Bambara groundnut plants grown on different soils. Differences with level of 5 % were considered significant.


3. RESULTS

After being reactivated, the strains in our collection were examined for purity. Each strain displayed well-isolated colonies on the petri dishes that shared the same morphological traits, such as color, diameter, appearance, etc. This demonstrates the colonies' homogeneity and uniformity, which reflects the reactivated strain's purity. The reactivated strains were stored at +4° for the remainder of the tests.

3.1. Phosphate solubilization

The ability of isolates to solubilize inorganic phosphorus was determined firstly in Phikovskaya agar plate containing tricalcium phosphate (Ca_3PO_4) (Figure 1a). All 34 isolates were shown positive for phosphorus solubilization and form a clear halo. The quantitative estimation of the phosphorus solubilization by these strains in liquid medium showed an important variability between strains highlighted by the different shades of yellow expressing the variation in the concentrations of solubilized phosphate (Figure 1b). Indeed, the concentration of solubilized phosphate were very significantly different ($P=0,0001$) with value ranging from 16.43 to 453.45 $\mu\text{g/ml}$ (Table 1). Otherwise, most isolates had the capability to solubilize the high concentration of phosphate, thereby the strains RFK27 *Pseudomonas poae*, RFK11 *Bacillus pocheonensis*, RFK8 *Bacillus subtilis*, RFK12 *Bacillus cereus*, RFK29 *Lysobacter* and RFK10 *Bacillus subtilis*, with respectively, 453.45 $\mu\text{g/ml}$, 439.66 $\mu\text{g/ml}$, 387.94 $\mu\text{g/ml}$, 361.64 $\mu\text{g/ml}$, 316.63 $\mu\text{g/ml}$ and 313.37 $\mu\text{g/ml}$ were identified as most efficient P-solubilizer (Table 1).

Commented [AB7]: PROVIDE ACCESSION NUMBERS FOR ALL STRAINS



B

Figure 1a: qualitative test

Figure 1b: quantitative test

Figure 1: Phosphate solubilization test

3.2. IAA production

The ability of the bacterial isolates to produce IAA was detected by the development of pink-red color after the addition of Salkowski reagent to the culture, firstly in YEM-tryptophan agar plate (Figure 2a). All isolates were able to synthesize this phytohormone using tryptophan as precursor. The quantitative estimation of the IAA revealed an important variability between strains highlighted by the different shades of pink expressing the variation in the concentrations of IAA produced (Figure 2b). In fact, there was a highly significant difference ($P = 0,0001$) between the quantities of synthesized IAA, which ranged from 16.3 to 275.5 $\mu\text{g/ml}$. The highest production was obtained by RFK12 *Bacillus cereus*, RFK34 *Curtobacterium pusillum*, RKF10 *Bacillus subtilis* and RFK22 *Pseudomonas* spp producing, respectively, 275.5 $\mu\text{g/ml}$, 254 $\mu\text{g/ml}$, 246 $\mu\text{g/ml}$ and 212 $\mu\text{g/ml}$ (table 1).

Commented [AB8]: ACCESSION NUMBER TO BE MENTIONED FOR STRAINS




Figure 2a: qualitative test





Figure 2b: quantitative test

Figure 2: IAA production test

3.3. Siderophore production

The ability of isolates to produce siderophore was determined by the development of a yellow halo around the bacterial colony growing in CAS agar plate (Figure 3). All isolates were able to produce Siderophore except two isolates. However, a highly significant difference ($P = 0.0001$) was observed between the calculated Siderophore production index, which ranged from 1.07 to 5.36. The isolate that showed the best capability in Siderophore production was RFK27 *Pseudomonas poae*, followed by RFK26 *Pseudomonas fluorescens* and RFK22 *Pseudomonas* sp that produce 5.36, 4.3 and 3.65 respectively (Table 1).

- ← Colorless control
- ← Light pink
- ← No coloring
- ← Dark pink
- ← Pink coloring



← Yellow Colony

← Yellow halo

Figure 3: Siderophore production on chrome azurol S medium

UNDER PEER REVIEW

Table 1: Plant growth-promoting abilities of representatives of root-nodule bacteria isolated from ivoirian bambara groundnut

strains	closest relative species	Amount of P (µg/ml)	Amount of IAA (µg/ml)	Siderophore production index
RFK1	<i>Bacillus megaterium</i> strain FBMAX18	233.6±9.89 g	112.50±3.87 no	1.16±0.01 l
RFK2	<i>Bacillus</i> sp. L105	104.41±2.91 qp	152.5±2.50 h	1.07±0.00 l
RFK3	<i>Bacillus</i> sp. L105	255.71±21.16 g	23.50±1.5 xy	2.30±0.03ghijk
RFK4	<i>Bacillus safensis</i> strain BXC22	167.53±9.21 ml	19.58±4.28 xy	0
RFK5	<i>Bacillus nealsonii</i> strain ASB-160	251.10±1.77 f	16.3±5.62 y	2.00±0.05 ijk
RFK6	<i>Bacillus megaterium</i> strain YN7 16S	185.76±9.12 hjk	136.13±9.87 jk	3.5±0.03 cde
RFK7	<i>Bacillus</i> sp. BAB-3563	60.83±1.47 ut	26.00±3.25 x	3.22±0.1cdef
RFK8	<i>Bacillus subtilis</i> strain THt3-1	387.94±5.69 c	122.38±3.12 lm	2.50±0.06fghij
RFK9	<i>Bacillus cereus</i> strain THt1-8	73.95±1.64 st	51.25±7.75 v	2.85±0.08 defgh
RFK10	<i>Bacillus subtilis</i> strain V90	313.37±5.43 e	246.00±8.46 c	2.75±0.11 efghi
RFK11	<i>Bacillus pocheonensis</i> strain P12	439.66±3.45 b	129.8±3.12 kl	1.08±0.02 l
RFK12	<i>Bacillus cereus</i> strain BCRh6	361.64±4.12 d	275.5±5.51 a	2.84±0.33 defgh
RFK13	<i>Bacillus cereus</i> strain BAB-6399	48.97±7.67 u	181.88±4.63 f	2.51±0.04fghij
RFK14	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> strain GC17	51.29±5.20 u	104.42±1.66 op	2.49±0.02ghij
RFK15	<i>Bacillus tequilensis</i> strain RBB6	160.29±3.69 m	74.83±8.37 st	2.51±0.07fghij
RFK16	<i>Bacillus megaterium</i> strain VITNJ1	135.40±6.21 n	93.75±3.51 q	.67±0.05 kl
RFK17	<i>Bacillus megaterium</i> strain E2-04	178.45±5.18 jkl	141.75±6.75 ij	2.57±0.01fghi
RFK18	<i>Bacillus megaterium</i> strain DS8	115.95±3.27 nop	54.87±5.13 v	3.6±0.03 cd
RFK19	<i>Bacillus subtilis</i> strain LWIS15	299.07±14.14 e	154.75±2.75 h	3.61±0.06 cd
RFK20	<i>Bacillus subtilis</i> subsp. <i>subtilis</i> strain AN3	94.28±8.39 qr	116.25±7.06 mn	2.1±0.1 hijk
RFK21	<i>Pseudomonas</i> sp. strain Fas20	84.05±3.89 rs	145.63±3.13 h	3.52±0.03 cde
RFK22	<i>Pseudomonas</i> spstrain 7	212.77±4.13 h	212.00±4.76 d	2.46±0.05fghij
RFK23	<i>Pseudomonas azotoformans</i> strain R2SsM3P2C7	111.01±5.93 nop	39.67±1.15 w	3.65±0.04 c
RFK24	<i>Pseudomonas</i> sp. strain Fas20	67.73±8.99 stu	97.58±3.75 pq	3.05±0.07 cdefg
RFK25	<i>Pseudomonas koreensis</i> strain PS3TA2	132,155±11,22 n	79.25±8,01 rs	2.00±0.04 ijk
RFK26	<i>Pseudomonas fluorescens</i> strain FC6846	125.78±7.1 no	174.62±4.12 g	4.3±0.09 b
RFK27	<i>Pseudomonas poae</i> strain 15-A1	453.45±4.13 a	82.75±2.75 r	5.36±0.03 a
RFK28	<i>Lysobactersp.</i> KNUC361	161.58 ±12.68 m	21.00±3.5 xy	1.32±0.02 l
RFK29	<i>Lysobactersp.</i> KNUC361	316.63±3.97 e	69.5±3.0 ut	2.92±0.02cdefgh
RFK30	<i>Rahnellainusitata</i> strain FOD 9/21	229.70±4.10 g	196.12±4.13 e	2.3±0.01 fghij
RFK31	<i>Microbacteriumsp.</i> HBUM179633	16.47±6.64 v	85.50±8.75 r	3.00±0.06cdefg
RFK32	<i>Microbacteriumsp.</i> strain JL3592	81.72±2.90 rs	119.0±3.88 mn	0
RFK33	<i>Paenibacilluslautus</i> strain E118	49.23±9.41 u	103.87±2.62op	2.72±0.12 efghi
RFK34	<i>Curtobacteriumpusillum</i>	68.23±3.12 stu	254.0±4.50 b	2.77±0.14efghi
Average		183,57±10,22	118,51±5,94	2,44±0,09
CV		65,41	58,94	45,37
Pr>F		<.0001	<.0001	<.0001

Values indicate mean values (±SD); different letters indicate significant differences within a column at the 5 % level according to the Newman-Keuls test

4. Discussion

The context of this study is rooted in the growing need to promote environmentally friendly agricultural practices. In this context, the evaluation of the PGPR potential of endophytic bacteria associated with Bambara groundnut as a promising response to improve crop productivity while reducing the use of chemical inputs. Thus, this study is positioned at the heart of current issues by contributing to the search for sustainable solutions for agriculture

Commented [AB9]: Why choosing this crop particularly

The majority of nodule endophyte in this study had potential for promoting plant growth by the production of IAA, production of siderophore and solubilization of phosphate. The capacities of these strains would be linked to their intrinsic characteristics, in particular their genetic lineage. Indeed, several researches have revealed that bacteria with PGP traits were generally affiliated with the genera *Bacillus*, *Pseudomonas*, *Paenibacillus*, *Rahnella* etc. included within Firmicutes and Proteobacteriaphyla (Bahroun et al., 2018; Pérez-Montaña et al., 2014). It confirms the results reported by Kuklinsky-Sobral et al. (2004) which studied nodule endophyte isolated from soybean

Phosphorous is one of the major second nutrient only to nitrogen in requirement for plants. It plays an important role virtually in all major metabolic processes in plant including photosynthesis, energy transfer, signal transduction, macromolecular biosynthesis and respiration (Khan et al., 2010).

All strains were able to solubilize phosphate in quantities ranging from 16 to 453.45 µg/ml. However, the of six (06) strains (RFK27 *Pseudomonas poae*, RFK11 *Bacillus pocheonensis*, RFK8 *Bacillus subtilis*, RFK12 *Bacillus cereus*, RFK29 *Lysobacter*, RFK10 *Bacillus subtilis*) that solubilized high concentrations of phosphate (quantities > 300 µg/ml), four (04) were *Bacillus* species. These strains are able to solubilize phosphate because soil microorganisms are naturally occurring biological agents that can dissolve inorganic phosphate and make it available. Similar results were found by Ferchichi et al. (2019) who showed that better quantity of solubilized phosphate was 320.27 µg /ml and *Bacillus subtilis* was well documented for their phosphate solubilizing abilities (Ferchichi et al., 2019; Mohamed et al., 2018). According to recent research, some of the genera with PGPR traits that enable them to dissolve phosphorus include *Arthrobacter*, *Bacillus*, *Burkholderia*, *Enterobacter*, *Microbacterium*, *Pseudomonas*, *Rhizobium*, *Mesorhizobium*, *Flavobacterium*, and *Serratia* (Riaz et al., 2021).

Other researchers reported that the endophytic bacteria of cocoa nurseries (41,28 % of the collection) from various soils were able to solubilize phosphate with Solubilisation index indices ranging from 20% to 200% (Ouattara et al., 2019). The ability of some microorganisms to convert phosphate to form solution is available, such as orthophosphate, is an important PGPR trait to increase crop yields (Chen et al., 2006).

IAA is one of the most important phytohormone and function as important signal molecule in the regulation of plant development (Compant et al., 2010).

All strains tested produced a detectable quantity of IAA, which ranged from 16.3 to 275.5 µg/ml. Of the four (04) strains (RFK12 *Bacillus cereus*, RFK34 *Curtobacterium pusillum*, RFK10 *Bacillus subtilis*, RFK22 *Pseudomonas sp*) with the best IAA productions (quantities > 200 µg/ml), two (02) belong to the genus *Bacillus*.

Commented [AB10]: Were they identified as strains ? were accession numbers received for all isolates ?

The ability of the tested strains to produce IAA may be attributed to the substantial presence of IAA in the nodules of leguminous plants. Indeed, this phytohormone may be involved in the development of legume-rhizobia symbiosis, a phenomenon that is the foundation for the formation of nodules. (Duca *et al.*, 2014).

Kumar and *al.*(2012)found that the range of IAA production was 0.2 to 213 µg/ml. On the contrary, several studies demonstrated the ability of nodule endophyte to produce IAA with value ranging from 0.89 to 63.55 µg/ml (Saïdi *et al.*, 2013)and from 0.67 to 74.51µg/ml (Ferchichi *et al.*, 2019). It has been reported that IAA production by PGPR can vary among different species and strains (Compant *et al.*, 2010).Also, several authors reported that *Bacillus* species used as biofertilizers probably have direct effects on plant growth through the synthesis of plant growth hormones (Amer & Utkhede, 2007).

Iron is an essential growth element for all living organisms. The scarcity of bioavailable iron in soil habitats and on plants surfaces foments a furious competition (Whipps, 2001).

This work showed that the majority of Bambara groundnut nodule endophytes were able to produce siderophore. However, it has been observed that the first best producers of siderophore were RFK27 *Pseudomonas poae*, RFK26 *Pseudomonas fluorescens* and RFK22 *Pseudomonas sp.* This PGPR trait has been observed in several in a variety of bacteria. Indeed, Research indicates that some plant growth-promoting bacteria (PGPB) produce low-molecular-weight chemicals (400–1500 Da) that have the ability to extract iron from the soil (Shanmugaiah *et al.*, 2015) It has also been reported that Gram-negative bacteria that produce siderophores are *Enterobacter* and *Pseudomonas*. In contrast, only 2% of Gram-positive species, such as *Bacillus* and *Rhodococcus*, can do the same (Czarneset *al.*, 2020). Other authors reported also the production of siderophore by *pseudomonas* (Sasirekha and Shivakumar, 2016).

The strain RFK27 *Pseudomonas poae* which was the best producer of Siderophore was also the best efficient P-solubilizer. and produced a significant amount of IAA. The strain RFK12 *Bacillus cereus* that was better producer of IAA was also best fourth efficient P-solubilizer. Researchers such as Backer *et al.* (2021) and Ramakrishnan *et al.* (2024) noted the importance of PGPRs by saying that PGPR can promote plant growth through mobilization of nutrients in soils, production of plant growth regulators, protection of plants from phytopathogens by controlling or inhibiting them, improvement of soil structure and enhanced tolerance to biotic and abiotic stresses.

The endophytic bacteria in this study have the potential to be used in sustainable agriculture as biofertilizers. These bacterial strains can help reduce the dependence on synthetic chemicals, which contributes to the preservation of the environment.

5. Conclusion

The endophytes isolated from Bambara groundnut nodules in Daloa soil demonstrated overall great capacities to solubilize phosphate and produce IAA and siderophores. Nevertheless, these bacteria showed a disparity in their capacity to execute these three (03) PGPR functions, thereby exhibiting varying potential in promoting plant growth. In total, 19 strains present multi PGPR characters very interesting for the improvement of plant productivity.

Commented [AB11]: What is AIA ?

In perspective, it would be interesting to test the 19 strains with multi-PGPR traits under field conditions.

6. References

- Vance, C. P. (2001). Symbiotic nitrogen fixation and phosphorus acquisition. Plant nutrition in a world of declining renewable resources. *Plant physiology*, 127(2), 390-397. <https://doi.org/10.1104/pp.010331>
- Soussou, S. (2013). Adaptation of the Fabaceae-rhizobium symbiosis to mining sites: Zinc uptake by *Anthyllis vulneraria* and analysis of the diversity of symbiotic bacteria of *Hedysarum coronarium*. Doctoral thesis in Microbiology, Parasitology Sustainable Agriculture. Montpellier Supagro (France), 230 p.
- Hussain, I., Ali, I., Ullah, S., Iqbal, A., Tawaha, A. A., Al-Tawaha, A. R., & Sirajuddin, S. N. (2021). Agricultural soil reclamation and restoration of soil organic matter and nutrients via application of organic, inorganic and bio fertilization (Mini review). In IOP Conference Series: Earth and Environmental Science (Vol. 788, No. 1, p. 012165). IOP Publishing. DOI:10.1088/1755-1315/788/1/012165.
- Imran, A., Khan, A. A., Bari, A., Ud Din, R., Ali, R., Ahmed, N., & Khan, G. R. (2019 a). Production statistics and modern technology of maize cultivation in Khyber Pakhtunkhwa Pakistan. *Plant science. Archives*, 2, 1-12. DOI: <http://dx.doi.org/10.33680/plsa.2019.014>.
- Imran, A., Bari, A., & ALI, R. (2019 b). Peach organic sources, phosphorous and beneficial microbes application enhance productivity and profitability of soybean. *Soybean Research*, 1, 39. DOI:10.20944/preprints201911.0159.v1
- Khan, I. (2018). Climate change is a threat toward agronomy (base of food, fiber system), and food security. *Food & Nutrition. Journal*, 3(10), 1-8. DOI:10.29011/2575-7091.100060
- Gupta G., Parihar, S. S., Ahirwar, N. K., Snehi, S. K. & Singh, V. (2015). Plant Growth Promoting Rhizobacteria (PGPR): Current and Future Prospects for Development of Sustainable Agriculture. *Microbial Biochemical Technology*, 7: 096-102. doi:10.4172/1948-5948.1000188.
- Ferchichi, N., Toukabri, W., Boularess, M., Smaoui, A., Mhamdi, R., Trabelsi, D., (2019). Isolation, identification and plant growth promotion ability of endophytic bacteria associated with lupine root nodule grown in Tunisian soil. *Arch Microbiol* DOI: 10.1007/s00203-019-01702-3
- Pérez-Montaña, F., Alías-Villegas, C., Bellogín, R.A. et al. (2014). Plant growth promotion in cereal and leguminous agricultural important plants: from microorganism capacities to crop production. *Microbiol Res* 169:325–336. <https://doi.org/10.1016/j.micres.2013.09.011>

- Zaatri, T., & Tagzirt, L. (2023). Study of Ceratonia siliqua mycorrhizae in the Tizi-ouzou region (Doctoral dissertation, Mouloud Mammeri University).
- Raïssa, G.N K., TAHA K., KOFFI A.G., GUEI A. M., KOFFI M., FILALI-MALTOUF A., & KONATE I. (2024). Morphological and Molecular Characterization of Endophytic Bacteria Isolated from Bambara Groundnut (*Vigna subterranea*) Nodules in Daloa, Côte d'Ivoire. *Biotechnology Journal International*. 28 (6):140-50. <https://doi.org/10.9734/bji/2024/v28i6757>
- Guei, N.K.R, Konate, I., Taha, K., Bamba, I., Beugre, G. A. M., Akaffou, D. S., & Filali-Maltouf, A. (2020). Molecular and Symbiotic Efficiency Characterization of Rhizobia Nodulating Bambara Groundnut (*Vigna subterranea* L.) from Agricultural Soils of Daloa Localities in Ivory Coast. *International Journal of Current Microbiology and Applied Sciences*. 9(07): 507-519. <https://doi.org/10.20546/ijcmas.2020.907.056>
- Vyas, P., Rahi, P., Chauhan, A., & Gulati, A. (2007). Phosphate solubilization potential and stress tolerance of *Eupenicillium parvum* from tea soil. *Mycol. Res.* 111, 931–938. DOI:10.1016/j.mycres.2007.06.003
- Pikovskaya, R.I. (1948). Mobilization of phosphorus in soil in connection with vital activity of some microbial species, *Microbiologia*, 1948, 17 edition. DOI:10.4236/ojss.2015.55010
- Sarwar, M., Muhammad, A., Dean, A. M., & Frankenberger, W. T. (1992). Tryptophan-Dependent Biosynthesis of Auxins in Soil. *Plant and Soil* 147 (2): 207-15. <https://doi.org/10.1007/BF00029072>.
- Gordon, S. A., & Weber, P. R. (1951). "Colorimetric estimation of indole acetic acid". *Plant Physiology* 26 (1): 192-95. <https://doi.org/10.1104/pp.26.1.192>
- Schwyn, B., & Neilands, J. B. (1987). Universal chemical assay for the detection and determination of siderophores; *Analytical Biochemistry*; 160(1); 47-56. DOI:10.1016/0003-2697(87)90612-9
- Bahroun, A., Jousset, A., Mhamdi, R., et al, (2018). Anti-fungal activity of bacterial endophytes associated with vegetables against *Fusarium solani*: assessment of fungi soil suppressiveness and plant protection induction. *Appl Soil Ecol* 124:131–140. <https://doi.org/10.1016/j.apsoil.2017.10.025>
- Pérez-Montaña, F., Alías-Villegas, C., Bellogín, R.A., et al, (2014). Plant growth promotion in cereal and leguminous agricultural important plants: from microorganism capacities to crop production. *Microbiol Res* 169:325–336.
- Kuklinsky-Sobral, A.J., Mendes, W.L., Geraldi, R., Pizzirani-Kleiner, I.O., & Azevedo, A.J.L. (2004). Isolation and characterization of soybean-associated bacteria and their potential for plant growth

promotion. *Environnement Microbiology* 6, 1244-1251. DOI : 10.1111/j.1462-2920.2004.00658.x

Khan, M.S., Zaidi, A., Ahemad, M., Oves, & Wani, A.P. (2010) : Plant growth promotion by phosphate solubilizing fungi current perspective, *Archives of Agronomy and Soil Science*, 56 :1, 73-<https://doi.org/10.1080/03650340902806469>

Ferchichi, N., Toukabri, W., Boularess, M., Smaoui, A., Mhamdi, R., & Trabelsi, D. (2019). Isolation, identification and plant growth promotion ability of endophytic bacteria associated with lupine root nodule grown in Tunisian soil. *Arch Microbiol* DOI: 10.1007/s00203-019-01702-3

Mohamed, E.A.H., Farag, A.G. & Youssef, S.A. (2018) Phosphate Solubilization by *Bacillus subtilis* and *Serratia marcescens* isolated from Tomato Plant Rhizosphere. *Journal of Environmental Protection*, 9, 266-277. DOI :10.4236/jep.2018.93018

Riaz, U., Murtaza, G., Anum, W., Samreen, T., Sarfraz, M., & Nazir, M. Z. (2021). Plant growth-promoting rhizobacteria (PGPR) as biofertilizers and biopesticides. *Microbiota and biofertilizers: a sustainable continuum for plant and soil health*, 181-196. DOI:10.1007/978-3-030-48771-3_11

Ouattara, A., Coulibaly, K., Konate, I., Kebe, I.B., Tidou A.S., Filali-Maltouf, A. (2019). Selection of Cocoa Tree (*Theobroma cacao* Linn) Endophytic Bacteria Solubilizing Tri-Calcium Phosphate, Isolated from Seedlings Grown on Soils of Six Producing Regions of Côte d'Ivoire. *Advances in Microbiology*, 9 (9), pp 842-852. DOI : 10.4236/aim.2019.99051

Chen, Y.P., Rekha, P.D., Arun, A.B., Shen, F.T., Lai, W.A. & Young, C.C. (2006). Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Applied Soil Ecology*; 34 (Suppl 1): 33-41. DOI : 10.1016/j.apsoil.2005.12.002

Compant, S., Clément, C., & Sessitsch A. (2010). Plant growth-promoting bacteria in the rhizo- and endosphere of plants : their role, colonization, mechanisms involved and prospects for utilization. *Soil Biol. Biochem.* 42, 669–678. <https://doi.org/10.1016/j.soilbio.2009.11.024>

Duca, D., Lorv, J., Patten, C. L., Rose, D., & Glick, B. R. (2014). Indole-3-acetic acid in plant-microbe interactions. *Antonie Van Leeuwenhoek*, 106, 85-125. <https://doi.org/10.1007/s10482-013-0095-y>

Kumar, A., Kumar, A., Devi, S., Patil, S., Payal, C., & Negi, S. (2012). Isolation, screening and characterization of bacteria from Rhizospheric soils for different plant growth promotion (PGP) activities: an in vitro study. *Recent research in science and technology*, 4(1), 1-5.

- Saïdi, S., Chebil, S., Gtari, M., & Mhamdi, R. (2013). Characterization of root-nodule bacteria isolated from *Vicia faba* and selection of plant growth promoting isolates. *World J Microbiol Biotechnol* 29 :1099–1106. DOI 10.1007/s11274-013-1278-4
- Amer A. G., & Utkhede R. S (2007). Development of formulation of biological agents for management of root rot of lettuce and cucumber. *Canada J. Microbiol.* 46:809-816. <http://dx.doi.org/10.1139/w00-063>
- Whipps, J.M. (2001). Microbial interactions and biocontrol in the rhizosphere. *J. Experi Bot.* 52(1) :487-511. http://dx.doi.org/10.1093/jexbot/52.suppl_1.487.
- Shanmugaiah, V., Nithya, K., Hari Krishnan, H., Jayaprakashvel, M., & Balasubramanian, N. (2015). Biocontrol mechanisms of siderophores against bacterial plant pathogens. Sustainable approaches to controlling plant pathogenic bacteria, 167-190. DOI:10.1201/b18892-9
- Czarnes, S., Mercier, P. E., Lemoine, D. G., Hamzaoui, J., & Legendre, L. (2020). Impact of soil water content on maize responses to the plant growth-promoting rhizobacterium *Azospirillum lipoferum* CRT1. *Journal of Agronomy and Crop Science*, 206(5), 505-516. <https://doi.org/10.1111/jac.12399>
- Sasirekha, B., & Shivakumar, S. (2016). Siderophore production by *Pseudomonas aeruginosa* FP6, a biocontrol strain for *Rhizoctonia solani* and *Colletotrichum gloeosporioides* causing diseases in chilli. *Agriculture and Natural Resources* <https://doi.org/10.1016/j.anres.2016.02.003>
- Backer, R., Rokem, J. S., Ilangumaran, G., Lamont, J., Praslickova, D., Ricci, E., & Smith, D. L. (2018). Plant growth-promoting rhizobacteria: context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Frontiers in plant science*, 9, 1473. DOI: 10.3389/fpls.2018.01473
- Ramakrishnan, P., Rangasamy, A., Ariyan, M., Raghu, R., Krishnamoorthy, R., Senthil Kumar, M., & Vincent, S. (2024). Rhizobial and passenger endophytes alleviate moisture stress in groundnut (*Arachis hypogaea*). *Plant Stress*, 14, 100590. <https://doi.org/10.1016/j.stress.2024.100590>