

ASSESSMENT OF ANTIMICROBIAL ACTIVITY OF *OCIMUM GRATISSIMUM* AND *CYMBOPOGON FLEXUOSUS* OIL AGAINST SOME SELECTED PATHOGENIC MICROORGANISM

ABSTRACT

Essential oils are concentrated substances extracted from plant components such as leaves, flowers, stems, roots, and bark. These oils contain volatile organic compounds that are responsible for their distinctive aroma and therapeutic effects. This study investigated the antimicrobial activity of essential oils from *Ocimum gratissimum* and *Cymbopogon flexuosus* against two gram-positive bacteria, two gram-negative bacteria, and one fungus, using the agar well diffusion method. Fourier-transform infrared spectroscopy (FTIR) analysis of selected essential oils revealed the presence of functional groups, such as aldehydes, ketones, and aromatic rings. The results showed that both oils exhibited significant antimicrobial activity against all tested microorganisms, with *Cymbopogon flexuosus* being more effective, completely inhibiting the growth of *Bacillus paramycoides* and *Bacillus altitudinus* at concentrations up to 40%. The zone of inhibition for *Cymbopogon flexuosus* against *Escherichia coli* and *Pseudomonas spp. RRC15* were 25.5 mm and 8.1 mm at 100% concentration, respectively. *Ocimum gratissimum L.* showed lower antimicrobial activity than *Cymbopogon flexuosus*, with the zone of inhibition in the decreasing order of *Bacillus paramycoides*, *Bacillus altitudinus*, *Escherichia coli*, and *Pseudomonas spp. RRC15*. Both essential oils completely inhibited the growth of *Colletotrichum gloeosporioides* at concentrations ranging from 100% to 40%. The antimicrobial activity was attributed to the presence of bioactive compounds, such as citral, eugenol, and other monoterpenes in the essential oils, which disrupt bacterial and fungal cell membranes, leading to cell death. These findings highlight the potential of *Ocimum gratissimum L.* and *Cymbopogon flexuosus* essential oils as natural antimicrobial agents for various applications.

Key words:

Antimicrobial activity, *Ocimum gratissimum*, *Cymbopogon flexuosus*, Essential oils, Fourier-transform infrared spectroscopy, Agar well diffusion method, Bioactive compounds, Monoterpenes

Commented [MG1]:
Include fruits

Commented [MG2]: An example of these applications?

Commented [MG3]: The summary is well centered, objective and in accordance with what was worked on in your manuscript.

Commented [MG4]: Are aligned with the theme of the manuscript

INTRODUCTION

Plant-derived essential oils are a significant product in agriculture-based industries. They are commonly utilised as flavouring agents in a variety of products including food, beverages, perfumes, pharmaceuticals, and cosmetics (1). At least 2000 plant species have been used to produce 3000 different types of essential oils with numerous benefits and out of these 300 oils are important from a commercial perspective (2). Primarily obtained through steam distillation, hydrodistillation, and solvent extraction, these substances are stored in plant oil ducts, resin ducts, glands, or trichomes. They contain 20-100 plant metabolites and are chemically complex mixtures mainly of low-molecular-weight compounds such as terpenoids and phenylpropanoids, along with some aromatic and aliphatic constituents (3). Monoterpenes, sesquiterpenes, and their oxygenated derivatives form the largest chemical groups in essential oils (4,5). These natural compounds exhibit various beneficial properties, including antimicrobial, antiviral, antimutagenic, anticancer, antioxidant, anti-inflammatory, immunomodulatory, and antiprotozoal activities. (6–9).

Phytopathogenic microorganisms reduce plant yield and compromise fruit quality through postharvest diseases, leading to deterioration of produce resulting in food wastage (10–14). Chemical agents are harmful and is detrimental to human health, also causes environmental contamination, resulting in increase in production costs, and promote pathogen resistance (15,16). As a result, many studies have explored plant extracts and essential oils in combating this.

Essential oils from the *Cymbopogon* genus (*Poaceae* family) are valuable in fragrance, flavouring, perfumery, and pharmaceuticals (17). This genus comprises around 140 species globally, with 45 found in India. Economically important oils of species include lemongrass, citronella, ginger grass, and rusa (18). There is subtle variations in oil content, composition, and quality, morphological differences within and between species (19). Lemongrass (*C. flexuosus*) is widely cultivated in Brazil, Mexico, Dominica, Haiti, Indonesia, and China, and in Indian states like Kerala, Assam, Maharashtra, and Uttar Pradesh. The essential oils from *Cymbopogon* species are rich in terpenoids, such as geraniol, citronellol, and citronellal. Citral, another terpenoid, is used in synthesizing ionone and vitamin A (20). *Cymbopogon flexuosus*, a tall perennial grass reaching 1.5m with a lemony scent and dark-green foliage, thrives in tropical and subtropical climates at temperatures between 10°C and 33°C, requiring abundant sunlight for oil production. It is sensitive to cold and cannot withstand frost. Oil is extracted

from fresh plant materials, mainly stalks and leaves, through hydrodistillation (21). Studies show lemongrass extract, oil, citral, and citral-derived compounds have antimicrobial, allelopathic, anthelmintic, anti-inflammatory, anticancer, and antioxidant properties, and can repel insects and mosquitoes.

Ocimum gratissimum L. (Lamiaceae), native to Africa, Asia, and South America, is part of a genus with about 30 species in tropical regions, producing essential oils for perfumery, cosmetics, pharmaceuticals, and food industries (22,23). Also known as clove basil, African basil, or wild basil, *O. gratissimum* can grow up to 3 m tall with a woody stem base, ovate leaves 5–13 cm long and 3–9 cm wide, and petioles 1–6 cm long. Its flowers form in inflorescences 5–30 cm long, either simple or branched (Matasyoh et al., 2007). The plant is used for food flavoring and medicinal purposes due to its anti-inflammatory, analgesic, hepatoprotective, antimutagenic, antihypertensive, and anticarcinogenic properties(22,23). It exhibits antinociceptive, antagonistic, antibacterial, and antifungal effects on intestinal motility (24,25). This study assessed the in vitro antimicrobial efficacy of *Cymbopogon flexuosus* and *O. gratissimum* L. essential oils against bacterial and fungal species.

MATERIALS AND METHODS

Collection of materials

The bacterial strains *Escherichia coli*, *Pseudomonas*, *Bacillus altitudinus*, and *Bacillus paramycoides* were obtained from the Department of Microbiology, College of Basic Sciences and Humanities, GBPUA&T, Pantnagar. The fungal strain was isolated from anthracnose-affected mango fruits at the Department of Plant Pathology, College of Agriculture, GBPUA&T, Pantnagar.

The essential oils utilised in the antimicrobial assays, *Ocimum gratissimum* and *Cymbopogon flexuosus*, were procured from Aromaaz International, Sahibabad, Ghaziabad, Uttar Pradesh.

Fourier transform infrared spectroscopy

The essential oils were analysed utilising a Fourier-transform infrared spectroscopy phase II spectrometer. FTIR spectra were obtained using Bruker software and subsequently analysed to determine the functional groups present in the essential oil (26). The IR spectra for the range of (600–3900 cm^{-1}) were employed. This range corresponds to the mid-infrared region, which exhibits specific vibrational bands associated with the functional groups. These groups facilitate the determination of the sample composition based on this mid-IR spectral fingerprint.

Commented [MG5]: The introduction presents relevant information about essential oils derived from plants, highlighting their applications, extraction methods, chemical composition and bioactive properties. However, the structure is dense, with abrupt transitions between topics and an excess of technical details that could be condensed to improve fluidity. There is a lack of a clear justification for the study and a more explicit connection between *Cymbopogon* and *Ocimum gratissimum* oils and their relevance in combating phytopathogens. Furthermore, there are grammatical and agreement problems that compromise the quality of the text. It is recommended to reorganize and summarize the introduction, highlighting the objective of the study in a clearer and more concise way.

Preparation of essential oil concentration: Various concentrations of essential oils, ranging from 10% to 100%, were prepared utilising DMSO as a solvent.

Antimicrobial assay

Antibacterial

An agar well diffusion assay was employed to determine the antimicrobial activity of the essential oils (27). The bacterial culture broth of 24 h old was prepared in a nutrient broth medium. Nutrient agar was aseptically poured into Petri plates and allowed to solidify. Wells were created using a sterile borer of 5 mm diameter. The spread plate method was utilised to transfer the inoculum to the plate. The bacterial culture (100 µL) was transferred onto a Petri plate and spread using a sterile L spreader. Subsequently, 20 µL of the essential oil was transferred into each well using a sterile micropipette. The plates were incubated at 37°C for 24 h. Streptomycin and tetracycline were used as positive controls.

Antifungal

Fungal spores from anthracnose-affected mango were purified through repeated culturing in PDA media. One-week-old fully grown fungal mycelium without contamination was utilised for the present study. Potato Dextrose Agar media was aseptically poured into Petri plates, and wells were created using a sterile borer of 5 mm diameter. Mycelium from the fully grown plate was excised using a sterile borer and subsequently transferred with a sterile loop to the centre of another plate containing a bored well. The bored wells were filled with 20 µL of essential oil using a micropipette. Carbendazim (0.2% concentration) served as a positive control.


RESULTS AND DISCUSSION

FTIR analysis of essential oils

Figure 1 illustrates the FTIR spectra of the essential oil of *Cymbopogon flexuosus*. Examining from higher wavelengths to lower wavelengths, the spectral graph exhibits peaks in the regions of 1743.61 and 1682.32, indicating the presence of C=O (aldehyde/ketone group). Additionally, the presence of peaks in the regions of 3.48.76, 1561.45, and 1513.88 suggests the presence of aromatic C=C stretching, while a peak at 1461.09 represents C-H asymmetrical bend and C-H bend methylene. The peak in the region of 1322.16 indicates the presence of an OH bond. The region from 1322.16 to 728.32 represents the skeletal vibration C-C. The peak at 728.32 demonstrates 'methylene rocking'. The essential oil is known to comprise the main components,

Commented [MG6]: The "Materials and Methods" section clearly describes the materials used, such as bacterial and fungal strains and essential oils, as well as the methods, including FTIR and antimicrobial assays. However, negative controls and justifications for the chosen concentrations (10%-100%) are lacking. The agar diffusion technique does not mention how the inhibition halos were measured and analyzed, and the scientific names, such as *Pseudomonas*, need to be corrected. Furthermore, the description of the FTIR methodology and the antifungal assay could be more detailed. Greater organization with clear subtopics is also suggested to improve readability and replicability.

Additionally, a peak in wavelength is observed in the regions of 1643.77 and 1575.91, indicating the presence of the C=C conjugate. Eugenol constitutes the major component present in the essential oil of *Ocimum gratissimum*. The chemical structure reveals that it is a cyclohexane with hydroxyl and ether functional groups.



Eugenol




Fig 3 FTIR spectra of *Ocimum gratissimum*

Antimicrobial activity of essential oils

Antibacterial activity

Table 1 demonstrates the effects of varying concentrations of essential oils on their antimicrobial activity. The results indicated that for all selected bacterial and fungal species, the concentration of the essential oil was directly proportional to the zone of inhibition. At concentrations up to 40%, *Cymbopogon flexuosus* essential oil completely inhibited the growth of *Bacillus altitudinus*, and at concentrations up to 20%, it inhibited *Bacillus paramycoides*. The zone of inhibition against *Pseudomonas spp* was observed to be 8.1 mm for 100 percent concentration of oil and 5.25 for 40 percent concentration. Similarly, against *Escherichia coli*, the zone of inhibition was observed to be 25.5 mm at 100 percent EO concentration and 8.5 mm at 40 percent concentration.

EO *Cymbopogon flexuosus* and its antimicrobial activity

The EO of *C. flexuosus*, commonly known as lemongrass, contains an active component, citral, geranial, neral, and other monoterpenes which are responsible for its antibacterial activity. In a study it was highlighted that citral and other constituents of lemongrass oil possess significant antibacterial properties, particularly against Gram-positive bacteria such as *Staphylococcus aureus* and *Bacillus subtilis* (28). A researcher emphasised lemongrass oil's health benefits, particularly its antimicrobial properties due to its high citral content (29). These results suggest that the EO of *C. flexuosus* is more effective against gram-positive organisms due to their simpler cell wall structure (30).

Mode of action of Citral a compound present in *Cymbopogon flexuosus* against Bacteria

The antibacterial action of citral against gram-negative bacteria disrupts their lipopolysaccharide-rich outer membrane, increasing permeability and leading to cell lysis, altered ATP levels, membrane hyperpolarisation, and reduced cytoplasmic pH, indicating compromised integrity (31–33). This results in leakage of essential components and cell death. Citral induces oxidative stress, generating ROS that damages DNA, proteins, and lipids, thereby enhancing antimicrobial efficacy (33,34). It disrupts metabolic pathways, inhibits cell wall synthesis, impairs homeostasis, and increases the susceptibility to other antimicrobials (35). Citral also inhibits quorum sensing in bacteria, such as '*Pseudomonas aeruginosa*' and '*Cronobacter sakazakii*', reducing pathogenicity and enhancing immune response vulnerability (33). In Gram-positive bacteria, citral disrupts the cytoplasmic membrane lipid bilayer,

increasing permeability and causing intracellular leakage and cell death (34,36,37). It induces morphological changes such as membrane deformation and porosity (38,39). Citral inhibits key metabolic processes, including mitochondrial complex IV, generates ROS, and causes oxidative stress that damages DNA and proteins (40,41). It also downregulates ergosterol biosynthesis, contributing to its antifungal activity (31,42). Furthermore, citral enhances the efficacy of other antimicrobial agents, particularly when combined with nanoparticles, thereby offering effective treatments against resistant bacterial strains (43). This synergistic effect highlights the potential of citral as a natural preservative and therapeutic agent against bacterial infections.

EO *Ocimum gratissimum* and its antimicrobial activity

The EO *Ocimum gratissimum* exhibited a zone of inhibition against these bacterial species in decreasing order as *Bacillus paramycooides*, *Bacillus altitudinus*, *E. coli*, and *Pseudomonas spp. RRC15*. *Ocimum gratissimum* is rich in various phytochemicals, including eugenol, thymol, and other phenolic compounds (44,45) found that essential oils possess notable antimicrobial properties, particularly against gram-positive bacteria such as *Staphylococcus aureus* and *Bacillus subtilis*. This is consistent with the findings of (46) who noted that *Ocimum gratissimum* exhibited strong antibacterial activity against common oral pathogens, highlighting its potential for dental applications. The effectiveness of *Ocimum gratissimum* essential oil against gram-negative bacteria is low. This can be attributed to the structural differences between Gram-positive and Gram-negative bacteria, particularly the presence of an outer membrane in Gram-negative bacteria that acts as a barrier to the penetration of hydrophobic compounds. For example, a study by (45) indicated that while eugenol and methyl eugenol showed some antimicrobial activity, their effectiveness against gram-negative bacteria was comparatively weaker owing to their chemical structure.

Mode of action of eugenol a compound present in *Ocimum gratissimum* against Bacteria

Ocimum gratissimum essential oil disrupts bacterial cell membranes, resulting in the leakage of intracellular components and subsequent cell death, as demonstrated by (47). The phytochemical composition of the oil, which includes flavonoids and tannins, enhances its antibacterial activity by targeting multiple bacterial pathways (44,48). Eugenol, a phenolic compound present in oil, disrupts gram-positive bacterial cell membranes, interferes with metabolism, and modulates biofilm formation. It integrates into lipid bilayers, increases membrane fluidity, and induces lysis, leading to cellular leakage in *Staphylococcus aureus* and

Streptococcus pneumoniae (49). High concentrations of eugenol significantly inhibit gram-positive bacteria (50). Eugenol forms hydrogen bonds with proteins and inhibits bacterial metabolic enzymes (51), which is crucial for the treatment of antibiotic-resistant infections (52). It disrupts biofilm formation and increases the bacterial susceptibility to antimicrobials (52). In gram-negative bacteria, eugenol penetrates lipid bilayers and lipopolysaccharides of *Escherichia coli* and *Pseudomonas aeruginosa*, causing cell lysis and leakage at high concentrations (53–55). Eugenol inhibits membrane-bound ATPases and reduces ATP levels and viability of gram-negative bacteria (53). The hydroxyl group disrupts metabolic pathways by forming hydrogen bonds with proteins (51). Eugenol disrupts biofilm formation and enhances the susceptibility to antimicrobial agents (45,56). When combined with other antimicrobials, eugenol exhibits synergistic effects, enhancing activity against resistant strains and reducing antibiotic dosages (55,57).

Compared to *Cymbopogon flexuosus*, *Ocimum gratissimum* exhibits a lower zone of inhibition for bacterial species. This can be attributed to the difference in chemical composition, and the high concentration of active compounds, such as citral, present in *Cymbopogon flexuosus* oil, which is highly effective in disrupting the bacterial cell structure.

Antifungal activity

Table 1 depicts the antifungal activity of the essential oils *Cymbopogon flexuosus* and *Ocimum gratissimum* against *Colletotrichum gloeosporioides*. Essential oils at concentrations of 100%, 80%, 60%, and 40 percent had completely inhibited the growth of fungus and exhibited a total zone of inhibition without any mycelium growth. *Cymbopogon flexuosus* essential oil is rich in bioactive compounds, particularly citral, which is known for its significant antifungal activity against various fungal pathogens. A researcher (58) demonstrated that the essential oil of *Cymbopogon flexuosus* possesses potent antifungal properties against *Aspergillus* species, which are common postharvest pathogens. In a study (59,60) it is reported that lemongrass essential oil effectively inhibits the growth of key postharvest pathogens, including *Penicillium digitatum* and *Aspergillus niger*. Researchers findings suggested that antifungal activity is concentration-dependent, with higher concentrations leading to greater inhibition of fungal spore germination and growth (59,60).

In a study by researchers (61,62) it is found that lemongrass oil exhibits significant antifungal activity against various *Candida* species, including *Candida albicans*. This study indicated that the antifungal effects were closely associated with the presence of citral in the essential oil,

which demonstrated a strong capacity to inhibit fungal growth (62). This aligns with the findings of (63) who reported that lemongrass oil in both powder and oil forms effectively inhibits *Candida albicans*, further emphasising its potential as a natural antifungal agent.

Mode of action of Citral a compound present in *Cymbopogon flexuosus* against fungus

Citral exerts antifungal effects by disrupting the fungal cell membrane, increasing its permeability, and causing intracellular leakage. This disruption results in morphological changes such as thinner filaments and fragile hyphae, leading to cell death (64,65). The ability of citral to form charge transfer complexes with membrane components such as tryptophan also contributes to its efficacy (64). Additionally, citral inhibits ergosterol biosynthesis, a key component of fungal cell membranes, thereby compromising membrane integrity and function (31,66). This inhibition enhances the susceptibility of fungi to other antifungal agents, particularly azole-resistant strains (31). Citral also inhibits spore germination and proliferation by interfering with essential metabolic pathways and induces oxidative stress through reactive oxygen species (ROS) generation, damaging cellular components and leading to cell death (64,65,67). Furthermore, citral can enhance the efficacy of other antifungal agents by working synergistically with conventional drugs to combat resistant strains (68).

In a study (69) it is demonstrated that the essential oil of *Ocimum gratissimum* is highly effective against *Colletotrichum gloeosporioides*, a pathogen responsible for postharvest anthracnose in mangoes. This study reported that the oil exhibited fungicidal effects on both mycelial growth and spore germination. (70) reported that hydrodistilled volatile oils from the leaves of *Ocimum gratissimum* demonstrated significant antifungal activity against *Candida albicans*, a common pathogenic fungus. This was further supported by (46), who noted that *Ocimum gratissimum* showed antifungal activity against various *Candida* species, reinforcing its therapeutic relevance.

Mode of action of eugenol a compound present in *Ocimum gratissimum* against fungus

The antifungal mechanism of *O. gratissimum* likely involves disrupting the fungal cell membrane, leading to leakage of intracellular components and cell death. This is attributed to eugenol, a major component with strong antifungal properties (71)

The compounds present in *O. gratissimum* disrupts membrane integrity and metabolic processes (72,73). Its lipophilic nature allows it to penetrate fungal lipid bilayers, increase permeability, and cause cell lysis, effectively inhibiting fungi, such as *Candida albicans*,

Trichophyton rubrum, and *Microsporum canis* (71,74). *O. gratissimum* essential oil, with eugenol as a significant constituent, demonstrates potent antifungal activity with an MICs of 31.25 µg/mL against various fungi (75). Eugenol reduces fungal viability by inhibiting key enzymes that are essential for growth and reproduction (72). It forms hydrogen bonds with proteins and enzymes, disrupting critical biochemical processes within the fungal cells (46). Furthermore, eugenol modulates biofilm formation, rendering biofilm-associated fungi more susceptible to antifungal agents (76,77), which is crucial in clinical settings for challenging biofilm-associated infections. *O. gratissimum* essential oil also exhibits synergistic effects with other antifungal agents, enhancing activity, reducing the required dosages, and minimising potential side effects and resistance development (78).

Commented [MG7]: The section is well structured and presents a detailed analysis of the antimicrobial and antifungal activities of *Cymbopogon flexuosus* and *Ocimum gratissimum* essential oils. The description of the FTIR spectra is clear and provides a good correlation between the identified functional groups and the biological properties of the essential oils. The discussion of antimicrobial and antifungal activities effectively highlights key compounds, such as citral and eugenol, and their mechanisms of action, such as disruption of bacterial and fungal cell membranes. The relationship between chemical composition and biological activity is well established, supported by references to previous studies, which strengthens the credibility of the results. However, it would be interesting to include a critical analysis on possible limitations or challenges of using these essential oils in practical contexts, such as in the pharmaceutical or food industry. Furthermore, small improvements can be made to ensure greater fluidity between sections and uniformity in citations. Overall, the section provides a well-researched analysis and contributes significantly to the understanding of the antimicrobial and antifungal properties of these essential oils.

Table 1 Zone of inhibition against *Escherichia coli*, *Pseudomonas spp.* RRC15, *Bacillus altitudinus*, *Bacillus paramycooides*, and *Colletotrichum gloeosporoides*

Essential oil	Conc.	<i>Escherichia coli</i>	<i>Pseudomonas spp.</i> RRC15	<i>Bacillus altitudinus</i>	<i>Bacillus paramycooides</i>	<i>Colletotrichum gloeosporoides</i>
<i>Cymbopogon flexuosus</i>	100	25.5	8.1	100.0 (CI)	100.0 (CI)	100.0 (CI)
	80	35.0	5.25	100.0 (CI)	100.0 (CI)	100.0 (CI)
	60	19.5	6.75	100.0 (CI)	100.0 (CI)	100.0 (CI)
	40	8.5	5.25	100.0 (CI)	100.0 (CI)	100.0 (CI)
	30	-	-	37.5	100.0 (CI)	100.0 (CI)
	20	-	-	23.5	100.0 (CI)	100.0 (CI)
	10	-	-	20.5	24.0	100.0 (CI)
<i>Ocimum gratissimum</i>	100	6.5	4.25	17.5	37.5	100.0 (CI)
	80	5.25	2.25	11.5	14.5	100.0 (CI)
	60	4.25	1.25	9.5	16.0	100.0 (CI)
	40	3.75	1.0	9.0	7.5	100.0 (CI)
DMSO*	NA	0	0	0	0	0
Tetracyclin (disc)	10mcg	21.0	4.0	34.5	22.75	NA
Streptomycin	0.25mg/ml	1.0	1.0	18.0	17.0	NA
Carbendazim	0.2%	NA	NA	NA	NA	24.2

*DMSO - Dimethyl sulfoxide, NA-Not applicable

CONCLUSION AND SUMMARY

This study investigated the antimicrobial activity of essential oils from *Ocimum gratissimum* and *Cymbopogon flexuosus* against two gram-positive bacteria, two gram-negative bacteria, and one fungus, using the agar well diffusion method. FTIR analysis revealed the presence of functional groups, such as aldehydes, ketones, and aromatic rings, in the essential oils. Both oils exhibited significant antimicrobial activity against all tested microorganisms, with *Cymbopogon flexuosus* being more effective, completely inhibiting the growth of *Bacillus paramycooides* and *Bacillus altitudinus* at concentrations up to 40%. *Ocimum gratissimum* showed lower antimicrobial activity than *Cymbopogon flexuosus*, with the zone of inhibition decreasing in the order of *Bacillus paramycooides*, *Bacillus altitudinus*, *Escherichia coli*, and *Pseudomonas spp.* Both essential oils completely inhibited the growth of *Colletotrichum gloeosporioides* at concentrations ranging from 40% to 100%. The antimicrobial activity was attributed to the presence of bioactive compounds, such as citral, eugenol, and other

monoterpenes in the essential oils, which disrupt bacterial and fungal cell membranes, leading to cell death.

The potential applications of this antimicrobial agent extend across various industries and offer promising solutions to address microbial challenges. In the food industry, it can serve as an effective natural preservative, contributing to the extension of shelf life for perishable products and the reduction of food waste. This application aligns with increasing consumer demand for clean-label products and natural food additives. In agriculture, the antimicrobial properties of compounds could contribute to sustainable crop management strategies, potentially reducing reliance on synthetic pesticides and promoting environmentally friendly farming practices. Furthermore, the pharmaceutical and textile industries stand to benefit significantly from this antimicrobial agent. In pharmaceuticals, it can be incorporated into new drug formulations or utilised to develop novel treatments for bacterial infections, addressing the growing concern of antibiotic resistance. The textile industry can employ this compound to create insect repellent and antimicrobial fabrics, which are increasingly in demand for medical textiles, sportswear, and everyday clothing. Such applications could enhance hygiene standards, reduce odour-causing bacteria, and potentially extend the lifespan of textile products. Future research should focus on optimising the efficacy of the compound, exploring its mechanisms of action, and conducting comprehensive safety assessments to ensure its suitability for diverse applications.

Commented [MG8]: The conclusion and summary clearly and concisely present the results obtained in the study on the antimicrobial activity of the essential oils of *Ocimum gratissimum* and *Cymbopogon flexuosus*. The FTIR analysis is well summarized, highlighting the main functional groups present in the essential oils, while the results of antimicrobial activities are presented objectively, with emphasis on the superior efficacy of *Cymbopogon flexuosus* compared to *Ocimum gratissimum*. The explanation of the mechanisms of action, involving the destruction of bacterial and fungal cell membranes, is also well integrated. Furthermore, the section provides a valuable discussion on the potential applications of these essential oils in various industries, including food, agriculture, pharmaceuticals, and textiles. The proposal to optimize the compound's effectiveness and carry out safety assessments in future research is pertinent and highlights the need for additional studies to ensure its safe and effective application. The text is well structured, but could be enriched with a mention of the limitations of the study, such as the scope of the research, number of strains tested or possible variations in results in different conditions. Overall, it is a solid conclusion that summarizes the findings well and points to directions for future investigation and applications.

References

1. Hassid A, Hassan H, El Khatib S, Krayem M, Khaled S, Salla M. A Review on the Versatile Applications of Plant-Based Essential Oils in Food Flavoring, Culinary Uses and Health Benefits [Internet]. 2024. Available from: <https://doi.org/10.20944/preprints202406.2073.v1>
2. Djilani A, Dicko A. The Therapeutic Benefits of Essential Oils [Internet]. 2012. Available from: <https://doi.org/10.5772/25344>
3. Raut JS, Karuppaiyl SM. A status review on the medicinal properties of essential oils. *Ind Crops Prod.* 2014;62(62):250–64.
4. Huong LT, Chau DTM, An NTG, Dai DN, Ogunwande IA. Essential Oils of Lauraceae: Antimicrobial Activity and Constituents of *Phoebe macrocarpa* C.Y. Wu Leaf Essent Oil Vietnam *J Essent Oil Bear Plants.* 2022;25(2):297–304.
5. Kaul VK, Vats SK. Essential oil composition of *Bothriochloa pertusa* and phyletic relationship in aromatic grasses. *Biochem Syst Ecol.* 1998;26(3):347–56.
6. Al-Harrasi A, Ahmed MM, Anwer K, Bhatia S, Sharma PB. Anti-Inflammatory, Antioxidant, and Immunomodulatory Effects of EOs [Internet]. 2022. p. 239–55. Available from: <https://doi.org/10.1201/9781003175933-18>
7. Reboredo-Rodríguez P, Varela-López A. Essential Oils from Aromatic Plants in Cancer Prevention and Treatment [Internet]. *springer*; 2021. 61–81 p. Available from: https://doi.org/10.1007/978-3-030-74035-1_4
8. Reichling J. Antiviral and Virucidal Properties of Essential Oils and Isolated Compounds - A Scientific Approach. *Planta Med.* 2021;88(8):587–603.
9. Shaaban HAE, El-Ghorab AH, Shibamoto T. Bioactivity of essential oils and their volatile aroma components: Review. *J Essent Oil Res.* 2012;24(2):203–12.
10. Grahovac M, Gvozdenac S, Lazic S, Indjic D, Hrustic J, Vukovic S, et al. Integrated management of causal agents of postharvest fruit rot of apple. *Pestic Fitomedicina.* 2011;26(4):289–99.
11. Khan GEL, Bashir S, Mehmood A, Ijaz R, Haneef N, Usman Raja M, et al. Essential Oils Based Nano Formulations against Postharvest Fungal Rots [Internet]. 2024. Available from: <https://doi.org/10.5772/intechopen.113834>
12. Kumar AP, Swamy JS, Santhoshini CNR, Gouthami P, Shahana S. A Review on Bio-Based Control of Post-Harvest Diseases. *Int J Plant Soil Sci.* 2023;35(19):212–22.
13. Sivakumar D, Bautista-Baños S. A review on the use of essential oils for postharvest decay control and maintenance of fruit quality during storage. *Crop Prot.* 2014;64:27–37.
14. Strano MC, Genovese F, Matera A, Paterna G, Allegra M, Altieri G, et al. Postharvest Technologies of Fresh Citrus Fruit: Advances and Recent Developments for the Loss Reduction during Handling and Storage. *Horticulturae.* 2022;8(7):612.
15. Harris SJ, Cormican M, Cummins E. Antimicrobial Residues and Antimicrobial-Resistant Bacteria: Impact on the Microbial Environment and Risk to Human Health—A Review. *Hum Ecol Risk Assess Int J.* 2012;18(4):767–809.

16. Lee J, Kim S, Jung H, Koo BK, Han JA, Lee HS. Exploiting Bacterial Genera as Biocontrol Agents: Mechanisms, Interactions and Applications in Sustainable Agriculture. *J Plant Biol.* 2023;66(6):485–98.
17. Ganjewala D, Luthra R. Essential Oil Biosynthesis and Regulation in the Genus *Cymbopogon*. *Nat Prod Commun.* 2010;5(1):1934578 1000500.
18. Kumar A, Lal R, Chanotiya C, Gupta A. Historical and contemporary development of novel chemotype varieties with high essential oil of lemongrass in India: A review. *J Med Aromat Plant Sci.* 2023;45(1):17–27.
19. Sangwan NS, Yadav U, Sangwan RS. Molecular analysis of genetic diversity in elite Indian cultivars of essential oil trade types of aromatic grasses (*Cymbopogon* species). *Plant Cell Rep.* 2001;20:437–44.
20. Khanuja SPS, Shasany AK, Pawar A, Lal RK. Essential oil constituents and RAPD markers to establish species relationship in *Cymbopogon* Spreng. *Poaceae Biochem Syst Ecol.* 2005;33:171–86.
21. Lawrence R, K. L, R. S, Gup-ta D. Antioxidant activity of lemon grass essential oil grown in North Indian plains. *Sci Temper.* 2015;IV:23–9.
22. Ajayi AM, Okhale SE, Ologe MO, Ben-Azu B, Ademowo OG, Adzu B. *Ocimum gratissimum* Linn. Leaf extract inhibits free radical generation and suppressed inflammation in carrageenan-induced inflammation models in rats. *J Basic Clin Physiol Pharmacol.* 2017;28(6):531–41.
23. Ugbogu OC, Emmanuel O, Agi GO, Ibe C, Ekweogu CN, Ude VC, et al. A review on the traditional uses, phytochemistry, and pharmacological activities of clove basil (*Ocimum gratissimum* L. *Heliyon.* 2021;7(11):08404.
24. Offiah VN, Chikwendu UA. Antidiarrhoeal effects of *Ocimum gratissimum* leaf extract in experimental animals. *J Ethnopharmacol.* 1999;68(1–3):327–30.
25. Oyebode O, Jeje O, Oyelade W. Phytochemical screening and antibacterial activities of *Ocimum gratissimum* against some clinical isolates. *GSC Biol Pharm Sci.* 2023;24(2):148–53.
26. Coates J. Interpretation of infrared spectra, a practical approach. *Encycl Anal Chem.* 2000;12:10815–37.
27. Osée Muyima NY, Nziweni S, Mabinya LV. Antimicrobial and antioxidative activities of *Tagetes minuta*, *Lippia javanica* and *Foeniculum vulgare* essential oils from the Eastern Cape Province of South Africa. *J Essent Oil Bear Plants.* 2004;7(1):68–78.
28. Gupta A, Muhury R, Ganjewala D. A study on antimicrobial activities of essential oils of different cultivars of lemongrass (*cymbopogon flexuosus*). *Pharm Sci.* 2016;22(3):164–9.
29. Júnior A. Influence of lemongrass essential oil (*cymbopogon flexuosus*) supplementation on diabetes in rat model. *Life.* 2024;14(3):336.
30. Lopes V, Carvalho A, Bertolucci S, Roza H, Figueiredo F, Pinto J. Macronutrient suppression in nutrient solution alters the growth and citral content of *cymbopogon flexuosus*. *J Agric Sci.* 2019;11(6):320.

31. Menezes C, Perez A, Sousa J, Ramos K, Neto H, Filho A, et al. Evaluation of antifungal activity and mode of action of citral against *Cladophialophora carrionii*. *Rev Colomb Cienc Quím Farm* [Internet]. 2023;51(3). Available from: <https://doi.org/10.15446/rcciquifa.v51n3.100452>
32. Patterson J, McElmeel L, Wiederhold N. In vitro activity of essential oils against gram-positive and gram-negative clinical isolates, including carbapenem-resistant enterobacteriaceae. *Open Forum Infect Dis* [Internet]. 2019;6(12). Available from: <https://doi.org/10.1093/ofid/ofz502>
33. Shi C, Song K, Zhang X, Sun Y, Sui Y, Chen Y, et al. Antimicrobial activity and possible mechanism of action of citral against *Cronobacter sakazakii*. *Plos One*. 2016;11(7):0159006.
34. Mokarizadeh M, Kafil H, Ghanbarzadeh S, Alizadeh A, Hamishehkar H. Improvement of citral antimicrobial activity by incorporation into nanostructured lipid carriers: a potential application in food stuffs as a natural preservative. *Res Pharm Sci*. 2017;12(5):409.
35. Rodrigues L, Coelho E, Madeira R, Teixeira P, Henriques I, Coimbra M. Food ingredients derived from lemongrass byproduct hydrodistillation: essential oil, hydrolate, and decoction. *Molecules*. 2022;27(8):2493.
36. Gao S, Liu G, Li J, Chen J, Li L, Li Z, et al. Antimicrobial activity of lemongrass essential oil (*Cymbopogon flexuosus*) and its active component citral against dual-species biofilms of *Staphylococcus aureus* and *Candida* species. *Front Cell Infect Microbiol* [Internet]. 2020;10. Available from: <https://doi.org/10.3389/fcimb.2020.603858>
37. Korenblum E, Goulart F, Rodrigues I, Abreu F, Lins U, Alves P, et al. Antimicrobial action and anti-corrosion effect against sulfate reducing bacteria by lemongrass (*Cymbopogon citratus*) essential oil and its major component, the citral. *Amb Express*. 2013;3(1):44.
38. Chatrath A, Gangwar R, Kumari P, Prasad R. In vitro anti-biofilm activities of citral and thymol against *Candida tropicalis*. *J Fungi*. 2019;5(1):13.
39. Liao Z, Lin K, Liao W, Xie Y, Yu G, Shao Y, et al. Transcriptomic analyses reveal the potential antibacterial mechanism of citral against *Staphylococcus aureus*. *Front Microbiol* [Internet]. 2023;14. Available from: <https://doi.org/10.3389/fmicb.2023.1171339>
40. Ouyang Q, Tao N, Zhang M. A damaged oxidative phosphorylation mechanism is involved in the antifungal activity of citral against *P* [Internet]. 2018. Available from: <https://doi.org/10.3389/fmicb.2018.00239>
41. Sanches L, Marinello P, Fagundes T, Morgado-Díaz J, de-Freitas-Junior J, Cecchini R, et al. Cytotoxicity of citral against melanoma cells: the involvement of oxidative stress generation and cell growth protein reduction. *Tumor Biol*. 2017;39(3):101042831769591.
42. Song W, Yin Z, Lu X, Shen D, Dou D. Plant secondary metabolite citral interferes with *Phytophthora capsici* virulence by manipulating the expression of effector genes. *Mol Plant Pathol*. 2023;24(8):932–46.
43. Wang W, Ren Z, Wang L, Cai Y, Ma H, Fang L, et al. Nanoparticle-stabilized encapsulation of borneol and citral: physicochemical characteristics, storage stability, and enhanced antibacterial activities. *J Food Sci*. 2021;86(10):4554–65.
44. Ashokkumar K, Dhanya MK, Pandian A, Murugan M, Vellaikumar S. Phytochemistry and pharmacological properties of *Ocimum gratissimum* (L.) extracts and essential oil - A critical review. *J Curr Opin Crop Sci*. 2021;2(1):138–48.

45. Lahiri D, Nag M, Dutta B, Dey S, Mukherjee D, Joshi S, et al. Antibiofilm and anti-quorum sensing activities of eugenol and linalool from *Ocimum tenuiflorum* against *Pseudomonas aeruginosa* biofilm. *J Appl Microbiol*. 2021;131(6):2821–37.
46. Varghese R. Antibacterial activity of herbal formulation against common oral pathogens. *Bioinformation*. 2023;19(5):663–9.
47. Chimnoi N, Reuk-Engam N, Chuysinuan P, Khlaychan P, Khunnawutmanotham N, Chokchaichamnankit D, et al. Characterization of essential oil from *Ocimum gratissimum* leaves: Antibacterial and mode of action against selected gastroenteritis pathogens. *Microb Pathog*. 2018;118:290–300.
48. Silva JC, Tavares DSA, Rodrigues FFG, Freitas TS, Macedo NS, Junior FNP, et al. Evaluation of antibacterial and toxicological activities of essential oil of *Ocimum gratissimum* L. and its major constituent Eugenol [Internet]. 2022. Available from: <https://doi.org/10.21203/rs.3.rs-1383046/v1>
49. Marchese A, Barbieri R, Coppo E, Orhan İ, Daglia M, Nabavi S, et al. Antimicrobial activity of eugenol and essential oils containing eugenol: a mechanistic viewpoint. *Crit Rev Microbiol*. 2017;43(6):668–89.
50. Bednarczyk P, Wróblewska A, Markowska-Szczupak A, Ossowicz-Rupniewska P, Nowak M, Kujbida M, et al. UV curable coatings based on urethane acrylates containing eugenol and evaluation of their antimicrobial activity. *Coatings*. 2021;11(12):1556.
51. Pei R, Zhou F, Ji B, Xu J. Evaluation of the combined antibacterial effects of eugenol, cinnamaldehyde, thymol, and carvacrol against *E. coli* using an improved method. *J Food Sci* [Internet]. 2009;74(7). Available from: <https://doi.org/10.1111/j.1750-3841.2009.01287.x>
52. Yadav M, Chae S, Im G, Chung J, Song J. Eugenol: a phyto-compound effective against methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* clinical strain biofilms. *Plos One*. 2015;10(3):0119564.
53. Biasi-Garbin R, Otaguiri E, Morey A, Silva M, Morguette A, Lancheros C, et al. Effect of eugenol against *Streptococcus agalactiae* and synergistic interaction with biologically produced silver nanoparticles. *Evid Based Complement Alternat Med*. 2015;1–8.
54. Devi K, Nisha S, Sakthivel R, Pandian S. Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhi* by disrupting the cellular membrane. *J Ethnopharmacol*. 2010;130(1):107–15.
55. Sedlaříková J, Janalíková M, Peer P, Pavlatkova L, Minařík A, Pleva P. Zein-based films containing monolaurin/eugenol or essential oils with potential for bioactive packaging application. *Int J Mol Sci*. 2021;23(1):384.
56. Ulanowska M, Olas B. Biological properties and prospects for the application of eugenol—a review. *Int J Mol Sci*. 2021;22(7):3671.
57. Shajari M, Ahmadi N, Zamani M, Rostamizadeh K, Shapouri R. Hospital wastewater treatment using eco-friendly eugenol nanostructured lipid carriers: formulation, optimization, and in vitro study for antibacterial and antioxidant properties. *Water Environ Res* [Internet]. 2022;94(7). Available from: <https://doi.org/10.1002/wer.10751>
58. Rezende DADCS, Lunguinho ADS, Nelson DL, Alves E, Ferreira VRF, Cardoso MDG, et al. on mycotoxin-producing *Aspergillus flavus* and *Aspergillus ochraceus* antifungal properties of

- essential oils. H. M, S C, editors. *Eff Essent Oils Satureja Mont L* [Internet]. 2021;368(19). Available from: <https://doi.org/10.1093/femsle/fnab137>
59. Irkin R, Korukluoglu M. Effectiveness of *Cymbopogon citratus* L. Essential Oil to Inhibit the Growth of Some Filamentous Fungi and Yeasts. *J Med Food*. 2009;12(1):193–7.
 60. Regnier T, Combrinck S, Veldman W, Plooy W. Application of essential oils as multi-target fungicides for the control of *Geotrichum citri-aurantii* and other postharvest pathogens of citrus. *Ind Crops Prod*. 2014;61:151–9.
 61. Al-Ghanayem AA. Phytochemical analysis of *Cymbopogon flexuosus* (lemongrass) oil, its antifungal activity, and role in inhibiting biofilm formation in *Candida albicans* MTCC854. *J King Saud Univ - Sci*. 2022;34(5):102072.
 62. Taweechaisupapong S, Aieamsaard J, Chitropas P, Khunkitti W. Inhibitory effect of lemongrass oil and its major constituents on *Candida* biofilm and germ tube formation. *South Afr J Bot*. 2012;81(81):95–102.
 63. Prajapati M, Thakkar S, Shah M, Acharya D, Ranginwala A, Agrawal P. Antifungal effects of tulsi, garlic, cinnamon and lemongrass in powder and oil form on *Candida albicans*: An in vitro study. *J Oral Maxillofac Pathol*. 2021;25(2):306–12.
 64. Leite M, Bezerra A, Sousa J, Guerra F, Lima E. Evaluation of antifungal activity and mechanism of action of citral against *C. albicans*. *Evid Based Complement Alternat Med* [Internet]. 2014;2014(1). Available from: <https://doi.org/10.1155/2014/378280>
 65. Wardana A, Wigati L, Van T, Tanaka F, Tanaka F. Antifungal features and properties of Pickering emulsion coating from alginate/lemongrass oil/cellulose nanofibres. *Int J Food Sci Technol*. 2022;58(2):966–78.
 66. Ouyang Q, Tao N, Jing G. Transcriptional profiling analysis of *penicillium digitatum*, the causal agent of citrus green mold, unravels an inhibited ergosterol biosynthesis pathway in response to citral. *BMC Genomics* [Internet]. 2016;17(1). Available from: <https://doi.org/10.1186/s12864-016-2943-4>
 67. Tang X, Shao Y, Tang Y, Zhou W. Antifungal activity of essential oil compounds (geraniol and citral) and inhibitory mechanisms against grain pathogens (*Aspergillus flavus* and *Aspergillus ochraceus*). *Molecules*. 2018;23(9):2108.
 68. de-la-Fuente I. In vitro and in vivo activity of citral in combination with amphotericin B, anidulafungin, and fluconazole against *Candida auris* isolates. *J Fungi*. 2023;9(6):648.
 69. Koffi Gaston K. Antifungal Activity of Essential Oils Extracted from *Monodora myristica* (Gaertn), *Ocimum gratissimum* L. and *Zingiber officinalis* Roscoe on Post-harvest Anthracnose of Mango Fruit (*Mangifera indica* L.) Variety Kent in Côte d'Ivoire. *Int J Sci*. 2015;1(11):8–18.
 70. Matasyoh LG, Matasyoh JC, Wachira FN, Kinyua MG. Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *Afr J Biotechnol*. 2007;6:760–5.
 71. Silva MRR, Lemos JA, Souza LKH, Paula JR, Costa CR, Fernandes OFL, et al. Antifungal activity of *Ocimum gratissimum* towards dermatophytes. *Mycoses*. 2005;48(3):172–5.

72. Faria TDJ, Souza JRPD, Ishikawa NK, Ferreira RS, Yassumoto L, Barbosa ADM. Antifungal activity of essential oil isolated from *Ocimum gratissimum* L. (eugenol chemotype) against phytopathogenic fungi. *Braz Arch Biol Technol.* 2006;49(6):867–71.
73. Tripathi RD, Sharma ML, Balasubrahmanyam VR, Nigam SK, Banerji R. Toxicity of essential oil from a new strain of *Ocimum gratissimum* (Clocimum) against betelvine pathogenic fungi. *Agric Biol Chem.* 1985;49(8):2277–82.
74. Pandey S. ANTIBACTERIAL AND ANTIFUNGAL ACTIVITIES OF OCIMUM GRATISSIMUM L. *Int J Pharm Pharm Sci.* 2017;9(12):26.
75. Mohr F, Lermen C, Gazim Z, Gonçalves J, Alberton O. Antifungal activity, yield, and composition of *ocimum gratissimum* essential oil. *Genet Mol Res [Internet].* 2017;16(1). Available from: <https://doi.org/10.4238/gmr16019542>
76. Akpo A. In vitro antifungal activity of aqueous extract and essential oil of african basil (*ocimum gratissimum* l. *North Afr J Food Nutr Res.* 2023;7(16):136–45.
77. Madjouko M, Tchameni S, Tchinda E, Jazet P, Kamsu P, Kamga V, et al. Inhibitory effects of essential oils from *ocimum basilicum* and *ocimum gratissimum* on *colletotrichum musae*: the causal agent of bananas anthracnose. *J Phytopathol.* 2019;167(5):257–64.
78. Matasyoh L, Matasyoh J, Wachira F, Kinyua M, Muigai A, Mukiama T. Antimicrobial activity of essential oils of *ocimum gratissimum* l. from different populations of kenya. *Afr J Tradit Complement Altern Med [Internet].* 2008;5(2). Available from: <https://doi.org/10.4314/ajtcam.v5i2.31272>