

Original Research Article

Impact of Drying Methods on the Chemical and Storage Qualities of Powdered 'Daddawa' From African Locust Bean (*Parkia biglobosa*) Seeds

ABSTRACT

The study investigated impact of different drying methods on the chemical and storage qualities of 'daddawa' powder produced from African locust bean (*Parkia biglobosa*) seeds. The seeds were processed using slight modification of the traditional method. Fermented seeds were divided into four portions, three parts were dried using cabinet, oven and sun giving samples CB1, OC2 and SD3 respectively. Dried samples were comminuted using attrition mill and packaged. The fermented undried seeds (UA0 - control) was refrigerated before analysis. The samples were subjected to chemical analysis using standard procedures. Results of proximate composition revealed that moisture content of CB1 ($13.04^b \pm 0.00$), OC2 ($12.64^{bc} \pm 0.47$) and SD3 ($12.03^c \pm 0.00$) were within range of flour (10-14%). Protein content showed that SD3 ($25.17^a \pm 0.09$) was significantly ($P = .05$) different to the rest samples. The result of vitamin content of 'daddawa' powder shows that SD3 ($8.88^a \pm 0.00$) was highest in vitamin B1 followed by UA0 ($8.67^a \pm 0.15$) with no significant ($P = .05$) difference. The results show an array of mineral content including calcium, potassium, iron, magnesium, phosphorus, zinc, manganese and sodium. pH value was highest in OC2 (6.9) and SD3 (6.9) and least in CB1 (6.2). The study revealed that cabinet dried samples could be more stable owing to its lowered pH value. Also, both water activity and microbial counts of bacteria and fungi increased with storage time. It is recommended that further studies should be carried out to determine the *invitro* and *invivo* digestibility of the protein for both the dried and undried (control) samples. Also, to ameliorate losses of nutrients during drying, lower temperature should be adopted.

Keywords: Daddawa; drying methods; chemical qualities; powder; microbial; *Parkia biglobosa*.

1. INTRODUCTION

Daddawa is an alkaline fermented condiment which is processed from the seeds of the African locust bean and has a final pH above 7 [1]. African locust bean (*Parkia biglobosa*) and other oil seeds such as melon seed, castor seed, soy-bean etc. are fermented to produce condiments [2]. Fermented locust bean is a well-known condiment with characteristic ammoniacal odour and flavor which enhance the taste of traditional soups and sauces especially those used as accompaniments to starchy foods. It is generally added to soups as low cost meat substitute by low income families in parts of Nigeria and is known to contribute to the caloric and protein intake [3]. *Daddawa* is common with the Hausa tribe in

Comment [U1]: P values should be well defined. That is it < and not = since the values gotten from the analysis are within range and not precise.

the northern part of Nigeria and the Middle belt. Fermented or local seasonings are good source of nutrients and could be used to produce complementary food supplements. Local seasonings can also add extra nutrients such as vitamins B due to traditional processes which they undergo.

Fermented products remain a thing of interest since they do not require refrigeration during distribution and storage. Traditionally, fermented locust beans are molded into paste and sundried [4]. However, some factors limit their production and acceptability by the generality of the people. As observed by [5], the traditional condiments have not attained commercial status due to the very short shelf-life, objectionable packaging materials, stickiness and the characteristics putrid odour.

One of the constraints to utilization of *daddawa* is the poor quality, short shelf life and high putrid odour which affect its acceptability and storage stability. Hence, the condiment has remained localized. Consequently, converting *daddawa* into powder will not only extend its shelf life but also reduce the high offensive odour associated with the product thereby improving the acceptability. Also, significant contributions have been made on local condiments from both legumes and wild trees, however, little or no knowledge is available on the impact of drying on the quality characteristics of *daddawa* powder.

The study was aimed at evaluating the chemical, sensory and storage properties of powdered *daddawa* produced from African locust bean (*Parkia biglobosa*) seeds using oven, cabinet and sundrying methods.

2. MATERIAL AND METHODS

2.1 Material

Processing raw materials such as African locust bean (*Parkia biglobosa*) ("*kalwa*" in Hausa) was purchased at Muda Lawal market in Bauchi metropolis. Drying machines such as cabinet (APV MITCHELL DRYERS CARLISLE ENGLAND ORDER NO. 5213/82) and oven (PROLABO NO. 35349) dryers were sourced from the Department of Food Science and Technology, Federal Polytechnic Bauchi where processing and analysis was carried out.

2.2 Formulation of samples and production process

The traditional method was adopted for the production of *daddawa* with slight modification [6]. Fermented *daddawa* was divided into four portions, three parts were dried using cabinet (80°C), oven (80°C) and sun (ambient) giving samples CB1, OC2 and SD3 respectively. The dried 'daddawa' was comminuted using attrition milling machine to get samples of powdered *daddawa* (Fig.1). The fermented undried seeds (UA0 - control) was refrigerated before analysis. Three samples of 'daddawa' powder including control (UA0) were subjected to chemical analysis which included proximate composition, vitamin content, mineral content, pH and titratable acidity, water activity, and color determination using standard analytical methods [7].

Comment [U2]: More literatures should be written on the storage and storage qualities of daddawa.

Comment [U3]: Sentence should be reference.

Comment [U4]: Which state and which country?

Comment [U5]: Three method of sun drying should be stated as well.

Comment [U6]: Figure 1 is a flow chart not comminution process. The reason(s) for the refrigeration of the fermented undried seeds should be stated. The undried seeds should be milled fermented undried seeds.




Figure 1: Flow chart for powdered 'daddawa' production from African locust bean seeds (*Parkia biglobosa*) with modification [6].

2.2 Determination of Chemical properties of powdered 'daddawa'

Proximate composition, vitamin content, mineral content, Titratable acidity, pH and color were determined using standard chemical methods [7].

2.3 Sensory Evaluation of 'Daddawa' Samples

Sensory evaluation was determined using 15 untrained panellists randomly selected with the 9-point hedonic scale[8].

2.5 Determination of Total Microbial Count

The Total Microbial Count (TMC) also known as the Total Viable Count (TVC) was determined using the procedure [9]. 23.4g of potato dextrose agar was weighed and dissolved in 300ml of distilled water in a conical flask. 5.6g of nutrient agar was also weighed and dissolved in 200ml of distilled water. 9ml of distilled water was pipetted into 5 test tubes each for the samples, and plucked with cotton wool. The prepared agars, distilled water, petri-dishes and pipettes was sterilized in the autoclave (Model: YX-280A. GULFEX MEDICAL AND SCIENTIFIC ENGLAND) at 121°C for 15 minutes at 15psi. After sterilizing,

Comment [U7]: Authors should state the method(s) used for storage quality of the product.

the test tubes was labelled 10^{-1} to 10^{-5} and the petri-dishes was also labelled appropriately. 10g of each sample was weighed into a homogenate containing 50ml of distilled water in a beaker. This was stirred using a stirrer to a homogenous solution which was later filtered to get a clear solution. 1ml of the homogenate was serially diluted into the 9ml of test tubes labelled 10^{-1} . After homogenizing, 1ml was taken from here and transferred to the 10^{-2} and continued up to the 10^{-5} dilution for each sample. 0.1ml aliquot from the 10^{-5} dilution was transferred into labelled petri-dishes in duplicate for each sample. The potato dextrose agar and nutrient agar were allowed to cool, after which each was poured into the petri-dish containing 0.1ml aliquot for each sample. The petri-dishes were rocked gently and allowed to solidify. The plate was then incubated by placing the nutrient agar plate inside the incubator (M^c Jefferson – UK) at 37°C for 24 hours and the potato dextrose agar plate was put on the shelf at room temperature for 72 hours. At the end of the incubation period, the colonies on each plate was counted with a colony counter (J-2 Colony Meter PEC MEDICAL USA) and results calculated as colony forming units (CFU) per ml.

Formula

$$c = n/vd$$

Where c = colony forming unit per ml (cfu/ml)

n = number of colonies

d = dilution blank factor

v = volume transferred to plate

2.6 Statistical Analysis

Results were presented as mean \pm standard deviations of duplicate determinations. The data obtained from the various determinations were subjected to analysis of variance using the GenStat Statistical Software, Seventeenth Edition (17.1). One way analysis of variance (ANOVA) was used for the comparison of the means. Differences between means was considered to be significant at (P = .05) using the Tukey multiple range test.

3. RESULTS AND DISCUSSION

3.1 Proximate composition of 'daddawa' powder

The proximate composition of 'daddawa' powder presented in "Table 1" showed that the control sample UA0 ($23.38^a \pm 0.06$) is significantly different to samples CB1 ($13.04^b \pm 0.00$), OC2 ($12.64^{bc} \pm 0.47$) and SD3 ($12.03^c \pm 0.00$) in moisture content at 5% confidence level. Whereas CB1 and SD3 shows significant difference (P = .05), OC2 appeared same with others in their moisture content. This result unarguably appears under range of moisture content of flour (10-14%) which makes it ideal for safe keeping. The result is equally lower than that reported for moisture content (26.61%) by [10]. Expectedly, protein, fat, ash, crude fibre and carbohydrate contents expressed higher values in the dried samples than the undried (control); with SD3 ($25.17^a \pm 0.09$) being the highest and UA0 ($19.71^c \pm 0.38$) as the least in protein value. Proximate composition result in general ranged from $2.94^c \pm 0.07$ (UA0) to $41.79^a \pm 0.44$ (OC2) in ash and carbohydrate respectively. The protein and carbohydrate contents were way above that reported by [10] which was given as 16.98 and 22.28% respectively. It can be deduced that drying help to condense the nutrient values especially the non-volatile ones.

Comment [U8]: Formula should be removed. Equation 1 was used to determined the colony forming of the ----- . Also, = should be removing when defining the terms of the equation e.g c is the colony forming etc,

Comment [U9]: The sub section is results and discussion. The first section should comprises of all the results of ifndings from the resaerch i.e 3.1. should be written as Tables ----- shows the ----- . While 3.2 should discussed the obtained results. Authors are advised to re-write this section accordingly.

Table 1: Results of Proximate Composition of Powdered “Daddawa” Samples

Parameter	UA0	CB1	OC2	SD3
Proximate composition (% 100g sample)				
Moisture	23.38 ^a ±0.06	13.04 ^b ±0.00	12.64 ^{bc} ±0.47	12.03 ^c ±0.00
Protein	19.71 ^c ±0.38	24.26 ^b ±0.06	23.80 ^b ±0.04	25.17 ^a ±0.09
Crude fat	9.50 ^d ±0.08	24.26 ^b ±0.06	12.04 ^c ±0.05	12.93 ^a ±0.02
Ash	2.94 ^c ±0.07	4.53 ^a ±0.59	3.28 ^{bc} ±0.07	3.89 ^{ab} ±0.01
Crude fibre	4.93 ^d ±0.01	6.16 ^b ±0.04	6.44 ^a ±0.01	5.20 ^c ±0.02
Carbohydrate	39.54 ^b ±0.44	39.63 ^b ±0.48	41.79 ^a ±0.44	40.79 ^a ±0.13

Values are means ± standard deviations of duplicate determinations. Means in same row with different superscript are significantly (P = .05) different.

Key: UA0 = Undried fermented ‘daddawa’ seeds

CB1 = Cabinet dried ‘daddawa’ powder

OC2 = Oven dried ‘daddawa’ powder

SD3 = Sundried ‘daddawa’ powder

3.2 Vitamin content of ‘daddawa’ powder

The result of vitamin content of ‘daddawa’ powder “Table 2” showed that SD3 (8.88^a±0.00) was highest in vitamin B1 followed by UA0 (8.67^a±0.15) with no significant difference (P = .05) and the least was expressed in CB1 (7.17^b±0.02) which has no significant difference (P = .05) with OC2 (7.29^b±0.08). ‘Daddawa’ powder showed highest value in vitamin D content, with UA0 (87.98^a±1.92) being significantly different (P = .05) to the rest of the samples. While vitamin E expressed least in vitamin content, sample UA0 (2.18^a±0.23) is significantly different (P = .05) to the rest of the samples. Vitamins B1 and D expressed values higher than that reported by [11] which were 0.77mg/100g and 0.29mg/100g respectively. However, the same author reported a value for vitamin E (23.34mg/100g) which was higher than that obtained for the experimental samples.

Table 2: Results of Vitamin analysis of Powdered “Daddawa” Samples

Parameter	UA0	CB1	OC2	SD3
Vitamin content (mg/100g)				
Vitamin B1	8.67 ^a ±0.15	7.17 ^b ±0.02	7.29 ^b ±0.08	8.88 ^a ±0.00
Vitamin D	87.98 ^a ±1.92	70.38 ^b ±0.08	66.23 ^c ±0.00	71.39 ^b ±0.07
Vitamin E	2.18 ^a ±0.23	1.30 ^b ±0.01	1.10 ^b ±0.01	1.30 ^b ±0.01
Vitamin K	12.50 ^a ±0.08	12.38 ^a ±0.05	12.61 ^a ±0.53	12.65 ^a ±0.01

Values are means ± standard deviations of duplicate determinations. Means in same row with different superscript are significantly (P = .05) different.

Key: UA0 = Undried fermented ‘daddawa’ seeds

CB1 = Cabinet dried ‘daddawa’ powder

OC2 = Oven dried ‘daddawa’ powder

SD3 = Sundried ‘daddawa’ powder

3.3 Mineral content of ‘daddawa’ powder

The 'daddawa' powder samples showed an array of mineral content "Table 3" including calcium, potassium, iron, magnesium, phosphorus, zinc, manganese and sodium. Potassium had the highest value with SD3 (174.90^a±2.22) being significantly different (P = .05) to the other samples including the control as seen in "Table 3". Calcium followed in the amount expressed by minerals and then magnesium. In calcium content UA0 (123.60^a±0.08) was significantly different (P = .05) to other samples. Iron was least expressed with no significant difference (P = .05) among the samples. All mineral values for the samples were higher than those reported by [11] for fermented oven dried *Parkia biglobosa* seeds except iron which was 14.13±0.01 mg/100g.

Table 3: Results of Mineral analysis of Powdered "Daddawa" Samples

Parameter	UA0	CB1	OC2	SD3
Mineral content (mg/100g)				
Calcium	123.60 ^a ±0.08	119.10 ^b ±1.27	106.80 ^c ±0.78	105.30 ^c ±0.78
Potassium	139.80 ^c ±2.14	150.80 ^b ±0.70	122.20 ^d ±0.00	174.90 ^a ±2.22
Iron	1.08 ^a ±0.09	1.15 ^a ±0.06	1.10 ^a ±0.01	1.23 ^a ±0.02
Magnesium	103.00 ^b ±0.56	102.30 ^b ±0.00	100.60 ^c ±0.53	108.60 ^a ±0.71
Phosphorus	32.95 ^c ±0.56	40.40 ^b ±0.08	40.50 ^b ±0.86	42.38 ^a ±0.07
Zinc	4.40 ^c ±0.05	4.65 ^b ±0.02	4.63 ^b ±0.00	4.75 ^a ±0.00
Manganese	4.29 ^b ±0.07	4.89 ^a ±0.01	4.07 ^b ±0.02	4.73 ^a ±0.15
Sodium	56.32 ^a ±0.62	56.51 ^a ±0.08	57.16 ^a ±1.16	56.56 ^a ±0.16

Values are means ± standard deviations of duplicate determinations. Means in same row with different superscript are significantly (P = .05) different.

Key: UA0 = Undried fermented 'daddawa' seeds
 CB1 = Cabinet dried 'daddawa' powder
 OC2 = Oven dried 'daddawa' powder
 SD3 = Sundried 'daddawa' powder

3.4 pH Content and Titratable acidity of 'Daddawa' Powder

The pH values expressed by the powdered 'daddawa' samples "Table 4" showed a similar range between the samples, except CB1 with a slightly lower value. While UA0, OC2 and SD3 had 6.8, 6.9, 6.9; CB1 was least with 6.2 respectively. This indicate that CB1 was more slightly acidic than the others.

Titratable acidity value also shown in "Table 4" was highest in CB1 (3.72%) and lowest in SD3 (2.30%). This indicate a little variation in the palmitic value of the samples. The pH result agrees with the titratable acidity content of the samples. As observed, CB1 with highest acidity value had lower pH value indicating a more slightly acidic content.

Table 4: Results of Chemical Property of Powdered "Daddawa" Samples

Parameter	UA0	CB1	OC2	SD3
pH	6.8	6.2	6.9	6.9
Titratable acidity (% palmitic acid)	2.72	3.52	2.43	2.30

Values are means ± standard deviations of duplicate determinations. Means in same row with different superscript are significantly (P = .05) different.

Key: UA0 = Undried fermented 'daddawa' seeds
 CB1 = Cabinet dried 'daddawa' powder
 OC2 = Oven dried 'daddawa' powder
 SD3 = Sundried 'daddawa' powder

3.5 Sensory evaluation of powdered 'daddawa' samples

The result of sensory evaluation is presented in "Table 5". The result showed that the mean values for the parameters of taste, smell, and general acceptability were: 8.53, 8.27, 7.60,

7.80; 8.07, 7.73, 7.60, 7.93; 8.07, 7.93, 7.93, 8.07 and 8.20, 7.87, 8.00, 7.47 for UA0, CB1, OC2 and SD3 respectively.

The result showed that there was significant difference ($P = .05$) among the samples only in taste, and it is between the control sample UA0 and OC2 as observed in the "Table". There was no significant difference ($P = .05$) between UA0, CB1, OC2 and SD3 in the parameters of smell, colour and general acceptability. Overall, the test panelist preferred UA0 based on taste.

Table 5: Result of sensory evaluation

Sample	Taste	Smell	Colour	General acceptability
UA0	8.53 ^a ±0.74	8.07 ^a ±0.96	8.07 ^a ±1.09	8.20 ^a ±0.94
CB1	8.27 ^{ab} ±0.70	7.73 ^a ±0.96	7.93 ^a ±0.96	7.87 ^a ±1.13
OC2	7.60 ^b ±0.99	7.60 ^a ±0.99	7.93 ^a ±0.96	8.00 ^a ±0.76
SD3	7.80 ^{ab} ±0.94	7.93 ^a ±1.09	8.07 ^a ±0.88	7.47 ^a ±1.06

Mean values along a column with different letter superscript are significantly different ($P = .05$)

Key: UA0 = Undried fermented 'daddawa' seeds

CB1 = Cabinet dried 'daddawa' powder

OC2 = Oven dried 'daddawa' powder

SD3 = Sundried 'daddawa' powder

3.6 Water activity (a_w) of powdered 'daddawa' samples

The result of water activity of 'daddawa' powder samples presented in "Table 6" indicate values obtained during three months storage period. In month one, CB1, OC2 and SD3 had 0.81, 0.80 and 0.89 water activity values respectively. In month two, the values increased as follows: 0.89, 0.87 and 0.91 for CB1, OC2 and SD3 respectively. In month three, the values obtained were 0.90, 0.90 and 0.945 for CB1, OC2 and SD3 respectively.

The result showed that there is significant difference ($P = .05$) between CB1 (0.81a) and SD3 (0.89b) and OC2 (0.80a) and SD3 (0.89b) in month one. There was no significant difference ($P = .05$) among the samples in month two, although the values increased significantly than that of month one. There was also slight increase in water activity of samples from month two to three of storage.

The result of water activity of the samples were slightly lower than the water activity requirement for bacteria growth which is 0.91 [12] with only sample SD3 expressing a value above this in month three. The increasing trend of the values with increase in storage time reflect a possible increase in microbial load with time. Water activity is a measure of water in a material that is available to react with or attach itself to other material; it is also called "free" water while the unavailable water is called "bound" water. Fungi require a water activity of at least 0.6 [12].

Table 6: Result of water activity (a_w) of powdered daddawa samples

Sample	Month 1	Month 2	Month 3
CB1	0.81 ^a ±0.01	0.89 ^a ±0.00	0.90 ^a ±0.00
OC2	0.80 ^a ±0.01	0.87 ^a ±0.00	0.90 ^a ±0.00
SD3	0.89 ^b ±0.00	0.91 ^a ±0.00	0.945 ^b ±0.10

Values are mean ± standard deviation of duplicate determinations. Mean values along a column with different letter superscripts are significantly different ($P = .05$).

Key: UA0 = Undried fermented 'daddawa' seeds

CB1 = Cabinet dried 'daddawa' powder

OC2 = Oven dried 'daddawa' powder

SD3 = Sundried 'daddawa' powder

3.7 Microbial analysis of powdered 'daddawa' samples

The result of microbial analysis of 'daddawa' samples are presented in "Table 7". The results shows values of total viable counts (TVC) and fungi counts obtained after production (zero month) and during period of storage (1-3 months). In zero month, UA0 expressed highest value of total viable count (TVC) and fungi count (FC) (4.7×10^4 cfu/ml and 4.8×10^1 cfu/ml) respectively than CB1, OC2 and SD3 as observed in Table 7. While fungi count had higher values than total viable count, during the storage period, both results are within safety limits. In month one, SD3 showed the highest value of TVC (7.1×10^1 cfu/ml) while OC2 (1.0×10^1 cfu/ml) was least. There is no significant difference between the samples. Similarly, there is no significant difference ($P = .05$) between the samples in FC values but OC2 (2.05×10^1 cfu/ml) was slightly higher.

There was no significant counts of the samples in TVC in month two but there was significant increase in TVC of samples in month three as CB1, OC2 and SD3 had 1.3×10^3 cfu/ml, 1.1×10^3 cfu/ml and 1.3×10^3 cfu/ml respectively. Also, FC gradually increased in month three as CB1, OC2 and SD3 had 1.9×10^1 cfu/ml, 2.4×10^1 cfu/ml and 2.4×10^1 cfu/ml respectively.

The total viable counts (TVC) of the powdered 'daddawa' samples during the storage period were less than that reported by [13] for locust bean-condiment powder produced in plastic ($5.03 \times 10^2 \log_{10}$ cfu/mL) and locust bean-condiment powder produced in calabash ($4.74 \times 10^2 \log_{10}$ cfu/mL).

Overall, the microbial results showed that both TVC and FC are within safety limit at 3 months of storage, as all the results were below minimum limit of standard (1.0×10^6 cfu/ml). It indicates that the products can keep for longer time on the shelf, being a dried product.

UNDER PEER REVIEW

Table 7: Results of microbial analysis (Month 0-3)

Sample	Month 0		Month 1		Month 2		Month 3	
	TVC*	FC*	TVC	FC	TVC	FC	TVC	FC
UA0	$4.7 \times 10^{4a} \pm 8485.2$ 5	$4.85 \times 10^{1a} \pm 53.5$ 0	-	-	-	-	-	-
CB1	$1.2 \times 10^{1b} \pm 0.00$	$2.85 \times 10^{1a} \pm 7.78$	$1.2 \times 10^{1a} \pm 0.00$	$1.9 \times 10^{1a} \pm 0.00$	$1.3 \times 10^{1a} \pm 0.0$ 0	$2.0 \times 10^{1a} \pm 0.00$	$1.3 \times 10^{3a} \pm 0.0$ 0	$1.9 \times 10^{1a} \pm 0.0$ 0
OC2	$1.0 \times 10^{1b} \pm 0.00$	$2.25 \times 10^{1a} \pm 0.71$	$1.0 \times 10^{1a} \pm 0.00$	$2.05 \times 10^{1a} \pm 0.7$ 1	$1.0 \times 10^{1a} \pm 0.0$ 0	$2.25 \times 10^{1b} \pm 0.7$ 1	$1.1 \times 10^{3a} \pm 0.0$ 0	$2.4 \times 10^{1a} \pm 0.0$ 0
SD3	$1.95 \times 10^{1b} \pm 2.12$	$3.25 \times 10^{1a} \pm 2.12$	$7.1 \times 10^{1a} \pm 83.4$ 4	$2.0 \times 10^{1a} \pm 0.00$	$1.3 \times 10^{1a} \pm 0.0$ 0	$2.3 \times 10^{1b} \pm 0.00$	$1.3 \times 10^{3a} \pm 0.0$ 0	$2.4 \times 10^{1a} \pm 0.0$ 0

Values are mean \pm standard deviation of duplicate determinations. Mean values along a column with different letter superscripts are significantly different (P = .05).

Key: *TVC = Total viable count. FC = Fungi count

UA0 = Undried fermented 'daddawa' seeds

CB1 = Cabinet dried 'daddawa' powder

OC2 = Oven dried 'daddawa' powder

SD3 = Sundried 'daddawa' powder

3.8 Color determination of 'daddawa' samples

The result of color determination of 'daddawa' samples is presented in "Table 8". It was determined using the L* a* b* color measurements. The result showed a significant difference (P = .05) in L* (values) between samples UA0, CB1, OC2 and SD3 as expressed: 20.77^c, 23.51^b, 18.17^d and 28.50^a respectively. The values ranged from 18.17-28.50 with OC2 being the least and SD3 being the highest respectively. As shown in the table, a* also expressed significant difference (P = .05) among the samples and ranged from 2.45-6.00 (SD3 and UA0) respectively. The values expressed for b* by the samples is zero except SD3 with -1. The value of b* obviously influenced the DC* and Hue outcome as seen in the table.

L* values (also called value) indicate degree of lightness from black to white (range = 0 – 100). Negative b* is blue, positive b* is yellow; negative a* is green and positive a* is red. Both a* and b* values range from -127 to 127 [14]. As observed in the table, the values of L* for all the samples is closely quarter away from black indicating that the product could be darker in colour. It was also observed from the results that the products leans towards red and yellow except for SD3 (b* = -1.00).

The Hue values of zero degrees obtained for samples UA0, CB1 and OC2 indicate a red hue direction for the product and a probability of a darker product as the hue moves from red to yellow. This analysis could suggest a depiction of the colour of 'daddawa' powder which is red-brown. The dark red-brown colour of the powdered samples was stable during the three months storage period. Colour is a sensory attribute of food that can be used to determine its quality and consumer acceptability. Colour evaluation in foods is an important aspect of safety and marketability in the food industry. It is so influential that it can greatly affect consumers' rating of food quality, taste and acceptability [13].

Table 8: Result of Color Determination of 'Daddawa' samples

Sample	L*	a*	a*	DC*	Hue (°)
UA0	20.77 ^c ±0.46	6.00 ^a ±0.00	0.00	6.00	0.00
CB1	23.51 ^b ±0.66	3.10 ^c ±0.00	0.00	3.10	0.00
OC2	18.17 ^d ±0.08	4.00 ^b ±0.00	0.00	4.00	0.00
SD3	28.50 ^a ±0.52	2.45 ^d ±0.07	-1.00	2.65	-22.21

Values are means ± standard deviations of duplicate determinations. Means in same column with different superscript are significantly (P = .05) different.

Key: UA0 = Undried fermented 'daddawa' seeds

CB1 = Cabinet dried 'daddawa' powder

OC2 = Oven dried 'daddawa' powder

SD3 = Sundried 'daddawa' powder

4. CONCLUSION

The study revealed that 'daddawa' powder dried by various methods expressed better quality in proximate composition than the control. However, drying affected the various vitamins analyzed but with little significant difference. Also, drying had no much effect on the minerals except calcium with much significant difference between the control and the rest. It also revealed that cabinet dried samples could be more stable owing to its lowered pH value. Either of the methods of drying could possibly give a nutritionally stable 'daddawa' powder as both methods compared favorably in the quality indices analyzed. Also, both water activity (a_w) and microbial counts of bacteria and fungi increased with storage time. The study indicated color stability during storage, however, sundried sample had a brighter color than the rest.

REFERENCES

1. Azokpota, P; Hounhouigan, D. J and Nago, M. C. Microbiological and chemical changes during the fermentation of African locust bean *Parkia biglobosa*, to produce *afitin*, *iru* and *sonru*, three traditional condiments produced in Benin. *Int. J. Food Microbiol* 2006;107: 304–309.
2. Omafuvbe, B. O; Falade, O. S; and Osuntogun, B. A. Chemical and biochemical changes in African. Locust Bean *Parkia biglobosa*, and melon *Citrullus vulgaris*, seeds during Fermentation to condiments. *Pakistan J. Nutrit.* 2004;3: 140–145.
3. Odunfa, S. A. Biochemical changes in fermenting African Locust Bean *Parkia biglobosa*, during *iru* Fermentation. *J Food Technol* 1985; 20: 295–303.

4. Simonyan, K. J. *African Locust Bean Seed Processing Prospects, Status and Challenges*. 1st Ed. Published by Naphtali Prints Nig. Ltd. 22, Market Street, Somolu Lagos. 2012; Pp 42-43.
5. Arogba, S. S., Ademola, A. and Elum, M. The effect of solvent treatment on the chemical composition and organoleptic acceptability of traditional condiments from Nigeria. *Plant Foods Human Nutrition*, 1995; 48:31-38.
6. Odunfa, S. A. African Fermented Foods: From Art to Science. *MIRCEN Journal of Applied Microbiology and Biotechnology*, 1988; 4: 259-273.
7. AOAC. Official Methods of Analysis of the Association of Official Analytical Chemists. Revised 21st edition. Washington D.C.; 2022; 19761-19776.
8. Ihekoronye AI and Ngoddy PO. Integrated Food Science and Technology for the Tropics. Macmillan Education Ltd. London and Oxford. 1985 Pp. 86, 92-94, 106-109, 178-189, 344, 353-357.
9. Ihekoronye AI and Ngoddy PO. Integrated Food Science and Technology for the Tropics. Macmillan Education Ltd. London and Oxford. 1985; Pp. 86, 92-94, 106-109, 178-189, 344, 353-357.
10. Muhammad, A. A., Egbebi, A. O. and Seidu, K. T. Nutritional and Microbiological Analyses of Fermented Locust Beans (*Parkia biglobosa*) and Fermented Melon (*Citrullus vulgaris*). *Savant Journal of Agricultural Research*, 2016; 2(1):1-6.
11. Termote, C; Odongo, N. O; Dreyer, B. S; Guissou, B; Parkouda, C. and Vinceti, B. Nutrient composition of *Parkia biglobosa* pulp, raw and fermented seeds: a systematic review, *Critical Reviews in Food Science and Nutrition*. 2020; Pp. 1-26
12. Allen, L. Quality Control: Water Activity Considerations for Beyond-use Dates. *International Journal of pharmaceutical compounding*. 2018; 22:288-293.
13. Awofadeju O. F. J., Ademola I. T., Alagbe O. A., Oyediran R. I. and Adekunle E. A. Studies of Harnessing Quality Attributes of Condiment Powder (Iru) Processed from Soy (Glycine Max) And African Locust Bean (*Parkia biglobosa*) Seeds. *Journal of Research in Forestry, Wildlife And Environment*, 2021; (13) 1:85-94. <http://www.aiol.info/index.php/jrfwe>
14. Eoptis, Colorimeter CLM-194 Interface version. 2014; 1.4.5.0 EoptisS.r.l Italy.