

Short Research Article

A Pilot Study Evaluating Medication Adherence in Geriatric Patients with Chronic Conditions

ABSTRACT:

Introduction: More established grown-ups are more inclined to ongoing circumstances and intermittent ailment like hypertension, diabetes mellitus, respiratory messes and so on which might require persistent prescription with different medications. Unfortunate consistency in this age bunch represents medicine wastage with the inflated cost of medical care and significant deterioration of the sickness with disability of death. A large portion of the human and financial expenses related to non-adherence can be tried not by further developing prescription adherence. Typically, geriatric patients have different co-morbidities, which can antagonistically influence drug consistence. Our review recognizes different reasons liable for low adherence.

Aim: To assess the level of medication compliance in the geriatric patients with chronic conditions and also analyze the reason behind the non-adherence.

Materials and methods: The study subjects were assessed by using 8 structured questionnaires as per modified morisky medication adherence scale (MMAS).

Results: A total of 88 subjects of geriatric patients with chronic conditions were assessed for the level of compliance for long term medications. The compliance level was assessed by interviewing patients with 8 item structured questionnaires as per modified morisky medication adherence scale (MMAS). The level of compliance was high in 34 and moderate in 39 and low in 15 of the study subjects.

Conclusion: To further develop drug adherence it is fundamental to recognize the purposes for the non-adherence like confounded systems, obliviousness about the sickness and inconveniences, physical and financial problems, forgetfulness, accessibility issues and absence of section of land providers and so on. Geriatric patients particularly tend to quit taking drugs by their own this can make it hard to anticipate improvement in the adherence. So, medication consistency is a vital element which increments helpful result.

KEY WORDS: Chronic illness, Geriatrics, Medication adherence, Factor influencing, Poor compliance.

INTRODUCTION:

Commented [DM1]: It must be relevant to the topic ' to Evaluate Medication Adherence in Geriatric Patients with Chronic Conditions

Commented [DM2]: Remove abbreviations from abstract

Commented [DM3]: Not relevant to topic ' Medication Adherence -Geriatric Patients - Chronic Conditions '

The elderly has turned into the fundamental survivors of persistent sicknesses, in light of the fact that they will generally experience the ill effects of multimorbidity which is characterized as people with at least two coinciding ongoing circumstances. Generally, the treatment of multimorbidity needs the drawn out taking different medicine and this drug intricacy might prompt unfortunate adherence which can decrease the adequacy of treatment and cause extra monetary misfortunes. WHO pronounced the explanation in its 2003 report taking drugs adherence that "expanding the ineffectualness of adherence mediations might greatly affect the wellbeing of the populace than any improvement in unambiguous clinical medicines. "Specialists have found that social help might be a successful defensive element of the more seasoned grown-up's medicine adherence both for purposeful and inadvertent nonadherence[1]. Adherence to therapy, a general medical problem, is of explicit significance in persistent sickness therapies. Prescription non-adherence is a typical and imperative general medical condition, explicitly among the geriatric individuals[2]. The personal satisfaction and the wellbeing related personal satisfaction of patients with ongoing sicknesses are fundamentally reduced[3,4]. a great many pharmacological treatments are right now accessible to draw out the existence of persistently sick people and to limit infection entanglements and incapacities. Since patients' adherence to both drug and suppliers' proposals expands the advantages of treatments, unfortunate adherence compromises the adequacy of regimens[5]. Non-adherence and abuse of meds might bring about bothersome results, for example, absence of medication viability, crisis visits, hospitalizations, inferior quality of life, or expanded medical services costs[6-8]. Understanding the hidden justification for non-adherence is fundamental to further develop prescription adherence] Many variables, for example, patient-related factors, condition-related factors, treatment related factors, medical services supplier factors, framework based factors, and financial elements, impact medicine adherence[9,10]. Taking drugs is a significant part of fitting medicine use. Patients with ongoing sicknesses frequently attempt various drug systems to deal with their circumstances, forestall entanglements and to keep up with their personal satisfaction. Numerous patients with ongoing sickness conditions are old and may battle in adhering to prescription directions, for example, taking the recommended measurements and following the organization plan. A patient's medicine information has been characterized as the familiarity with the medication name, reason, organization plan, unfavorable impacts or incidental effects, or extraordinary organization guidelines[11,12]. Patients who need to take numerous prescriptions are typically lacking in medicine information particularly the older[13]. Unfortunate medicine information can adversely affect drug adherence which might bring about expanded utilization of clinical assets, for example, doctor visits, lab tests, superfluous medicines, crisis division visits, clinic affirmations and treatment disappointment [14]. Unfortunate prescription information can likewise prompt drug wastage. Patient-related factors prompting unused and disposed of medicine stays obscure, but there are tremendous expense suggestions. We assess patients' medication knowledge and also assessed factors affecting their knowledge.

Commented [DM4]: 'world health organization' Define it for first time

Commented [DM5]: It is so far insert the newest report

Commented [DM6]: Not consistent with the paragraph

Commented [DM7]: Start with upper case

Commented [DM8]: Define it ?

Commented [DM9]: Older adults

Commented [DM10R9]: Finally the introduction need to flow systematically and define the all meaning completely

2. MATERIALS AND METHODS:

2.1. Study Type

This was a Prospective observational study. All the patients who met the inclusion criteria were taken as study population.

Commented [DM11]: design

2.2. Study Site

This study was done in a Tertiary care hospital, Virudhunagar, India

Commented [DM12]: setting

Commented [DM13]: Add more regarding the setting

2.3. Study Population

The study population for this study was Elderly patients (aged 50 years and above) with chronic illnesses attending the outpatient department of geriatrics. The inclusion and exclusion criteria were as follows:

2.3.1. Inclusion Criteria

- Geriatric patients aged 50 years or older.
- Diagnosed with chronic illnesses.
- Receiving treatment in the outpatient department.

2.3.2. Exclusion Criteria

- Patients admitted to the inpatient ward.
- Patients with mental illnesses.

2.4. Sample Size

A total of **88 patients** were included in the study.

Data Collection

- **Method:** Data was collected using a structured, 8-item questionnaire based on the **Modified Morisky Medication Adherence Scale (MMAS)**.
- **Procedure:** Patients were categorized into three adherence levels: low, moderate, and high, based on their responses.
- **Consent:** Written informed consent was obtained from all participants after explaining the study's purpose and procedures.

Commented [DM14]: Mention how are you perform your data collection, as well the duration of data collection,

Commented [DM15]: Ethical committee “?? The consent

Data Entry and Analysis




FIGURE 1 NUMBER OF SUBJECTS BASED ON AGE:

PEER REVIEW




FIGURE 2 NUMBER OF SUBJECTS BASED ON DISEASE:

Commented [DM16]: The figure not applied well

FIGURE 3 NUMBER OF NON-ADHERENT PATIENTS BASED ON REASONS:

Commented [DM17]: In percentages???




FIGURE 4 NUMBER OF SUBJECTS AFFECTED BASED ON GENDER

Commented [DM18]: Older / geriatric patients

UNDER PEER REVIEW




FIGURE 5 MEDICATION ADHERENCE LEVEL BASED ON MMAS SCORE:

Commented [DM19]: Clarify the numbers & percentages




FIGURE 6 ASSESMENT OF MEDICATION ADHERENCE BASED ON AGE:





FIGURE 7 ASSESMENT OF MEDICATION ADHERENCE BASED ON GENDER



UNDER PEER REVIEW

TABLE 1 OVERALL NUMBER OF GERIATRIC PATIENTS WITH CHRONIC ILLNESSES

CATEGORY	SUB CATEGORY	TOTAL	MALES	FEMALES
Gender		88	35(40%)	53(60%)
Age Group (Years)	50-60	54	16	38
	61-70	28	15	13
	71-80	4	3	1
	Above 80	2	1	1

In this current study, 88 geriatric patients with chronic illnesses were evaluated for their level of adherence to long-term drugs, and the many factors affecting adherence to medication were examined. Out of 88, males were 35(40%) and females were 53(60%). In the age group of 50-60 years -54 people, followed by 61-70 years-28 people, followed by 71-80 years-4 people and above 80 years-2 people were assessed. Out of 88 people, 50-60 years has males-16 & females-38, 61-70 years has males-15 & females-13, 71-80 years has males-3 & female-1 and above 80 years has male-1 & female-1. Out of 88 people, hypertension-14 people, diabetes mellitus-18 people, asthma-4 people, COPD-3 people, psoriasis-1 person, dermatophytosis-10 people, osteoarthritis-28 people and others-10 people. In each disorder, males and females were hypertension:males-6 ,females-8 ,diabetes mellitus:males-11,females-7,asthma:males-1,females-4,COPD:males-1,females-2,psoriasis:males-1,females-0,dermatophytosis:males-6,females-4,osteoarthritis:males-9.females-19 and other:males-4,females-6.

Commented [DM20]: Relate all table to its comment by numbering the table and comment

DISCUSSION:

In the present study 88 subjects of geriatric age group with chronic illness were assessed for the level of compliance for multiple co-morbidities and also various reasons influencing medication compliance were analyzed level of compliance was high in 34 subjects, moderate in 39 subjects and low in 15 subjects. Good compliance was observed in 50-70 year old was 31 and 71-80 was 3. Medium compliance was observed in 50-70 years old was 35 and 71-80 years old was 35 and 71-80 year old was 4. Low compliance was observed in 50-70 year old was 15 and 71-80 year old was 0.

Commented [DM21]: Where is the reason

These observations suggest that the compliance level has decreased progressively with increasing age. It says that the age may be an important reason to affect medication compliance. The missing of medications was mainly attributed to polypharmacy, financial constraints, forgetfulness, lack of time, dependency of care taker, on-availability of medications, and negligence this negligence was reason for high level of non-adherence. The patient's compliance can be improved by reducing the above reasons by giving pill box, medication calendar, simplifying the regime, avoiding the trouble giving adverse events and giving proper motivation of the patients and giving patient counselling about medications and illness. Measuring adherence and patient compliance is quite difficult and is patient dependent most of the time. Medication adherence is difficult in treating chronic diseases and non-adherence among elderly patients is a major issue on health care providers[15-18].

In age 50-60, 16 males and 38 females , in age 61-70, 15 males and 13 females , in age 71-80, 3 males and 1 female and above 80, 1 male and 1 female were affected.

About 38 males and 50 females participated in the study in tertiary care hospital. Among this 4 males and 11 females had low adherence. Then 10 males and 10 females had moderate adherence and then 13 males and 21 females had high adherence.

For Low adherence females were high, for moderate adherence males were high and for high adherence females were high. However, in general, the effect of gender on the rate of adherence to medication in other studies is contradictory. Some researcher found that female patients have better adherence than male patients. While other studies could not find a relationship between gender and adherence to medication[19,20].

CONCLUSION:

This study affirms the presence of unfortunate compliance to the recommended medicine among geriatric subjects with ongoing infections. There are variations in the degree of consistency (poor/moderate/great compliance). Due to the contrast in the review strategies. Adherence to prescription is significant in guaranteeing that restorative advantages are conveyed to patients. However, adherence to drugs has forever been an issue for older. It is critical to keep the patients, Medications medical care providers, healthcare framework, and financial variables as a primary concern. So, we had the option to think of reasonable and individualized answers to defeat these issues[21]. There are portion of the systems to further develop adherence on the grounds that many variables that related to drug adherence and related well-being in more established adults[22].

Statement of informed consent

Informed consent was obtained from all the individual participants included in the study.

REFERENCES:

- 1]Lu J, Zhang N, Mao D, Wang Y, Wang X. How social isolation and loneliness effect medication adherence among elderly with chronic diseases: An integrated theory and validated cross-sectional study. Archives of gerontology and geriatrics. 2020 Sep 1;90:104154.
- 2] Feda MA, Faidah NN, Allam AT, Aloithimen BA, Alsharif AA, Alqarni MA, Saeedi AY, Felemban BN, Awan HM, Alansari MA. Medication adherence among geriatric patients with chronic diseases while living alone. Journal of Pharmaceutical Negative Results. 2022 Nov 4:832-42.
- 3]Fernandez-Lazaro CI, Adams DP, Fernandez-Lazaro D, Garcia-González JM, Caballero-Garcia A, Miron-Canelo JA. Medication adherence and barriers among low-income, uninsured patients with multiple chronic conditions. Research in Social and Administrative Pharmacy. 2019 Jun 1;15(6):744-53.
- 4]Yu H. Universal health insurance coverage for 1.3 billion people: What accounts for China's success?. Health policy. 2015 Sep 1;119(9):1145-52.
- 5] Tøttenborg SS, Lange P, Johnsen SP, Nielsen H, Ingebrigtsen TS, Thomsen RW. Socioeconomic inequalities in adherence to inhaled maintenance medications and clinical prognosis of COPD. Respiratory medicine. 2016 Oct 1;119:160-7.
- 6] Widyakusuma NN, Suryawati S, Wiedyaningsih C, Probosuseno. What Do Seniors Believe About Medication Adherence? A Qualitative Study Among Seniors with Chronic Conditions in Yogyakarta, Indonesia. Patient preference and adherence. 2023 Dec 31:1381-92.

- 7] Zelko E, Klemenc-Ketis Z, Tusek-Bunc K. Medication adherence in elderly with polypharmacy living at home: a systematic review of existing studies. *Materia socio-medica*. 2016 Apr;28(2):129.
- 8] Burnier M, Polychronopoulou E, Wuerzner G. Hypertension and drug adherence in the elderly. *Frontiers in cardiovascular medicine*. 2020 Apr 7;7:49.
- 9] Clark M. *Management Sciences for Health. MDS-3: Managing Access to Medicines and Health Technologies*, Arlington, VA: Management Science for Health Drug Supply.
- 10] Hugtenburg JG, Timmers L, Elders PJ, Vervloet M, van Dijk L. Definitions, variants, and causes of nonadherence with medication: a challenge for tailored interventions. *Patient preference and adherence*. 2013 Jul 10:675-82.
- 11] FW, Wong FY, So WY, Kung K, Wong CK. How much do elders with chronic conditions know about their medications?. *BMC geriatrics*. 2013 Dec;13(1):1-7.
- 12] Ascione FJ, Kirscht JP, Shimp LA. An assessment of different components of patient medication knowledge. *Medical care*. 1986 Nov 1;24(11):1018-28.
- 13] Williford SL, Johnson DF. Impact of pharmacist counseling on medication knowledge and compliance. *Military medicine*. 1995 Nov 1;160(11):561-4.
- 14] Bergman-Evans B. AIDES to improving medication adherence in older adults. *Geriatric Nursing*. 2006 May 1;27(3):174-82.
- 15] Yap AF, Thirumoorthy T, Kwan YH. Systematic review of the barriers affecting medication adherence in older adults. *Geriatrics & gerontology international*. 2016 Oct;16(10):1093-101.
- 16] Chisholm-Burns MA, Spivey CA. The cost of medication nonadherence: consequences we cannot afford to accept. *Journal of the American Pharmacists Association*. 2012 Nov 1;52(6):823-6.
- 17] Osterberg L, Blaschke T. Adherence to medication. *New England journal of medicine*. 2005 Aug 4;353(5):487-97.
- 18] Brundisini F, Vanstone M, Hulan D, DeJean D, Giacomini M. Type 2 diabetes patients' and providers' differing perspectives on medication nonadherence: a qualitative meta-synthesis. *BMC health services research*. 2015 Jun;15:1-23.
- 19] Jin J, Sklar GE, Min Sen Oh V, Chuen Li S. Factors affecting therapeutic compliance: A review from the patient's perspective. *Therapeutics and clinical risk management*. 2008 Feb 29;4(1):269-86.
- 20] Gast A, Mathes T. Medication adherence influencing factors—an (updated) overview of systematic reviews. *Systematic reviews*. 2019 Dec;8:1-7.
- 21] Yap AF, Thirumoorthy T, Kwan YH. Medication adherence in the elderly. *Journal of Clinical Gerontology and Geriatrics*. 2016 Jun 1;7(2):64-7.
- 22] Murray MD, Morrow DG, Weiner M, Tu W, Deer MM, Brater DC, Weinberger M. A conceptual framework to study medication adherence in older adults. *The American journal of geriatric pharmacotherapy*. 2004 Mar 1;2(1):36-43.