

## Case report

# Laparoscopic repair of ileal perforation post blunt abdominal trauma: A Case Report.

### Abstract

Blunt trauma to the abdomen is considered one of the most leading causes of death among all age groups. Delay in the diagnosis of small bowel perforation can result in serious complications and death, making timely diagnosis critical in the treatment of these patients.

We present a male patient aged 35 years old, a case of road traffic accident (RTA), diagnosed as perforated viscus. We did a diagnostic laparoscopy, and laparoscopic repair of 1 cm perforation of the terminal ileum was done. post-operative recovery was smooth as the patient gradually improved and was discharged after 8 days. The patient was reviewed in the surgical outpatient clinic (SOPD) after one and two months with no complaints.

**Keywords:** Blunt abdominal trauma, Ileal perforation, laparoscopic repair, Road traffic accident.

### Introduction

Blunt trauma to the abdomen is considered one of the most leading causes of death among all age groups. Thus, it is an important topic of discussion and research(1). There are many causes of blunt abdominal trauma such as assault, falls from height, and animal attacks. However, road traffic accidents are

As this term appeared only once :[1S]Comment  
in the Abstract, no need to abbreviate it

See the previous comment :[2S]Comment

As this term is frequently found :[3S]Comment  
in the manuscript, the authors can abbreviate it  
with BAT

considered the most common cause of blunt trauma to the abdomen. (2,3). The gastrointestinal perforation diagnosis after blunt abdominal trauma is infrequent as the injury needs to be severe to cause perforation. A hollow viscus injury is not usually suspected unless the clinical condition is highly suggestive (3). A delay in the diagnosis and treatment of the hollow viscus injury results in early peritonitis, and hemodynamic instability leading to increased mortality and morbidity (2). Although small bowel perforation is rare, it is a clinically complex medical condition with high mortality and should be diagnosed and managed early for a better prognosis (4). Small bowel perforation is usually managed by early stabilisation, antibiotic therapy, and surgical source control depending on the case and strong indicators for prompt operative intervention. The mortality rates among patients diagnosed with GIT trauma and small bowel perforation are linked to the delay in initial diagnosis, development of multi-organ failure, and sepsis (5).

I think it could be better to :[4S]Comment  
change this word to "the development of"

Paraphrase this, please :[5S]Comment

On the first mention in the text, :[6S]Comment  
an abbreviation is to be given also in full

### Case report

Here, we present a male patient aged 35 years old, who came to the ER department in an emergency center away from our hospital by about 80 Kilometers at 2:00 am as an RTA case, he was complaining of right-side abdominal pain. He was vitally stable, and his abdominal examination showed right-sided abdominal tenderness. His labs were normal (WBCs 8.5, Hb 12.4). He was managed according to the ATLS protocol. His chest and pelvic X-rays were normal. Pan computed tomography (PAN-CT) (brain, chest abdomen and pelvic) was requested, which revealed minimal pelvic free fluid with mild ascending colon oedema (Figure 1). At 5:00 am, the patient was admitted to our hospital

As this term is not going to :[7S]Comment  
appear in the following paragraphs, there is no need  
to abbreviate it, give it in full

for observation. A serial abdominal examination was started, but the abdominal examination revealed board-like rigidity at 9:00 am. The patient was shifted to the operating theatre (OT) at 9:30 am as an erect chest x-ray revealed massive air under the diaphragm (Figure 2). Diagnostic laparoscopy (DL) was started, and we found intestinal contents inside the peritoneal cavity then running of the small bowel was done and revealed a 1 cm perforation at the ileum, about 120 cm from the ileocaecal junction (Figure3). Laparoscopic repair was done by Vicryl 2/0 in 2 layers and generous peritoneal washing with warm saline and 2 drains were fixed one in the pelvis and one subdiaphragmatic above the liver. post-operative recovery was smooth with some paralytic ileus, and the patient was discharged after 8 days. After 1 month and 2 months, the patient was reviewed in the surgical outpatient clinic, and he has no complaints.

= was complicated by :[8S]Comment






Figure 1: CT abdomen, **NO**pneumoperitoneum was detected.



**FIGURE 2:ERECT CHEST X-RAYshowing bilaterally massive pneumoperitoneum.**



**FIGURE 3: Intraoperative finding of intestinal contents and the perforation of the ileum.**

## Discussion

Because of modern life, developments in industry, various transport systems and violence, there is an increase in the number of patients with blunt trauma to the abdomen (6). The gastrointestinal tract injury following blunt abdominal trauma is frequently undetected, so the diagnosis is usually too late despite advanced imaging techniques such as focused abdominal sonography for trauma and computed tomography scans (7). Apart from our patient, he was diagnosed within hours and was in OT for DL in about 7-8 hours. Traumatic causes of

I think this is to be changed into :[9S]Comment something confirming the above statements like, for example, "Also in the case under discussion, ..." as there was a delay in diagnosis despite CTscan

small bowel perforation can be due to blunt or penetrating injuries (our case after blunt abdominal trauma). The nature of the small intestines being coiled and occupying the largest portion of the abdominal cavity makes them the most injured intra-abdominal hollow organ, particularly from penetrating injuries such as knife stabbing wounds or gunshots (8). Blunt trauma due to traffic accidents is one of the most common causes of small bowel perforation (as our patient), especially in seat-belt users accounting for approximately 75% of gastrointestinal trauma with potential perforation (9). Small bowel perforation can be of two types: free or contained perforation. Free perforations create an open hole that allows the intestinal material to escape into the normally sterile peritoneal cavity and can cause infection, inflammation, and other complications to the surrounding tissues and organs (as you can see in our case). On the other hand, contained perforations are those in which an ulcer creates a full-thickness hole, but the adjacent organs seal it and prevent the free spillage of the intestinal content (10). A high degree of suspicion of intra-abdominal injuries, even in cases following minor trauma will prevent the diagnostic errors. Early diagnosis and effective management of gastrointestinal perforations (as you can see in our case) following blunt trauma to the abdomen helps in decreasing overall morbidity and mortality among these patients (11). Small bowel perforations can present acutely with symptoms ranging from localised abdominal pain to systemic symptoms like fever, nausea, vomiting, and even shock. Additionally, rigidity, guarding, and other peritoneal signs may develop as the condition progresses (12). Although erect plain radiographs are usually the first imaging modality to start with and can detect signs of perforation, they cannot localise the site of perforation (13). The diagnosis of small bowel perforation is usually made by imaging techniques that detect the presence of fluid or gases in the peritoneum or mediastinum. A computed tomography (CT) scan is regarded as the imaging modality of choice for hemodynamically stable patients; however, negative results don't rule out the diagnosis of small bowel perforation (14) (as our patient, NO pneumoperitoneum in the abdominal computed tomography). In the EAST trial, 13% of the study samples who were discovered to have a small bowel perforation at laparotomy had no positive findings before surgery (15). Delays in the diagnosis of small bowel perforation can result in serious complications and death, making timely diagnosis critical in the treatment of these patients

= intestine : [10S] Comment

=GIT : [11S] Comment

was detected : [12S] Comment

(16). In trauma patients, laparoscopic interventions with normal haemodynamic parameters are an excellent modality to identify any abdominal injuries. Laparoscopy is more efficient and cost-effective and associated with fewer complications in comparison with traditional laparotomies (17). Although there are different treatments for the perforation of the small bowel at various locations, there are still no established and reliable therapeutic recommendations for small bowel perforation. In the management of small bowel perforation, conservative treatment is not favoured (18). Laparoscopy and open surgery (primary repair, anastomosis, and stoma) are mainly used, which may be related to timely preoperative management, good resuscitation of septic shock, and timely application of broad-spectrum antibiotics in the perioperative period (19). The mortality rates among patients diagnosed with GIT trauma and small bowel perforation are linked to the delay in initial diagnosis, development of multi-organ failure, and sepsis as emphasized by the **EAST trial**, which reported a rise in mortality rate from 4% to 16% when the diagnosis is delayed for more than 24h. It was further highlighted that when small bowel perforation was present the mortality rate rose from 14% to 19% (9).

## ETHICAL APPROVAL

As per international standards or university standards, written ethical approval has been collected and preserved by the author(s).

## CONSENT

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

DISCLAIMER : **NO generative AI technologies were used.**

## References

1. **Dauterive AH, Flancbaum L, and Cox EF.** Blunt Intestinal Trauma A Modern-day Review. *Ann Surg J.* 1985;201(02):198-203.
2. **Hughes TMD and Elton C.** The Pathophysiology and Management of Bowel and Mesenteric Injuries Due to Blunt Trauma. *Ann Surg J.* 2002;33:120-8.

I need to comment this : **[13S]Comment** statement of the author(s). First, small bowel perforation (both trauma-induced and caused by a pathological condition) is a highly-discussed topic in surgical society, with time-proven and quite reliable recommendations for the management clearly expressed in the literature. The other thing is that with science and technology development the treatment options have been constantly modified. The management becomes more selective and less invasive aiming to avoid non-therapeutic laparotomies and at the same time to reduce the risk of missed injuries. Second, the issue of the recommendations for the treatment is out of scope of the article presented, and in my opinion, it would be better to leave it for further large-scale research.

Is it a relevant reference, as : **[14S]Comment** the article cited (written in German) describes the management of intestinal obstruction?

3. **Kurane SB and Ugane SP.** A clinical study of hollow viscus injury due to blunt trauma abdomen. *Int J Res Med Sci.* 2017;5(11):5017.
4. **Onken F, Senne M, Königsrainer A, and Wichmann D:** Classification and treatment algorithm of small bowel perforations based on a ten-year retrospective analysis. *J Clin Med.* 2022, 11:5748. 10.3390/jcm11195748
5. **Fakhry SM, Allawi A, Ferguson PL, Michetti CP, Newcomb AB, Liu C, and Brownstein MR:** Blunt small bowel perforation (SBP): an Eastern Association for the Surgery of Trauma multicenter update 15 years later. *J Trauma Acute Care Surg.* 2019, 86:642-650.
6. **Bouzat P, Valdenaire G, Gauss T, Charbit J, Arvieux C, Balandraud P, Bobbia X, David JS, Frandon J, Garrigue D, and Long JA.** Early management of severe abdominal trauma. *Anaesthesia Critical Care & Pain Medicine.* 2020 Apr 1;39(2):269-77.
7. **Yavaş Ö, Çelik A, Altuntaş G, Altuntaş M, Bilir Ö, and Ersunan G.** The role of repeated extended FAST in patients with stable blunt thoracoabdominal trauma. *Turkish Journal of Trauma & Emergency Surgery/Ulusal Travmatoloji Cerrahi Dergisi.* 2023 May 1;29(5).
8. **Aljehaiman F, Almalki FJ, Alhusain A, Alsalamah F, Alzahrani K, Alharbi A, and Alkhulaiwi H.** Prevalence, Pattern, Mortality, and Morbidity of Traumatic Small Bowel Perforation at King Abdulaziz Medical City: A Retrospective Cohort Study. *Cureus.* 2024 Jan 15;16(1):e52313.
9. **Staab V, Naganathan S, McGuire M, Pinto JM, and Pall H.** Gastrointestinal Perforation with Blunt Abdominal Trauma in Children. *Children (Basel).* 2024 May 21;11(6):612.
10. **Hafner J, Tuma F, Hoilat GJ, and Marar O.** Intestinal perforation. *InStatPearls [Internet]* 2023 Aug 8. StatPearls Publishing.
11. **Gurav SP, Quraishi AH, Umare G, and Tongse PS.** A prospective observational study of gastrointestinal perforations following blunt trauma abdomen. *International Surgery Journal.* 2024 Apr 29;11(5):737-743.
12. **Katwal S, Ghimire A, and Joshi N.** Uncommon presentation: isolated small bowel perforation after blunt abdominal trauma in a pediatric patient; a case report. *Annals of Medicine and Surgery.* 2023 Dec 1;85(12):6222-6226.
13. **Kulinna-Cosentini C, Hodge JC, and Ba-Ssalamah A.** The role of radiology in diagnosing gastrointestinal tract perforation. *Best Practice & Research Clinical Gastroenterology.* 2024 Jun 7:101928.

14. **Bini R, Ronchetta C, Picotto S, Scozzari G, Gupta S, Frassini S, and Chiara O.** Importance of CT-scan predicting clinical outcomes in gastro-intestinal perforation. *Annals of Translational Medicine.* 2020 Nov;8(21).
15. **Loftus TJ, Morrow ML, Lottenberg L, Rosenthal MD, Croft CA, Smith RS, Moore FA, Brakenridge SC, Borrego R, Efron PA, and Mohr AM.** Occult bowel injury after blunt abdominal trauma. *Am J Surg.* 2019 Aug;218(2):266-270.
16. **Fakhry SM, Allawi A, Ferguson PL, Michetti CP, Newcomb AB, Liu C, and Brownstein MR.** Blunt small bowel perforation (SBP): an Eastern Association for the Surgery of Trauma multicenter update 15 years later. *Journal of Trauma and Acute Care Surgery.* 2019 Apr 1;86(4):642-650.
17. **EI-Bendary YB, Al-Adawi J, and Al-Qadhi H.** The Use of Laparoscopy in the Management of Trauma Patients: Brief review. *Sultan Qaboos Univ Med J.* 2016 Feb;16(1):e9-e14.
18. **Listle H, and Gutt C.** Conservative and surgical ileus treatment. *Chirurg* 2017;88:629-644.
19. **Li Z, Li S, Wu C, Liu Z, Zhao Y, Luan J, Miao S, and Wang P.** Diagnostic and therapeutic procedures of small bowel perforation: A retrospective monocentric cohort study. *World Journal of Surgical Infection.* 2023 Jan 1;2(1):7-11.