

Original Research Article

Study of the resurgence of typhoid fever in the commune of doba in Chad

ABSTRACT

Aims: The aim of this study was to determine the factors behind the increase in the number of cases of typhoid fever in the commune of Doba.

Place and Duration of Study: Our study was carried out in the District and Provincial hospital and the Christian Assemblies in Chad (ACT) health center of Doba, from July to August 2022.

Methodology: A total of 756 Widal and Félix tests according to the slide agglutination technique described by Raobijaona and Ranaivo-Harisoa were carried out in three health facilities in order to confirm the reliability of the test within these facilities. Surveys, using survey forms, were carried out in 120 households, with 95 households participating in our study.

Results: The results showed a fairly high typhoid fever prevalence rate (72.52%), with all households surveyed (100%) using self-medication antibiotics. We also noted a high frequency of antibiotic use without a doctor's prescription. Non-compliance with hygiene and sanitation rules was noted in the majority of households (80%), which ate food without carefully washing their hands with soap and water. Most of the households surveyed (52%) had no toilet, which encourages open defecation.

Conclusion: Despite its unreliability, the Widal and Félix serodiagnosis is the test frequently requested and carried out. Failure to comply with hygiene measures and excessive self-medication, at the root of the selection of resistant bacteria, have been partly identified as factors that have contributed to the increase in the number of cases of typhoid fever in Doba.

Keywords: typhoid fever, Widal et Félix, antibiotic resistance, Salmonella, hygiene and sanitation.

1. INTRODUCTION

Typhoid fever is an acute infectious disease caused by ingestion of Eberth's bacillus, *Salmonella typhi* or, more rarely, *S. paratyphi* A, B or even C. Transmission is essentially fecal-oral, and can be direct from human to human via dirty hands or indirect via drinking water and various foods contaminated with faeces [1].

It is fatal and febrile, with young people at greater risk of contracting the disease. Without treatment, its case-fatality rate is 10-30%, falling to 1-4% with appropriate therapy [2]. Common symptoms include long-lasting fever, chills and abdominal pain. On the other hand, a non-specific symptom profile complicates the clinical diagnosis, with symptoms common to other diseases occurring in typhoid fever endemic areas. The mainstay of laboratory confirmation of cases remains blood culture, with a limited sensitivity of around 40-60% [3], partly due to widespread self-medication prior to visiting a health centre.

It is estimated that there are between 11 and 21 millions cases of typhoid fever, with between 128,000 and 161,000 deaths, and 6 millions cases of paratyphoid fever, with 54,000 deaths annually [4]. The majority of cases occur in South-East and South Asia and sub-Saharan Africa.

While the incidence of typhoid fever has fallen sharply in countries with a high standard of living, to less than 1 case per 100,000 inhabitants, it remains high in developing countries, at 130 cases. Typhoid fever is endemic in the tropics and the Third World, with epidemics of varying severity [5].

The emergence of antimicrobial resistance is a significant challenge, with several recent major epidemics caused by multi-antibiotic-resistant *S. Typhi* in Africa and Asia [6].

In Chad, typhoid fever is a public health problem that needs to be addressed in view of the high demand for research into this pathology in the relevant departments within hospitals. In 2017, reference consultation services notified 25,946 probable NC [7]. Adults aged 15 and over (302 NC/100,000 adults) are 7 times more affected than children aged 5-14 (42 NC/100,000). Infants are less affected than other age groups [7].

In Logone Oriental province, the typhoid fever detection rate was 39% in 2017 [7]. Today, there is a considerable increase in cases of typhoid fever in the province of Logone Oriental, particularly in the commune of Doba. This worrying situation should be of concern to all those involved in health, as well as to the population itself. To this end, a better understanding of the epidemiological situation of *Salmonella Typhi* and *Paratyphi* in the province of Logone Orientale and in the commune of Doba in particular, as well as the causes that contribute to it, remains essential for appropriate control of this disease.

2. MATERIAL AND METHODS

2.1. Ethical considerations.

This study was approved by the National health policy 2016-2030 [8].

2.2. Study area

The study was carried out in the commune of Doba, capital of the Province of Logone Oriental. It is situated between 8.666° North Latitude and 16.850° East Longitude. The commune of Doba is divided into 4 arrondissements with 24 districts. It also has two (02) hospitals and three (03) health centres for a population of 106,403 inhabitants, spread over 15,200 concessions [9].




Figure 1 :Location map of the study area

2.3. Sampling

Our work involved clinical tests and surveys using survey forms between July and August 2022, a period of high demand for Widal and Félix examinations. It involved both hospitals and households.

2.3.1. In households

A total of 120 households were selected on a voluntary basis to take part in the survey. All arrondissements were covered, with the majority of households concentrated in the 4th arrondissement.

Within the concessions, the survey targeted mothers or guardians of children, i.e. the heads of household. And if there were several households in the concession, in each household the woman or the child's guardian, i.e. the head of the household, was asked to answer the questionnaire. On the other hand, if there were several women in the same household, only one should be interviewed.

2.3.2. In health facilities

Patients diagnosed with typhoid fever and referred for the Widal and Félix tests were systematically collated. Subjects were identified in two (02) hospitals (the District hospital and the Provincial hospital) and one (01) health center (the Christian Assemblies in Chad (ACT) health center) in the commune.

It should be noted that our survey covered adults and children, regardless of sex, with an age range of between 85 and 1 years.

Table I shows the variables defined for the study.

Table I : Variables studied

Variable	Type	Collection technique
Age	Quantitative	Interview
Sex	Qualitative	Direct observation
Ethnicity	Qualitative	Interview
Profession	Qualitative	Interview
Duration of symptoms	Quantitative	Interview
Result of Widal and Félix serodiagnosis	Qualitative	Direct observation
Result of control test	Qualitative	Direct observation

2.3.3. Sample size

The minimum sample size was calculated using the following formula :

$N = [Z^2 \times p \times (1-P)] / M^2$, where : N = required sample size ; Z = 95% confidence level (standard value of 1.96) ; P = estimated prevalence of typhoid fever cases between July and August 2022 in the commune of Doba ; M = 5% margin of error (standard value of 0.05).

The prevalence (P) of typhoid fever between July and August 2022 was :

$P = (\text{Number of positive cases} / \text{Number of suspected cases}) \times 100$

2.4. Widal and Félix serodiagnosis

2.4.1. Test rationale

The choice of this test is based on its multiplicity of tests and its use as a mass screening and peripheral diagnostic test. Its performance compared with the reference serological test should also be noted.

2.4.1.1. Performance of the test

A control test was used to check the reliability of the examinations carried out in the health facilities concerned. It consisted of collecting sera that had already been tested at the health centres in order to confirm its reliability.

For our study, 37 sera stored in dry tubes and closed at -20°C or at lower temperatures were thawed and homogenised before being tested on 746 samples.

At the Assemblées Chrétiennes health centre in Chad, the slide agglutination technique described by Raobijaona and Ranaivo-Harisoa (2000) was used. To a drop of serum corresponding to 50µl on a clean slide, a drop of antigenic suspension O or H (2.5µl) of salmonella T, A, B or C respectively is added. The mixture obtained is then homogenised on the plate and placed on a rotary shaker at 100 rpm for 3 minutes. Positive reactions result in lumpy agglutination of the mixture containing the antigen.

At the district hospital, on the other hand, we used the agglutination technique in tubes after dilution 1:10. To 900µl of distilled water, 100µl of serum was added and then 1/100 of the mixture was taken for each O and H tube. Next, 900µl of the antigenic suspensions O or H of Salmonella T, A, B or C respectively was added to the corresponding tubes and centrifuged for 5 minutes at 3000rpm. The positive tubes showed clumpy agglutination for the O antigen and dusty smoke for the H antigen, observable under light after gentle shaking of the tube.

2.4.1.2. Test evaluation criteria

- Internal validity criteria

Sensitivity and specificity attest to the intrinsic validity of diagnostic tests. In the case of typhoid and paratyphoid fevers, our analysis focused on the validity of the Widal et Félix test or plate agglutination test.

The reference diagnostic test used for our study is the dilution agglutination test, described as a control test [10].

Sensitivity :the sensitivity of the test defines its ability to correctly identify individuals carrying the disease. It correspond to the ratio of the number of subjects recognised as ill by the plate agglutination test to the total number of ill subjects (identified by the reference test). It was determined using the formula : $\text{Sensitivity} = \text{VP} / (\text{VP} + \text{FN})$

where : VP = True Positives (positive with the reference test and the plate agglutination test) and FN = False Negatives (positive with the reference test but negative with the plate agglutination test).

Specificity :specificity is the ability of the test to correctly identify individuals who are not carriers of the disease.

The formula used: $\text{Specificity} = \text{VN} / (\text{FP} + \text{VP})$

with : VN = True Negatives (negative with the reference test and the plate agglutination test); FP = False Positives (positive with the agglutination test but negative with the reference test), corresponds to the ratio of the number of subjects recognised as not sick by the plate agglutination test to the total number of subjects recognised as not carrying the disease (identified by the reference test).

- External validity criteria

The Positive Predictive Value (PPV) indicates the probability of a subject actually being ill when the test is positive. It is a function of the sensitivity, specificity and prevalence of the disease. It is calculated using the formula :

$$\text{PPV} = (\text{Preval.} \times \text{Sensitivity}) / [(\text{Preval.} \times \text{Specificity})] + [(1 - \text{Preval.}) \times (1 - \text{Specificity})]$$

The Negative Predictive Value (NPV) is the probability that a subject is actually not ill when the test gives a negative result. It depends on the sensitivity, specificity and prevalence of the disease and is calculated by the formula :

$$\text{NPV} = ((1 - \text{Preval}) \times (\text{Specificity})) / [(1 - \text{Preval}) \times (\text{Specificity})] + [(\text{Preval}) \times (1 - \text{Sensitivity})]$$

The positivity threshold, an arbitrary variable that separates positive results from negative results, is characterised in our study by the observation of the presence of agglutination on the slide.

2.5. Characterisation of behaviours favouring the transmission of germs responsible for typhoid fever in the population

Direct observation of the level of hygiene and sanitation in the various households surveyed was used to support or reject the information collected from patients diagnosed in hospital facilities.

The various points covered in the interview focused mainly on the state of health of the children, in particular the presence of diarrhoeal diseases, the supply of drinking water and the overall level of individual and collective hygiene within the household.

To this end, a direct observation guide was used in the various neighbourhoods.

2.6. Identification of practices favouring the selection of resistant bacteria within the population

The literature on the use of antibiotics in human populations was used to identify the risk factors that could be involved in the emergence of resistant bacteria responsible for typhoid fever. For practical reasons, the study was limited to two (02) main risk factors for which further details are required for their analysis. These were the presence of self-medication and non-compliance.

The survey was conducted among randomly selected residents of the commune of Doba, whether or not they attended the consultation.

2.7. Statistical analysis

We used a 2-stage random sampling method (health facilities (hospitals and health centres) and households).

The database was created and managed using Access software (Microsoft office corporation 2019). Quantitative data were entered using Excel (Microsoft office corporation 2019) and analysed using Statistical Package for the Social Sciences 17.0 (SPSS).

3. RESULTS AND DISCUSSION

3.1. Survey results

Of the 120 households identified, only 95 gave their consent to participate in our study and to answer the proposed questionnaire. This represents a participation rate of 79.16%. The remaining 25 households (20.84%) categorically refused to participate.

3.2. Typhoid fever prevalence rate

From July to August 2022, 746 patients suspected of having typhoid fever were diagnosed. Of these, 541 tested positive for the Widal et Félix serology, giving a prevalence rate of 72.52%. These results are shown in table II below.

Table II : Typhoid fever prevalence rate

Periods	Suspected cases	Number of positive cases	Percentage (%)	Negative cases	Percentage (%)
July	377	293	77.72%	84	22.28%
August	369	248	67.21%	121	32.79%
Total	746	541	72.52%	205	27.48%
Value of P			0,349		

Our results show a high incidence of typhoid fever, which is similar to those of Sauch *et al.* (2010), who found an annual distribution revealing a summer peak from July to October, in relation to rainfall. Two phenomena could explain this high incidence of typhoid fever in the hot, rainy season. Firstly, the rainy season is marked by major problems with drinking water supplies and sanitation. On the other hand, so-called 'water-borne' diseases, including typhoid fever, are rife at this time of year [1, 11]. Our results are in line with those obtained in 2002 by Sako in Yopougou, Abidjan, and Lefebvre in Senegal.

3.3. Patient characteristics

Age (Table III) and sex (Table IV) were used to characterise the patients diagnosed.

The presence of salmonella was observed in all the age groups identified (Table V), with the majority of cases observed after the incubation period (Table VI).

Table III : Typhoid fever prevalence rate according to age

Age groups	Effectif			
	Positive	Percentage	Négative	Percentage
1 to 10 years	146	73,73%	52	26,27%
11 to 40 years	298	72,68%	112	27,32%
41 years and over	97	70,28%	41	29,72%
P - Value			0,039	

The age group between 11 and 40 was the most represented, with a frequency of 55.08%, and an average age of 25. The minimum age was 1 and the maximum 85. The incidence of typhoid fever is much higher in young adults. This could be explained by the fact that older children and adults can become infected outside the family environment.

Table IV shows that women are more affected than men, with a frequency of 58.96%. The work of Theveniau *et al.* (2000) in France and Teyssier (2000) in Senegal showed a male predominance, which is contrary to our results. This is contrary to our results. There seems to be no explanation for these differences.

Table IV:Prevalence rate of typhoid fever according to sex

Widal and Félix serodiagnosis	Gender	
	Male	Female
Positive	222 (41,03%)	319 (58,96%)
Negative	94 (45,85%)	111 (54,14%)
P value	0,067	

As shown in Table V, all the salmonella serotypes responsible for typhoid fever were found. However, there was a predominance of *S. Typhi* strains (44.17%), confirmed by the Widal et Félix serodiagnosis. This is the only confirmatory test for typhoid fever carried out in hospitals and health centers in the commune of Doba.

Table V : Distribution of Salmonella serotypes identified

Sérotypes	Number positive	Percentage
<i>S. Typhi</i>	239	44,17%
<i>S. Paratyphi C</i>	49	9,05%
<i>S. Paratyphi B</i>	116	21,44%
<i>S. Paratyphi A</i>	137	25,32%

The data in Table VI show the reliability of the Widal et Félix test with advanced cases (73.19%) of typhoid fever.

Tables VI : Typhoid fever prevalence rate by incubation period

Widal and Félix serodiagnosis	Less than 8 days	More than 8 days	Total
Positif	145 (26,80%)	396 (73,19%)	541 (100%)

3.4. Practices favoring the selection of resistant bacteria in the population

The results in table VII show the practices that encourage the selection of resistant bacteria. They reveal the use of self-medication antibiotics in all households surveyed (100%). In 11 (11.58%) households, antibiotics are used without a doctor's prescription, due to a lack of means, and in 84 (88.42%), due to experience or habit.

We can thus note a high frequency in the use of antibiotics without a doctor's prescription in view of the percentage of households systematically resorting to it (56%) in the presence of a case of illness, compared with 44% referring to a doctor. For 88.42% of households, the main reasons for self-medication are related to experience, to which can be added the economic factor for the remaining households (11.58%).

Tables VII :Risky practices in the selection of resistant bacteria

Practices	Households	
	Yes	No
Practice of self-medication	95 (100%)	0 (0%)
Systematic self-medication in the event of illness	56 (59%)	39 (41%)
Antibiotic use by habit / experience	84 (88,42%)	11 (11,58%)
Choice of antibiotic by habit / experience	58 (61%)	37 (39%)
Referral to a doctor after failure to self-medicate	38 (40%)	57 (60%)
Compliance with doctor's prescription	27 (28,42%)	68 (71,58%)
Discontinuation of treatment after improvement in state of health	85 (89,47%)	10 (10,53%)

Our surveys also revealed a reliance on habit and experience in the choice of antibiotic molecules. This practice can be a significant risk factor, resulting in antibiotic treatments that are inappropriate, ineffective or even useless in the face of the pathogens present. Such as unsuitable spectra, non-bacterial microbial agents and/or ill-defined dosages [6, 12, 13].

It is highly regrettable to note that in our study, 60% (n=57) of households implement a second treatment before seeking medical attention. Indeed, any use of antibiotics again exposes bacteria to selection pressure [14, 15] and the multiplicity of treatments is a risk for the selection of multi-resistant bacteria [16].

Our study also shows that 28.42% of patients adhere to their doctor's prescriptions, as opposed to 71.58% who, due to lack of resources, fall back on lower-cost generic antibiotic molecules and, once their health has improved, stop treatment or modify the dosage and frequency of antibiotic administration. It should not be forgotten that changes in antibiotic dosage, duration or frequency of administration are risk factors for the emergence of resistant bacteria. It can also lead to a lack of bacteriological cure for the patient, and be responsible for chronic carriage of virulent bacteria [14, 17].

Our surveys revealed, among other things, abundant use of ciprofloxacin with a proportion of 56.4%, followed by Ceftriaxone (31.4%). According to the information gathered, these antibiotics are used because of their broad spectrum of action, availability and easy administration.

3.5. Behaviours favouring the transmission of the typhoid fever germ within the population

We noted that very few households (11.58%) regularly used soap to wash their hands before eating, which is well below the rate of households (80%) simply using water without soap, but higher than that of households (8.42%) using it occasionally (table VIII).

Tables VIII :Tools and frequency of hand disinfection before eating

Hand disinfection tools and frequency	Workforce	Percentages
Plain water	76	80%
Water with soap always	11	11.58%
Water +Soap occasionally	08	8.42%
P-value		0,057

This situation calls for a wide-ranging information and awareness campaign to link the level of hand hygiene in particular with the onset of diarrheal diseases such as salmonellosis among the population.

The main sources of household water supply in the commune of Doba during the rainy season are wells (80%), pumps or boreholes (33.68%) and taps (45.26%) (table IX). In the dry season, water comes from traditional wells and pumps.

Table IX: Hygiene and sanitation factors in households

Hygiene and sanitation	Existing	Percentage	Non existing	Percentage
Wells	76	80%	19	20%
Latrines	43	45.03%	52	54.73%
Garbage cans	13	13.68%	82	86.32%
Sumps	0	0%	0	0%
Boreholes	32	33.68%	63	66.32%
Taps	43	45.26%	52	54.74%

Traditional wells are the primary source of drinking water in both wet and dry seasons. This water is very often consumed directly in households, without prior treatment. It is only treated in 29 households (38.15) and bleached in 30 (39.47%).

The conditions under which drinking water is collected, transported, stored and handled can contribute to contamination at several levels: drawing water, buckets and ropes lying on the ground, the container or transport vessel, the cup used to draw water from the canary, and poor maintenance of the cup [18].

To circumvent these sources of contamination, containers should be covered for transport and home storage. These containers must also be kept out of the reach of children, pets and insects. Water should be removed from storage containers by siphoning, rather than using cups, which are often exposed to insects and dust [19].

The length of time water is stored in the home seems to be a major factor in its quality deterioration.

quality deterioration. It can be seen that the longer the water is stored, the greater the deterioration in water quality. After three days, over 80% of water samples analyzed were of poor quality [20]. Similarly, there is a low presence of latrines (48%) in households, which would favour the dissemination of germs in the wild, a major factor in the spread of typhoid fever [1, 11] in the commune.

4. CONCLUSION

In sum, our study shows that typhoid fever has a high incidence, with a prevalence rate of 72.52%. Although known to the population (especially in urban areas), this disease continues to cause public health problems in Chad, and particularly in the commune of Doba, the resolution of which requires multi-sectoral collaboration for epidemic control, prevention and vaccination. Despite its unreliability, the Widal et Félix serodiagnosis is the most frequently requested test performed by our practitioners. Failure to comply with hygiene measures and excessive self-medication, which are at the root of the selection of resistant bacteria, are partly identified in this study as factors that have contributed to the increase in the number of typhoid fever cases in the commune of Doba.

REFERENCES

1. Mbula, W., Odio, K., Kashongwe, M. & Mizero. (1993). Epidemiological aspects of typhoid fever in Kinshasa. About 208 observations. *Med. D'Afrique Noire*, 40 (11) 673-678.
2. World Health Organization (WHO). (2018). Surveillance standards for vaccine-preventable diseases. Geneva: World Health Organization. Last updated on September 5, 15p.
3. Parry, CM., Wijedoru, L., Arjyal, A. & Baker, S. (2011). The utility of diagnostic tests for enteric fever in endemic locations. *Expert Rev Anti Infect Ther.*, 9(6):711-25. <https://doi.org/10.1586/eri.11.47>.
4. GBD. (2020). Occupational Risk Factors Collaborators. Global and regional burden of disease and injury in 2016 arising from occupational exposures: a systematic analysis for the Global Burden of Disease Study 2016. *Occup Environ Med.*, 77(3):133-141. <https://doi.org/10.1136/oemed-2019-106008>. PMID: 32054817; PMCID: PMC7035694.

5. Kernbaum, S. (1988). Elements of infectious pathology. Edition. SIMEP/Specia., pp. 213-215.
6. ANSES. (2014). Assessment of the risks of emergence of antibiotic resistance related to the use patterns of antibiotics in the field of animal health. Report of the collective expertise Referral No. 2011-SA-0071, 218p.
7. Ministry of Public Health (MSP). (2017). General report on infectious diseases (F.T), 183p.
8. Minister of Public Health (MSP). (2016). National health policy 2016---2030. 45p.
9. Ministry of Planning and Development (MEP). (2009). Second population and housing census (RGPH2): Overall results. 88p.
10. Raobijaona, A. & Ranaivo-Harisoa, HA. (2000). Typhoid fever in children in Antananarivo. Black African Medicine, 47 (5): 244-246. <https://api.semanticscholar.org/CorpusID:58705361>.
11. Essayagh, M., Rhaffouli, AE., Essayagh, S. & Essayagh, T. (2020). Epidemiology profil of fever typhoid in Meknes (Morocco) 2013-2016. Rev Epidemiol Sante Publique, 68(1):45-49. French. <https://doi.org/10.1016/j.respe.2019.09.006>. Epub 2020 Jan 6. PMID : 31917023.
12. Chardon, H. & Brugere, H., (2014). Use of antibiotics in livestock farming and meat industries. Animal Health and Safety Notebooks of the Meat Information Center, 36p.
13. Guienne, V., Marquis, C., David, M., Fleuret, S. & D'halluin, E. (2017). Self-medication in question: a socially and territorially situated DIY. International and Multidisciplinary Conference of Nantes, 306p.
14. Sanders, P., Bousquet-Melou, A. & Chauvin, C. (2011). Toutain P, Use of antibiotics in livestock farming and public health issues. INRAE Animal Productions, 24(2), 199–204. <https://doi.org/10.20870/animal-productions.2011.24.2.3254>
15. Sanders, P., Perrin-Guyomard, A. & Moulin, G. (2017). Evolution of the use of antibiotics in animal production. Cahiers de nutrition et de diététique, 11p.
16. Kathleen, V. (2019). Use of antibiotics in dairy cattle farming: survey of farmers on risky practices. Veterinary doctoral thesis, Université Claude-Bernard - Lyon I. 65p.

17. Haenni, M., Jouy, E., Madec, J-Y. & Laurent, F. (2012). Methicillin-resistant *Staphylococcus aureus* (MRSA): shared between humans and animals? *Epidemiological, animal health and food bulletin* (n°53) Special Antibiotics and Antibiotic Resistance, pp. 40-43.
18. Degbey, C. (2011). Factors associated with the problem of drinking water quality and the health of populations in the commune of Abomey-Calavi in Benin. Doctoral thesis in Public Health Sciences. Université Libre de Bruxelles (ULB). General, 112p.
19. Grondin, PM. (2005). Chlorination in rural areas in developing countries. Proceedings of the meeting organized by pS-Eau. Water Solidarity Program. 99 p.
20. Ngnikam, E., Mougoue, B. & Tietche, F. (2007). Water, sanitation and impact on health: case study of an urban ecosystem in Yaoundé. Proceedings of the JSIRAUF, Hanoi (Vietnam), 13p.