

Original Research Article

Impact of Irrigation Regimes and Organic Amendments on Soil Physical Properties, Nutrient Availability, and Productivity in Calcareous Soil

ABSTRACT

The scarcity of irrigation water in Egypt requires addressing severe water scarcity, such as deficit irrigation and using organic amendments to improve water use efficiency in agriculture. A field experiment was conducted at El-Nubaria Agricultural Research Station in the summer of 2022 for maize and the winter of 2022/2023 for wheat to study the impact of organic amendments and irrigation regimes on calcareous soil properties and productivity. The experiments involved a split plot design with three replicates, irrigation regimes were placed in main plots (50% (I_{50}), 75% (I_{75}), and 100% (I_{100}) of ET_c) and organic amendments in subplots included five treatments: 1) 9.6ton ha^{-1} (B_1), 2) 14.4 ton ha^{-1} (B_2) of biochar, 3) potassium humate at the rate of 1.5% (KH_1), 4) 3% (KH_2) and 5) compost (12 ton ha^{-1}). The obtained results indicated that compost, followed by B_2 (14.4 ton ha^{-1}), achieved the highest soil organic matter value. Applied I_{100} had the highest soil available NPK and micronutrients, followed by I_{75} . Organic amendments significantly affected the soil availability of N, P, K, Fe, Mn and Zn. The maximum potassium content in grains was achieved using I_{100} , followed by I_{75} . The influence of organic amendments on field capacity and available water took the descending order: $B_1 > KH_1 > \text{compost}$. Water use efficiency was negatively impacted by increasing the amount of applied irrigation water. Applying I_{100} resulted in a significant increment in wheat and maize grain yields, followed by I_{75} . The influence of organic amendments on wheat grain yield took the descending order: $\text{compost} > \text{potassium humate} > \text{biochar}$. The effect on maize grain yield took the descending order: $\text{potassium humate} > \text{compost} > \text{biochar}$.

1. INTRODUCTION

Considering over 80% of Egypt's water sources are used for agriculture, the availability of water is the greatest obstacle to growing the country's food production. The increase in irrigated areas, however, is increasing the demand for water sources. Since irrigation water resources are scarce, great efforts should be made to rationalize the use of each unit of water. One of the major tactics to save water and improve water use efficiency is irrigating the crops with a water amount less than the optimal requirement, which refers to water deficit. However, the application of a deficit water pattern in irrigating crops is associated with low yields and quality [1]. Physiologically, low water supply, as an abiotic stress, causes critical issues in cell water integrity, stomata performance, plant pigments, and

photosynthesis, hence a reduction in productivity is anticipated [2]. Therefore, several practices are required to be applied under water shortage to maintain better plant growth and productivity [3]. Maize (*Zeamays L.*) is an important food and feed crop that ranks third in the world, with multiple uses such as human food, animal feed, and industrial raw materials [4].

Wheat (*Triticum aestivum L.*) is a major cereal crop in Egypt, with a planted area of 1.43 million hectares. The total production was 9.3 million tons, with an average of 6.5 tons per hectare [5]. Another important cereal crop is wheat, which grains contain a range of protein (6–21%), fats (1.5–2.0%), cellulose (2.0–2.5%), minerals (1.8%), and vitamins (1.8%), according to [6]. Egypt is the world's largest importer of wheat, providing 40% of the population's needs, with 10 million tons produced annually from 1.5 million hectares [7]. However, growing wheat in calcareous soils under deficit irrigation resulted in noticeable decreases in yield and quality. In saline or calcareous soils, crop plants are subjected to physiological drought, which negatively alters the physiological status and nutrient balance, suppressing growth. Organic soil amendments enhance soil properties, maintain fertility, increase productivity, preserve the environment, and produce safe organic food [8]. Organic materials improve the chemical and physical properties of sandy soil by retaining moisture and recycling nutrients [9]. Organic additions enhance soil quality and increase carbon storage, potentially aiding in climate change mitigation efforts [10]. Soil amendments, when properly prepared and applied, can enhance plant productivity, and soil stability [11]. They cause increased population sizes and proportions of bacteria associated with nutrient cycling [12].

Compost is a fully decomposed, stabilized, homogeneous substance of animal or plant origin, free from harmful elements that could harm humans, animals, plants, or the environment [13]. It is a humic-rich amendment that enhances soil properties through the controlled biological decomposition of organic resources [14, 15]. Compost can enhance the growth of plants, lower the need for fertilizer, and reduce soil-borne plant diseases [16]. Composting is an environmentally friendly method of converting organic waste into organic nutrients [17].

Biochar is a porous, high-carbon material produced through pyrolysis in a no-oxygen environment, which serves as a storage medium for carbon [18, 19] and raises aromatic structures and essential minerals in clay minerals over time [20]. Biochar is gaining popularity as a soil conditioner due to its carbon storage, soil fertility enhancement, water retention, and increased crop yield [21, 22, 23]. Also, the use of biochar in agricultural soils offers numerous benefits including improved soil health, nutrient retention, efficient transfer, and climate change mitigation. Its high surface area, porosity, and high levels of COO-, OH-, P-O-, and R-OH groups contribute to its benefits [24]. Furthermore, biochar enhances soil properties, relocates fertilizers, and sequesters carbon, while maintaining stability is crucial for climate change mitigation [25]. Biochar in alkaline soils was an effective, reasonable, and environmentally friendly approach [26]. Compost and biochar can be used to improve soil quality and productivity by absorbing nutrients, improving nutrient availability, and stimulating the microbial. Biochar-blended compost can act as a partial replacement for mineral fertilizers [27, 28]. Also, Biochar and compost can enhance maize yield and soil quality, with their combined application potentially enhancing the properties of soil [29, 30].

Humic substances (HS) promote plant growth through anatomical and biochemical changes in the root system via chelation [31]. Humic acid, a plant growth regulator, is broken down by living microorganisms, allowing plants with strong chemical and biological activity to easily absorb it [32]. HS comprising 60% of soil organic compounds are crucial for the agroecosystem and are responsible for various complex chemical reactions [33]. K-humate is more efficient than K-Si and compost due to its high solubility, smaller dose, and ease of

field application [34]. HS are commonly used in agriculture due to their hormone-like effects, which promote growth, enhance plant yield, and improve quality [35]. Humates can improve soil structure and water retention capacity to enhance maize quality [36, 37, 38]. Potassium humate, 75% N and 100% Reference evapotranspiration (ETO) improved soil chemical properties, leading to improved wheat yield [39]. Also, foliar applications of potassium humate can mitigate stress-related damage in soybean by promoting antioxidant growth and protecting proteins and chlorophyll [40]. The study found that adding 75% NPK, biochar, and KH to soil improved growth and water efficiency in olive trees under a half-irrigation water regime [41, 42].

The most effective treatment for crop water reduction was 75% full irrigation and 6 tons of biochar, saving 25% of the water needed [43]. Four irrigation treatments had the longest maturity period, while no irrigation treatments had the shortest [44]. The physicochemical and fertility properties of sandy soils were enhanced by optimizing water and nutrients using an 80% water regime with a 45 kg ha⁻¹ HA rate [45]. An alternate water-saving method for growing maize is 70% ETC [46]. Apply irrigation with a 4200 m² fed level and 4 tons of compost to achieve acceptable maize yield and water utilization [47]. This study aimed to investigate the impact of soil amendments (biochar, potassium humate, and compost) on soil physical properties, nutrient status, growth, and yield of wheat and maize cultivated in calcareous soil under varying irrigation regimes

2. MATERIALS AND METHODS

2.1 Field experiments location and design

The field experiment was conducted at the EL-Nubaria research farm, located in the Behaira Governorate in Egypt. The study examined the impact of soil amendments and irrigation regimes on soil physical properties, nutrient status, and productivity of maize and wheat during the 2022 summer and 2022/2023 winter seasons, under the Ministry of Agriculture and Land Reclamation's Agricultural Research Center. The geographical coordinates of the farm are 30° 90 N, 29° 96 E, with an altitude of 25 m above sea level.

The experiment was laid out in a split design with three replicates. The experimental plot area was 10.5 m² (3m x 3.5m). Irrigation regimes were placed in the main experimental plots, while organic amendments were in the subplots. The treatments were for irrigation water requirements (100, 75, and 50% of ETC), and organic amendments included five treatments: 1) biochar 9.6 ton ha⁻¹; 2) biochar 14.4 ton ha⁻¹; 3) potassium humates 1.5% (KH₁); 4) potassium humates 3% (KH₂) and 5) compost (12 ton ha⁻¹).

Some physical and chemical characteristics of the surface soil under investigation, as well as organic amendments, were analyzed before cultivation according to [48], as shown in Tables 1 and 2.

Table 1. Physical and chemical properties in soil study

Coarse sand (%)	Silt (%)	Clay (%)	Texture	SP (%)	OM (%)	CaCO ₃ (%)		
53.5	40.0	6.5	Sandy loam	48	0.75	33.65		
pH (1: 2.5)	EC (dSm ⁻¹)	Cations (meq l ⁻¹)			Anions (meq l ⁻¹)			
		Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	HCO ⁻³	Cl ⁻	SO ⁻⁴
7.96	2.4	7.5	4.50	11.45	0.25	1.5	13.5	8.97
Macronutrients (mg kg ⁻¹)				Micronutrients (mg kg ⁻¹)				
N	P	K	B	Fe	Mn	Zn		
17.8	10.12	224.38	3.14	1.89	1.37	17.8		

Table 2. Analysis of organic amendments used in the experiment.

Amendments	Bulk density (kg/m ³)	Moisture Contents (%)	pH	EC dS/m	O.C %	O.M %	C/N Ratio	Total (%)		
								N	P	K
Biochar	1060	3	7.2	2.7	0.62	1.07	1.19	0.53	0.001	0.38
Compost	580	11	6.6	2.4	26.5	45.7	33.5	0.79	0.350	0.48
k-humate	ND	ND	8.2	10	10.7	18.4	15.89	0.67	0.220	4.5

ND: not determined

2.2 Plant materials and planting

Maize hybrid triple white (Giza 310) was sown on June 1st and wheat (Misr 1) was planted on November 18th obtained from the Field Crops Institute. Nitrogen fertilizers such as ammonium sulfate (20.6%) were added to the soil. Phosphorus fertilizer as superphosphate (15.5% P₂O₅) and potassium fertilizer as potassium sulfate (48% K₂O), micronutrients, weeds, and diseases were added according to the Ministry of Agriculture and Land Reclamation of Egypt's recommendation. Potassium humate treatments were sprayed to coat the leaf surface and drenched the soil around plants at a rate of 1440 L/ha. The plants were sprayed three times at 30, 60 and 90 days after planting.

2.3 Plant-water and irrigation water measurements

The E_{Tc} value of crops in the Nubaria region was determined using reference evapotranspiration and crop factor. Table 3 shows the average values of standard evapotranspiration. The climatic data from the EL-Nubaria Agricultural Research Station, Ministry of Agriculture and Land Reclamation, Egypt, was used to calculate the E_{To} using CROPWAT 8[49]. The crop evapotranspiration was calculated using the following equation [50] E_{Tc} = E_{To} × K_c, where: E_{Tc} is crop evapotranspiration (mmd-1), E_{To} is the reference evapotranspiration (mmd-1) and K_c: crop coefficient.

Applied irrigation water was calculated according to the following equation: AIW = E_{Tc} / E_a (1 - LR) where: AIW: applied irrigation water (mmd-1), E_{Tc} evapotranspiration values (mmd-1) and E_a is the irrigation application efficiency. LR is the leaching requirement (assuming that

10% of the calculated applied irrigation water is additionally applied per irrigation during the growing season for leaching purposes)

Drought Tolerance Efficiency (DTE): was calculated using formula [51]:

$$\text{DTE (\%)} = \frac{\text{Yield under stress}}{\text{Yield under non stress}} \times 100$$

Water utilization efficiency (WUE): was estimated for each crop in according to [52] as follows:

$$\text{WUE (Kg m}^{-3}\text{)} = \frac{\text{Seed Yield (Kg)}}{\text{Water applied (m}^3\text{)}}$$

2.4 Plant and soil sampling and analysis

At harvest time, maize and wheat plants (after 120 and 150 days from planting, respectively) from each plot were cut and air-dried, and grain yield and straw yield (ton ha^{-1}) were estimated. Also, 1000 grain weights for wheat and 100 grains for maize were weighted. In addition, plant height (cm) and ear weight (gm) of maize were determined as yield components of maize. Oven-dried plant samples were analyzed for nitrogen (N), phosphorus (P), potassium (K), iron (Fe), manganese (Mn) and zinc (Zn) as described by [53].

Soil samples were analyzed for available N, P, K, Fe, Mn and Zn according to methods described by [48]. Organic carbon was determined by the modified Walkley-Black method [54]. Soil hydraulic conductivity (HC) was determined using undisturbed samples from the cores [55]. Soil moisture retention curves were determined by exposing the completely saturated samples to constant suction levels of 0.1, 0.33, 1.0 and 15 atmospheres using a pressure cooker and membrane [56]. The moisture percentages at each suction level were calculated volumetrically.

2.5 Statistical Analysis

The significant differences among means were tested using the least significant differences (L.S.D.) [57].

3. RESULTS AND DISCUSSION

3.1 Organic matter

Data in Tables 4 and 5 revealed the effect of irrigation treatments and organic amendments on the soil organic matter (O.M) % in calcareous soil under wheat and maize crops. It was observed that the maximum value of O.M% was obtained by compost (12 ton ha^{-1}), followed by B2 (14.4 ton ha^{-1}). This increment may be due to the high content of organic matter in compost as compared to biochar (Table 2). Thus, the soil O.M content resulting from applying soil amendments can be arranged in the following descending order after wheat or maize crops: compost > B₂ > KH₂.

These results are consistent with those obtained by [28] who reported that biochar and compost enhance soil organic matter. Besides, the results achieved by [58] showed that compost organic amendments markedly increased soil organic matter compared to control treatments. Additionally, the results achieved by [59], who mentioned that biochar application markedly intensified the values of soil organic matter in calcareous sandy soil.

3.2 Available NPK of calcareous soil

The results in Tables 4 and 5 indicated the effect of irrigation treatments and organic amendments on the soil availability of NPK. It was observed that applying 100% of Etc (I₁₀₀) significantly affected the soil available NP, followed by applying 75% of Etc (I₇₅). However,

the available K was not significantly affected by irrigation treatments. Applying 1100 increased soil nitrogen availability possibly due to its effective chelating properties, reducing nutrient loss [39]. The study revealed that organic amendments significantly impacted the

UNDER PEER REVIEW

Table 4. Effect of irrigation water regime and organic amendments on the organic matter, availability of NPK and their content in wheat under calcareous soil.

Irrigation water regime	Organic amendments	Soil				Straw			Grain		
		OM (%)	N (mg kg ⁻¹)	P (mg kg ⁻¹)	K (mg kg ⁻¹)	N (%)	P (%)	K (%)	N (%)	P (%)	K (%)
I ₅₀	B ₁	1.92	18.85	10.27	157.55	0.60	0.08	0.98	1.14	0.27	0.98
	B ₂	2.01	21.61	10.74	161.54	0.73	0.10	1.03	1.26	0.31	1.12
	KH ₁	1.74	23.72	11.14	169.52	0.90	0.11	1.12	1.54	0.37	1.13
	KH ₂	1.82	24.79	12.76	171.51	0.93	0.14	1.16	1.66	0.39	1.15
	Compost	2.17	26.95	13.25	187.46	0.98	0.16	1.23	1.72	0.43	1.18
	Mean	1.93	23.18	11.63	169.52	0.83	0.12	1.10	1.46	0.35	1.11
I ₇₅	B ₁	1.95	21.80	11.59	155.56	0.81	0.14	1.09	1.26	0.29	1.27
	B ₂	1.97	24.92	12.84	171.51	0.82	0.16	1.12	1.33	0.34	1.43
	KH ₁	1.84	25.93	13.77	191.50	0.88	0.19	1.15	1.47	0.42	1.45
	KH ₂	1.89	26.46	14.20	195.44	0.92	0.20	1.19	1.82	0.45	1.54
	Compost	2.09	27.53	15.05	205.41	0.95	0.20	1.26	1.84	0.47	1.66
	Mean	1.95	25.33	13.62	183.88	0.88	0.18	1.16	1.54	0.39	1.47
I ₁₀₀	B ₁	2.13	23.91	15.40	159.52	0.96	0.14	1.14	1.12	0.30	1.67
	B ₂	2.35	27.08	16.10	176.33	1.01	0.16	1.17	1.40	0.33	1.68
	KH ₁	1.96	29.11	14.20	195.44	1.02	0.18	1.20	1.55	0.44	1.71
	KH ₂	2.21	31.10	15.60	205.42	1.04	0.22	1.23	1.70	0.48	1.75
	Compost	2.67	33.81	16.88	216.72	1.18	0.23	1.28	2.18	0.51	1.77
	Mean	2.26	29.01	15.64	190.69	1.04	0.19	1.20	1.59	0.41	1.71
LSD at 0.05	Irrigation	ns	0.97	2.81	ns	0.11	ns	0.07	ns	ns	0.21
	Organic	0.20	0.52	1.13	10.7	0.11	0.04	ns	0.12	ns	0.13
	Interaction	ns	***	ns	ns	ns	ns	ns	ns	ns	ns
Mean of Organic amendments	B ₁	1.85	21.52	12.64	157.54	0.79	0.12	1.07	1.17	0.29	1.31
	B ₂	1.97	24.25	13.08	169.79	0.85	0.14	1.10	1.33	0.33	1.41
	KH ₁	2.00	20.25	13.23	185.49	0.93	0.16	1.16	1.52	0.41	1.43
	KH ₂	2.11	27.45	14.19	190.79	0.96	0.19	1.19	1.72	0.44	1.48
	Compost	2.31	29.43	15.06	203.20	1.04	0.20	1.26	1.91	0.47	1.54

Note: ** and *** refer to significance level of 5% (p≤0.05) and 1% (p≤0.01). ns:no-significant.

Table 5. Effect of irrigation water regime and organic amendments on the organic matter, availability of NPK and their content in maize under calcareous soil.

Irrigation water regime	Organic amendments	Soil				Straw			Grain		
		OM (%)	N (mg kg ⁻¹)	P (mg kg ⁻¹)	K (mg kg ⁻¹)	N (%)	P (%)	K (%)	N (%)	P (%)	K (%)
I ₅₀	B ₁	2.24	22.66	11.40	162.11	1.12	0.12	1.04	1.21	0.32	1.51
	B ₂	2.39	24.50	12.54	171.65	1.32	0.14	1.05	1.30	0.36	1.64
	KH ₁	2.08	26.80	13.49	194.69	1.49	0.17	1.24	1.40	0.43	1.71
	KH ₂	2.13	29.77	14.13	209.72	1.55	0.18	1.30	1.59	0.52	1.84
	Compost	2.69	33.70	15.19	223.93	1.74	0.21	1.43	1.68	0.58	1.94
	Mean	2.31	27.49	13.35	192.42	1.44	0.16	1.21	1.44	0.44	1.71
I ₇₅	B ₁	2.29	24.50	11.61	183.77	1.18	0.19	1.15	1.33	0.32	1.65
	B ₂	2.45	30.71	12.27	202.40	1.36	0.20	1.16	1.41	0.44	1.74
	KH ₁	2.15	31.86	13.69	214.03	1.60	0.21	1.34	1.56	0.48	1.82
	KH ₂	2.24	33.70	13.95	225.67	1.78	0.22	1.36	1.31	0.53	1.90
	Compost	2.79	35.42	16.12	232.63	2.05	0.25	1.48	1.77	0.66	2.14
	Mean	2.38	32.14	13.53	211.50	1.60	0.21	1.30	1.48	0.49	1.85
I ₁₀₀	B ₁	2.52	28.55	13.25	199.20	1.53	0.23	1.38	2.17	0.35	1.73
	B ₂	2.72	30.66	14.46	207.03	1.59	0.28	1.41	2.54	0.45	1.80
	KH ₁	2.36	34.31	14.94	226.00	1.79	0.25	1.46	2.84	0.56	1.97
	KH ₂	2.40	36.23	15.87	245.41	1.86	0.31	1.82	2.74	0.64	2.15
	Compost	3.31	39.95	16.64	260.62	2.59	0.34	1.90	2.97	0.73	2.27
	Mean	2.66	35.04	15.03	227.65	1.87	0.28	1.59	2.65	0.54	1.98
LSD at 0.05	Irrigation	ns	ns	0.85	17.46	0.15	0.03	0.18	0.11	ns	ns
	Organic	0.25	9.28	0.89	10.61	0.24	ns	0.22	0.12	0.07	0.19
	Interaction	ns	ns	ns	ns	ns	ns	ns	**	ns	ns
Mean of Organic amendments	B ₁	2.20	25.42	12.09	181.69	1.28	0.18	1.19	1.57	0.33	1.63
	B ₂	2.26	25.62	13.09	193.36	1.43	0.21	1.21	1.75	0.42	1.73
	KH ₁	2.35	33.23	14.04	211.57	1.63	0.20	1.35	1.88	0.49	1.83
	KH ₂	2.52	36.02	14.65	226.93	1.73	0.24	1.49	1.93	0.56	1.96
	Compost	2.92	37.66	15.98	239.06	2.13	0.27	1.60	2.14	0.66	2.12

Note: ** and *** refer to significance level of 5% (p≤0.05) and 1% (p≤0.01). ns:no-significant.

availability of NPK in calcareous soil, with compost being the most effective, followed by KH₂ and B₂. [34] found that compost significantly increased soil nitrogen (104.4%), followed by K-H, which increased soil available nitrogen by 81.8%. Besides, [60] found that biochar application significantly increased soil available nitrogen. In this respect, biochar and compost enhance soil phosphorus [29]. Moreover, the highest available phosphorus value was found in 100% of irrigation requirements, followed by 80% and 60% WR. [30] achieved a significant increase in soil macronutrient availability by applying high rates of compost and biochar to their soil. Concerning the effect of interaction between irrigation regimes and soil organic amendments on soil macronutrient availability, the statistical analysis showed that there's only interaction between irrigation treatments and soil organic amendments on soil available N under wheat crop. So, applying compost and I100 gave the maximum value of soil available N.

3.3 NPK contents of straw

Data in Tables 4 and 5 revealed that irrigation treatments had a significant effect on the N and K contents of wheat straw, but there was no significant effect of irrigation regimes on the P contents of wheat straw. The highest values of N and K in wheat straw were achieved by applying I₁₀₀, followed by applying I₇₅ (Table 4). While irrigation regimes significantly affected the NPK content of maize straw (Table 5), the lowest NPK contents of maize straw were achieved by applying I₅₀. In this respect, organic amendments significantly affected the NP contents of wheat straw. The maximum values of NP were obtained by compost, followed by KH₂ (Table 4). On the other hand, the NK contents of maize straw were significantly affected by organic amendments. KH₁ achieved lower values of NK compared to compost (Table 5). However, the lowest values of NK were achieved by applying 9.6 ton ha⁻¹ of biochar (B₁). These results agree with those reported by [63] who mentioned that organic materials achieved the highest nitrogen, phosphorus, and potassium contents in wheat straw compared to the control

3.4 Macro-nutrients contents of grains

Data in Tables 4 and 5 reveal that there was no significant effect of irrigation requirements on the N and P contents of grains. However, the K content of grain is significantly affected by irrigation regimes. The maximum K content of grain was achieved by I₁₀₀, followed by I₇₅. On the other hand, organic amendments significantly affect the N and K contents of grain. Although the phosphorus content of grain is not significantly affected by organic amendments, the highest values of NK contents of grain were under the wheat crop (Table 4). Additionally, the maximum values of NPK contents of maize grains were achieved by applying compost, followed by another attained by applying KH. This increase may be attributed to the high values of available NPK (Table 4). In contrast, the lowest values of NPK in wheat or maize crops were obtained by applying B₁ in calcareous soil. These results were in accordance with those obtained by [34], who mentioned that compost significantly increased the N content of plants cultivated in calcareous soil compared to KH.

3.5 Available micro-nutrients of the calcareous soil

The results obtained in Tables 6 and 7 show that the available Fe, Mn, and Zn significantly increased as irrigation requirements increased from 50% to 100% of of ET_c. Consequently, the highest values of available Fe, Mn, and Zn were obtained with I₁₀₀. This increase may be attributed to the application of soil organic amendments, which increase the organic matter content of the soil, improving water retention and plant growth [11, 45]. The study also found that compost and potassium humate significantly enhance soil micronutrient availability, with compost providing the highest values of Fe, Mn, and Zn. Potassium humate enhances soil micronutrient availability by chelating and releasing them [61]. It was observed that the availability of micronutrients affected by applying soil organic materials followed this descending order: compost > potassium humate > biochar. Applying biochar may enhance the soil's capacity to retain nutrients, potentially increasing the availability of micronutrients [62]. Moreover, biochar's porous structure and large surface area enhance soil nutrient availability and reduce leaching [13]. Regarding the interaction between irrigation regimes and soil amendments on the availability of micronutrients in the soil, highly significant values of available Fe, Mn, and Zn were obtained by applying compost and I₁₀₀ under wheat and maize crops. On the other hand, the least significant values of soil available Fe, Mn, and Zn were achieved by applying B₁ (9.6 ton ha⁻¹) and I₅₀ (50% of ET_c).

3.6 Micro-nutrients contents of straw

The results in Tables 6 and 7 reveal that the Fe, Mn and Zn contents of straw behaved similarly to those obtained in wheat and maize grains, as mentioned above. Thus, the maximum significant increment in Fe, Mn and Zn contents of wheat and maize straw was achieved by applying I₁₀₀. In this respect, applied compost or KH

Table 6. Effect of irrigation water regime and organic amendments on the availability of Fe,Zn, Mn and their content in wheat under calcareous soil.

Irrigation water regime	Organic amendments	Soil (mg kg ⁻¹)			Straw (mg kg ⁻¹)			Grain (mg kg ⁻¹)		
		Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn	Zn
I ₅₀	B ₁	3.22	2.21	1.12	10.60	11.00	10.15	32.40	15.60	15.82
	B ₂	3.54	2.30	1.15	14.60	15.16	14.42	33.60	19.17	16.00
	KH ₁	3.58	2.38	1.33	15.70	16.37	14.65	34.00	23.80	17.68
	KH ₂	3.69	2.47	1.37	16.50	16.92	15.40	35.00	36.56	19.45
	Compost	4.18	2.53	1.52	18.30	18.48	17.80	47.40	36.97	25.20
	Mean	3.64	2.38	1.30	15.14	15.59	14.48	36.48	26.39	18.83
I ₇₅	B ₁	3.26	2.57	1.39	11.20	13.24	13.31	35.60	19.42	14.92
	B ₂	3.54	2.62	1.44	12.90	15.61	15.86	39.60	22.30	16.80
	KH ₁	3.73	2.66	1.57	16.40	19.34	16.60	41.20	24.35	17.63
	KH ₂	3.88	2.77	1.95	17.40	21.40	19.26	43.40	29.14	21.48
	Compost	4.27	3.25	2.38	19.20	22.21	22.20	54.60	35.83	33.40
Mean	3.74	2.77	1.75	15.42	18.6	17.45	42.88	26.21	20.85	
I ₁₀₀	B ₁	3.36	3.13	1.48	13.90	13.52	12.20	38.00	21.60	16.32
	B ₂	3.75	3.29	1.77	15.60	14.71	12.94	40.60	28.80	18.40
	KH ₁	3.89	3.31	2.19	16.80	19.69	16.62	45.40	29.20	27.75
	KH ₂	4.03	3.47	2.53	17.80	23.36	19.80	49.40	33.60	29.84
	Compost	4.63	3.58	2.73	21.20	26.72	27.80	57.80	36.90	35.00
	Mean	3.93	2.36	2.14	17.06	19.60	17.87	46.24	30.02	25.46
LSD at 0.05	Irrigation	0.13	0.11	0.30	0.57	0.74	0.70	1.63	0.95	0.77
	Organic	0.06	0.05	0.04	0.27	0.31	0.29	0.73	0.47	0.40
	Interaction	**	***	***	***	***	***	***	***	***
Mean of Organic amendments	B ₁	3.28	2.64	1.33	11.9	12.59	11.89	35.33	18.87	15.69
	B ₂	3.61	2.74	1.45	14.37	15.16	14.41	37.93	23.42	1.6
	KH ₁	3.73	2.78	1.60 ¹	16.30	18.45	15.96	40.20	25.74	21.02
	KH ₂	3.87	2.90	1.95	17.23	20.56	18.15	42.60	33.10	23.59
	Compost	4.36	3.12	2.21	19.57	22.47	22.6	53.27	36.57	31.20

Note: ** and *** refer to significance level of 5% (p≤0.05) and 1% (p≤0.01). ns:no-significant.

Table 7. Effect of irrigation water regime and organic amendments on the availability of Fe,Zn,Mn and their content in maize under calcareous soil.

Irrigation water regime	Organic amendments	Soil (mg kg ⁻¹)			Straw (mg kg ⁻¹)			Grain (mg kg ⁻¹)		
		Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn	Zn
I ₅₀	B ₁	3.30	2.25	1.19	24.32	21.66	19.97	39.24	30.80	31.13
	B ₂	3.50	2.41	1.24	25.95	23.22	18.36	40.88	32.20	20.37
	KH ₁	3.80	2.48	1.46	27.64	25.47	20.13	42.50	34.75	24.28
	KH ₂	3.99	2.74	1.52	33.23	26.97	21.18	43.87	35.27	26.76
	Compost	4.72	3.06	1.72	34.22	29.23	29.84	47.30	36.69	39.26
	Mean	3.86	2.60	1.43	29.07	25.31	21.90	42.76	34.97	28.36
I ₇₅	B ₁	3.37	2.85	1.44	25.94	23.80	15.65	44.51	32.33	18.23
	B ₂	3.83	2.96	1.52	28.54	26.26	17.29	45.35	33.38	24.22
	KH ₁	4.00	3.03	1.68	32.38	27.52	20.74	48.23	34.40	25.36
	KH ₂	4.23	3.22	2.13	35.40	28.38	22.59	51.18	35.05	29.04
	Compost	4.78	3.80	2.69	36.90	31.31	34.67	53.33	39.30	41.75
	Mean	4.04	3.24	1.89	31.83	27.45	22.19	48.52	36.28	27.72
I ₁₀₀	B ₁	3.54	3.50	1.53	33.54	24.54	17.19	45.46	33.14	22.99
	B ₂	4.06	3.78	1.91	36.66	25.72	19.09	49.26	34.05	27.15
	KH ₁	4.24	3.84	2.35	37.12	27.58	28.50	50.90	35.14	36.19
	KH ₂	4.47	4.09	2.76	43.83	29.21	30.75	56.49	36.75	40.19
	Compost	5.31	4.37	3.06	44.79	33.05	38.98	60.23	41.34	45.18
	Mean	4.32	3.92	2.32	39.19	28.02	26.90	52.47	33.66	34.34
LSD at 0.05	Irrigation	0.15	0.14	0.33	1.90	0.98	1.65	2.28	1.20	2.96
	Organic	0.07	0.15	0.04	2.16	0.49	1.52	1.22	0.62	2.91
	Interaction	***	ns	***	ns	*	***	**	***	***
Mean of Organic amendments	B ₁	3.41	2.87	1.39	27.93	23.33	17.60	43.07	32.09	24.12
	B ₂	3.80	3.03	1.56	30.38	25.07	18.25	45.16	33.21	23.91
	KH ₁	4.01	3.23	1.83	32.38	26.86	23.12	47.21	34.76	28.61
	KH ₂	4.23	3.35	2.14	37.49	28.19	24.84	50.51	35.67	31.99
	Compost	4.94	3.74	2.49	38.64	31.20	34.50	53.62	39.11	42.07

Note: *, ** and *** refer to significance level of 10 (p≤0.1), 5% (p≤0.05) and 1% (p≤0.01).

realized a highly significant increment of Fe, Mn and Zn in wheat and maize crops (Tables 6 and 7). These results were in agreement with those obtained by [30, 63] who reported that biochar and compost significantly improved wheat plant straw micronutrient content.

3.7 Micro-nutrient contents of grains

Data in Tables 6 and 7 indicated that the Fe, Mn and Zn contents of grains followed the same trend as soil available Fe, Mn and Zn in the studied calcareous soil. The maximum increment of Fe, Mn and Zn achieved by I100. However, the lowest values of Fe, Mn and Zn were realized by applying I50. Similarly, applied compost attained the highest content of Fe, Mn and Zn in wheat and maize crops, followed by applying KH₂ (3%). This increment in micronutrient concentrations of wheat and maize grains may be due to the increase in soil availability of micronutrients in the studied calcareous soil resulting from the application of compost or potassium humate as soil conditioners. The results were in accordance with those obtained by [61, 62, and 30]. The interaction between irrigation regimes and organic amendments on the Fe, Mn and Zn contents of wheat and maize grains, as shown in Tables 6 and 7 indicated a highly significant increase in micronutrients was achieved by applying compost and I100 followed by those obtained by applying KH₂ and I100. The lowest values of micro-nutrients contents of wheat and maize grains were realized by applying B1 and I50 (Tables 6 and 7).

3.8 Physical properties of calcareous soil.

Data in Table 8 revealed the effect of irrigation regimes on the hydraulic conductivity (HC) of the studied calcareous soil. The results indicated that the HC (m/day) values gradually improved as the water requirements increased from I50 to I100. So, the maximum values of HC were obtained by applying I100, followed by applying I75. This may be attributed to the increased soil organic matter, which caused an enhancement of soil water retention by applying I75 and I100 (Tables 4 and 5). These results were in agreement with those obtained by [64]. The obtained results cleared that the effect of organic amendments on HC took the following descending order: compost > biochar > potassium humate. The beneficial effect of biochar on soil moisture retention may be due to the porosity of biochar as reported by [65]. These results were also in accordance with [25, 66].

The statistical analysis indicated that the combination of irrigation regimes and soil organic amendments influenced soil hydraulic conductivity (HC) under wheat and maize crops. The interaction between irrigation regimes and organic amendments significantly affected HC, with compost and I100 showing the highest impact, followed by B₂ (14.4 ton/ha) and I100. In contrast, applying KH₁ (1.5%) and 50% of ET_c had the least significant effect on HC. The results also demonstrated the influence of irrigation treatments and organic materials on soil physical properties such as field capacity (FC %), wilting point (WP%), and available water (AW%) under wheat and maize crops. Soil moisture content at FC% decreased with increasing water requirements, while WP% increased. Available water decreased gradually with increasing water requirements from 50% to 100% of ET_c. These findings align with previous studies. [43] found that applying biochar at a rate of 14.4 ton/ha and using 75% of full irrigation saved 25% of water requirements. [45] demonstrated that applying 45 Kg ha⁻¹ of humic acid under an 80% water regime improved soil physicochemical and nutrient status in sandy soils. Additionally, [46] suggested that utilizing 70% of crop evapotranspiration (ET_c) is an effective water-saving strategy for maize production. Regarding the influence of organic amendments on FC% and AW% of the studied calcareous soil, it can be considered in the following descending order: Biochar (B₁) > potassium humate (KH₁) > compost. The increase in soil moisture at FC and AW% resulted from applying biochar, which may be due to biochar becoming a soil conditioner due to its ability to store carbon, improve soil fertility, increase water retention, and increase crop yield [21, 22, 23]. Biochar has many advantages for agricultural soils, such as its high chemical and physical properties, nutrient retention, improved soil health, and the removal of pollutants. It has a high surface area, porosity, ion exchange, and water-holding capacity [24]. Moreover, compost and biochar can improve soil quality and productivity by enhancing nutrient absorption, cycling, and availability.

Regarding the interaction between irrigation regimes and applied organic amendments on F.C.% and AW% of the studied calcareous soil, data in Table 8 revealed that at 50% of ET_c, the effect of applied organic amendments on F.C.% and AW% took the following descending order: KH₁ > B₁ > compost. This increment resulted from the application of potassium humate, which may have enhanced the physicochemical and fertility properties of the studied calcareous soil by optimizing water and nutrients [45]. Meanwhile, at 75% of ET_c, the effect of applied organic amendments on the available water of the studied calcareous soil took the following descending order: B₂ > KH₂ > compost (Table 8). However, at 100% ET_c, the obtained results in Table 8 showed that both applied compost (12 ton/ha) and B₁ (9.6 ton/ha) maximized the values of field capacity and available water of the studied calcareous soil, followed by the first rate of potassium humate. These results were in accordance with the results obtained by [27] and [28], who reported that compost and biochar can improve soil quality and productivity by absorbing nutrients.

Table 8. Effect of irrigation water regime and organic amendments on some physical soil properties under wheat and maize crops.

Irrigation water regime	Organic amendments	Maize			Wheat	Maize
		FC %	WP %	AW %	HC (m d ⁻¹)	HC (m d ⁻¹)
I ₅₀	B ₁	32.31	6.10	26.21	1.62	1.31
	B ₂	28.69	5.60	22.79	1.80	1.43
	KH ₁	32.60	5.60	27.00	0.85	1.22
	KH ₂	27.65	5.60	22.05	0.87	1.29
	Compost	30.40	6.60	23.80	2.24	1.47
	Mean	30.33	5.9	24.37	1.46	1.34
I ₇₅	B ₁	29.82	5.60	24.22	1.96	1.49
	B ₂	30.40	5.60	24.80	2.12	1.57
	KH ₁	30.05	6.60	23.45	1.04	1.38
	KH ₂	30.15	6.30	23.85	1.12	1.45
	Compost	27.44	6.40	21.04	2.26	1.80
	Mean	29.57	6.1	23.47	1.70	1.56
I ₁₀₀	B ₁	29.13	6.31	24.23	2.08	1.84
	B ₂	28.46	9.64	18.82	2.19	2.11
	KH ₁	28.74	7.51	21.23	1.74	1.77
	KH ₂	27.80	6.73	21.07	1.44	1.78
	Compost	30.54	6.31	24.23	2.45	2.14
	Mean	28.93	7.29	21.92	1.98	1.93
LSD at 0.05	Irrigation	0.30	0.07	0.30	0.13	0.31
	Organic	0.35	0.09	0.41	0.11	0.04
	Interaction	***	***	***	***	***
Mean of Organic amendments	B ₁	30.38	6.06	24.77	1.92	1.56
	B ₂	29.18	6.95	22.14	2.04	1.70
	KH ₁	30.46	6.56	23.89	1.21	1.46
	KH ₂	28.53	6.21	22.32	1.14	1.51
	Compost	29.39	6.43	22.93	2.33	1.82

Note: *, ** and *** refer to significance level of 10 (p<0.1), 5% (p<0.05) and 1% (p<0.01).

3.9 Drought tolerates efficiency (DTE):

The results in Fig 1 illustrate that increasing applied irrigation water had a positive effect on the DTE values, which increased from 75.90% at I₅₀ to 85.0% at I₇₅. In addition, increasing the application rate of organic amendments had a positive effect on water stress in the I₇₅ treatments, where DTE were 88.3, 87.4, 82.3, 85.9, and 82.0% for I₇₅B₁, I₇₅B₂, I₇₅KH₁, I₇₅KH₂, and I₇₅compost, respectively. Biochar recorded the highest values, at 88.3% at I₇₅B₁ and 79.0% at I₅₀B₁. While, the irrigation water regime noted that I₇₅ had high-value drought tolerance efficiency for wheat in Fig 1. In addition, results in Figure 1 show that with maize, the highest value of drought tolerance efficiency was 97.3% at I₇₅, followed by 92.2% at I₅₀ for the irrigation water regime. Soil amendments increased by 98.4, 95.3, 96.4, 97.8, and 97.1% for I₇₅B₁, I₇₅B₂, I₇₅KH₁, I₇₅KH₂, and I₇₅compost, respectively. Biochar exhibited the highest drought tolerance efficiency at 98.4% due to its ability to retain water and reduce evapotranspiration. These findings align with previous reports [67] that drought tolerance efficiency is influenced by irrigation treatment and calcium-humate. The treatment of I₇₅ Ca-H₂O had the highest drought tolerance efficiency at 96.72%, while the irrigation water regime had 93.07%. As for the highest drought tolerance efficiency; it was higher in corn than in wheat under the influence of all treatments (Fig. 1)

3.10 Water use efficiency (WUE)

Results in Fig.2 for wheat illustrate that the highest water use efficiency value (6.1 kg/m³) was recorded for I₅₀ with compost. Decreasing the amount of applied irrigation water and increasing soil amendment applications have a highly positive effect on water use efficiency. Also, increasing the amount of irrigation water used negatively affects the value of the efficiency of water use, where WUE was better for I₅₀ compared to I₇₅ and I₁₀₀, respectively. The best additions for soil amendments were compost, followed by a high rate of potassium humate under an irrigation water regime. Also, the data in Fig. 2 for maize illustrates that the highest value of water use efficiency was I₅₀, followed by I₇₅, then I₁₀₀ of ETC. As for soil amendments, the best additives were compost, followed by potassium humate, and then

biochar under an irrigation water regime. Soil amendments are materials added to soil to improve its physical, chemical, or biological properties. They can have various effects on water retention, depending on their composition. For example, organic matter such as compost can improve water retention by increasing soil porosity and enhancing its ability to hold onto moisture. Irrigation water level at 150 with compost have significantly impacted water use efficiency, with a maximum value of 6.1 kg m⁻³. Water use efficiency was higher with maize than with wheat in all treatments under 50% irrigation water regimes (Fig. 2.). Using compost and increased potassium rates enhances water storage in the root zone, while potassium humate fertilizer mitigates water stress. Stomata rich in K keep stomata closed, reducing transpiration rate but allowing plant roots to absorb more water without additional water.




Fig.1: Drought tolerates efficiency as affected by irrigation water regime and soil amendment of wheat and maize




Fig.2: Water use efficiency as affected by irrigation water regime and soil amendments of wheat and maize

3.11 Wheat and maize yields

Results in Table 9 showed that applying I₁₀₀ significantly increased wheat and maize yields, followed by I₇₅. This increase may be attributed to improved soil properties such as organic matter and macro- and micronutrient availability, as well as increased soil hydraulic conductivity. These results were obtained by [46], who reported that 70% ET_c is an alternative water-saving strategy for maize production. Concerning the effect of organic amendments on wheat yield, the data in Table 9 indicates that this effect followed a descending order: applied compost > potassium humate > biochar. However, the effect of organic amendments on maize yield as affected by organic amendments followed a descending order: potassium humate > compost > biochar. The lowest values of wheat and maize grain yields were achieved by applying the first rate of biochar (Table 9). These findings align with previous studies by [14] and [15], who suggest that compost as a humic-rich amendment can enhance soil's chemical, physical, and biological properties. Humic substances, comprising 60% of soil organic compounds, can be utilized as soil conditioners to enhance soil structure and water retention capacity, thereby reducing fertilizer usage and enhancing plant growth. Moreover, potassium humate can also improve maize quality [36-38]. The study revealed that there was no significant difference in wheat grain yield as a result of the interaction between irrigation regimes and organic materials. However, the highest maize grain yields were observed when I₁₀₀ and compost were applied, followed by compost and I₇₅. The lowest yield was achieved with the first rate of biochar (B₁) and I₅₀ (Table 9).

3.12 Wheat and maize straw yields

Results in Table 9 show that the maximum wheat straw yield was achieved with I₇₅ irrigation, followed by I₁₀₀, while maize straw yield was affected in descending order, with 100% irrigation, 50% irrigation, and 75% irrigation resulting in the highest yield. The data in Table 9 reveals that organic amendments impact wheat straw yield in descending order, compost and KH1 above B₂, and maize straw yield in descending order: compost > potassium humate > biochar. These results were in accordance with those obtained by [14, 15].

Regarding the results in Table 9, the first rate of potassium humate and 75% of ET_c yielded significantly more wheat straw than the second rate of biochar and 75% of ET_c. The study found that the highest wheat straw yield was achieved with the second rate of potassium humate and 50% of ET_c, but the maximum significant maize straw yield was achieved with compost and I₁₀₀, followed by potassium humate and I₁₀₀. While the minimum value of maize straw yield is achieved by applying the first rate of biochar (B₁) and 50% of ET_c.

3.12.1 Weigh 1000 grains of wheat and 100 grains of maize

The obtained results in Table 9 revealed that the effect of irrigation regimes on the 1000 grain weight of wheat and the weight of 100 maize grains caused a highly significant increment in both the 1000 grain weight of wheat and the weight of 100 maize grains by increasing irrigation water requirements. The maximum significant increment of both the grain weight of wheat and the weight of 100 maize grains was achieved by applying I₁₀₀, followed by applying 75% of ET_c. The increase in the grain weight of wheat may be attributed to increased K, Fe, Mn, and Zn contents in wheat grains (Tables 4 and 6). While the increase in weight of 100 maize grains may be due to increased N, Fe, Mn, and Zn contents in maize grains, which may reflect on improving plant growth (Tables 5 and 7). The study found that increasing irrigation water requirements led to a significant increase in the weight of wheat and maize grains. Therefore, the application of I₁₀₀ and 75% of ET_c resulted in the maximum significant increase in the 1000 grain weight of wheat and 100 maize grains. The increase in grain weight of wheat may be attributed to increased K, Fe, Mn, and Zn contents in wheat grains (Tables 4 and 6). While the increase in the weight of 100 maize grains may be due to increased N, Fe, Mn, and Zn contents in maize grains, which may reflect on improving plant growth (Tables 5 and 7). Data in Table 9 clarified that the effect of applying organic amendments on 1000 grain weights of wheat and 100 grain weights of maize grains took the following descending order: compost > potassium humate > biochar. This increment in applying compost and potassium humate may be due to an increase in macro- and micronutrient contents in wheat and maize grains, which resulted in improved plant growth (Tables 4, 5, 6 and 7). Also, soil amendments can improve soil properties, maintain fertility, increase productivity, and preserve the environment [8].

Table 9: Effect of irrigation water regime and organic amendments on yield and its components of wheat and maize under calcareous soil.

Irrigation water regime	Organic amendments	Wheat			Maize				
		1000-grain weight (g)	Grain yield (ton ha ⁻¹)	Straw yield (ton ha ⁻¹)	Plant height (cm)	Ear weight (g)	Weight of 100 grains (g)	Grain yield (ton ha ⁻¹)	Straw yield (ton ha ⁻¹)
I ₅₀	B ₁	34.07	5.65	10.85	230.90	334.78	38.33	9.63	14.10
	B ₂	38.30	5.67	10.63	232.90	373.09	40.00	10.05	14.67
	KH ₁	39.80	6.00	9.63	240.97	374.78	42.00	10.59	15.93
	KH ₂	39.93	6.28	8.27	245.47	381.55	42.33	10.86	16.42
	Compost	40.20	6.71	10.51	237.00	416.66	43.00	11.19	18.25
	Mean	38.46	6.06	9.98	237.45	376.16	41.13	10.46	15.87
I ₇₅	B ₁	39.73	6.31	11.78	239.87	349.22	41.00	10.12	14.57
	B ₂	40.00	6.63	13.56	240.57	374.78	42.33	10.71	14.69
	KH ₁	42.51	6.74	13.70	249.23	410.56	42.67	11.36	15.29
	KH ₂	44.55	7.08	12.77	257.77	420.33	44.33	12.06	15.98
	Compost	53.26	7.15	12.20	245.20	436.01	45.00	12.47	16.45
	Mean	44.01	6.78	12.80	246.53	392.01	43.07	11.03	15.39
I ₁₀₀	B ₁	42.53	7.15	10.62	228.37	352.56	41.33	10.28	13.71
	B ₂	42.67	7.60	10.84	246.90	355.22	42.33	11.24	17.79
	KH ₁	55.64	8.19	10.35	249.10	364.00	44.33	11.79	18.01
	KH ₂	58.37	8.24	9.67	253.43	423.89	46.33	12.33	18.12
	Compost	57.55	8.72	10.98	248.33	464.33	52.33	12.84	20.01
	Mean	51.35	7.98	10.49	245.23	398.19	45.33	11.34	17.53
LSD at 0.05	Irrigation	3.25	0.13	0.30	3.72	10.07	1.55	0.329	0.389
	Organic	3.01	0.22	0.50	4.93	14.20	0.730	0.263	0.218
	Interaction	**	ns	***	ns	**	***	***	*
Mean of Organic amendments	B ₁	38.78	6.37	11.09	233.04	345.52	40.22	10.01	14.13
	B ₂	40.32	6.63	11.68	240.12	367.7	41.56	10.67	15.71
	KH ₁	45.98	6.98	11.23	246.43	383.11	43.00	11.24	16.41
	KH ₂	47.62	7.20	10.24	252.22	408.59	44.33	11.75	16.84
	Compost	50.33	7.53	11.23	243.51	439.00	46.78	11.05	18.24

Note: ** and *** refer to significance level of 5% ($p \leq 0.05$) and 1% ($p \leq 0.01$). *ns:no-significant.

Regarding the effect of the interaction between irrigation regimes and organic amendments on the grain weight of wheat, the results obtained in Table 10 revealed that the highly significant weight of 1000 wheat grains was obtained by applying the second rate of potassium humate (KH₂) and I₁₀₀. It was followed by applying compost and I₁₀₀. However, the lowest weight was achieved by applying the first rate of biochar (B₁) and 50% of ETc. The effect of the interaction between irrigation regimes and organic amendments on the weight of 100 maize grains is shown in Table 10. The highly significant value of 100 maize grain weight obtained by compost and I₁₀₀ is followed by applying the second rate of potassium humate and I₁₀₀. However, the least weight of 100 maize grains was achieved by applying B₂ and 50% of ETc.

3.12.2 Plant height of maize

The obtained results in Table 9 revealed that the effect of the irrigation regime on maize plant height was significantly increased by applying I₇₅, followed by applying I₁₀₀. Concerning the effect of organic amendments on maize plant height, the data in Table 10 showed that a significant increment in plant height was achieved by applying KH₂, followed by KH₁. However, the lowest height was obtained by applying B₁. These findings agree with those reported by [36, 29] who, found that combined and single-application biochar or compost additions

increased the plant height of two maize varieties. Higher plant heights (206.8 cm) were found in plots irrigated five times compared to lower irrigation regimes.

3.12.3 Ear weight of maize

Data in Table 9 indicated that irrigation regimes affected the ear weight of maize. It was observed that an increase in irrigation water requirements from I50 to I100 caused a highly significant increment in the ear weight of maize. Thus, the maximum value of the ear weight of maize was achieved by applying I100, followed by applying I75. This high value of ear weight in maize may be attributed to improved plant growth under applying I100, which resulted in increasing N, Fe, Mn, and Zn contents in maize grains, as mentioned above (Tables 5, 7). Concerning the effect of organic application on amendments application on ear weight of maize, the obtained results in Table 9 revealed that applying compost resulted in the highest significant increment of ear weight of maize, followed by applying KH₂. Whereas, the lowest one achieved B₁. This increment in ear weight of maize may be due to improving some soil characteristics and maintaining fertility, resulting in the application of soil amendments that increase the macro- and micronutrient contents of the maize grains (Tables 5 and 7). These findings agree with those reported by [68] found that reducing water stress and ensuring moisture during the maize growing season increased ear weight, possibly due to improved photosynthesis and growth. The study found that the maximum significant maize ear weight was achieved with compost with I100, followed by KH₂ with I100, and the least significant value was achieved with B₁ and 50% Etc (Table 9).

4. CONCLUSION

In calcareous soil at El-Nubaria Agricultural Research Station, the applied irrigating water (AIW) of maize was 3677.1 m³ha⁻¹ with 100% ETC for the application of compost, giving the highest yield of 12.84 ton ha⁻¹ and the highest water use efficiency of 6.1 kg m⁻³ of applied water at 50% ETC and compost. Drought tolerance efficiency is affected by irrigation water regimes and soil amendments. Therefore, I75B₁ recorded the highest values, at 98.4%. Also, the irrigation water regime showed that I75 had high-value drought tolerance efficiency, which was 97.3%. While irrigating wheat at 2218 m³ ha⁻¹ with 100% ETC at the application of compost gives the highest yield of 8.72 ton ha⁻¹ and the highest water use efficiency of 6.1 kg m⁻³ of applied water were achieved at 50% ETC and compost. Drought tolerance efficiency is affected by irrigation water regimes and irrigation water regime noted that I75 had high value drought tolerance efficiency, which was 85.0%. At the end of this study, the recommendation was to apply irrigation water at 75% of the ETC is saving 25% of the water required for crops in Egypt. Further research is needed to evaluate the impact of biochar, potassium humate and compost amendments on enhancing salt and drought resilience in soil and plants. This will aid in recommending suitable application methods considering soil management and cropping systems.

REFERENCE

1. Salem, E.M.M., Kenaway, M.K.M., Saady, H.S. & Mubarak, M. (2021). Soil Mulching and Deficit Irrigation Effect on Sustainability of Nutrients Availability and Uptake, and Productivity of Maize Grown in Calcareous Soils. *Comm. Soil Sci. Plant Anal.*, 52: 1745-1761.
2. Ali, M.A.A., Nasser, M.A., Abdelhamid, A.N., Ali, I.A.A., Saady, H.S. & Hassan, K.M. (2024). Melatonin as A Key Factor for Regulating and Relieving Abiotic Stresses in Harmony With Phytohormones in Horticultural Plants — A Review. *J Soil Sci Plant Nutr.*, 2024; 24:54-73.
3. Doklega, S.M.A., Saady, H.S., El-Sherpiny, M.A. Abou El-Yazied, A., Abd El-Gawad, H.G. Ibrahim, M.F.M. et al. (2024) Rhizospheric Addition of Hydrogel Polymer and Zeolite Plus Glutathione Mitigate The Hazard Effects Of Water Deficiency On Common Bean Plants Through Enhancing The Defensive Antioxidants. *Gesunde Pflanzen.*, 76(1): 235-249.
4. Madhavi, P., Sailaja, V., Prakash, T. R., & Hussain, S. A. (2017). Characterization Of Biochar And Humic Acid And Their Effect On Soil Properties In Maize. *Int. J. Curr. Microbiol. App. Sci.*; 6: 449-457.
5. Faostat, F. (2016). Statistics division. Food and Agriculture Organization of the United Nations..

6. Malav, A. K., Monpara, B. A., Gaur, A., & Bhati, S.S.(2017). Character Association Analysis in Yield and Yield Components in Bread Wheat (*Triticum aestivum* L.) Genotypes”, J. Plant Develop. Sci., 9: 77-83.
7. Asseng, S., Kheir, A. M., Kassie, B. T., Hoogenboom, G., Abdelaal, A. I., Haman, D. Z., & Ruane, A. C. (2018). Can Egypt Become Self-Sufficient In Wheat?, Environmental Research Letters.; 13: 094012.
8. Singh, K., Vasava, H.B., Snoeck, D., Das, B.S., Yinil, D., Field, D., & Panigrahi, N. (2019) Assessment of cocoa input needs using soil types and soil spectral analysis”, Soil Use and Management.; 35: 492-502.
9. Elshony, M., Farid, I. M., Alkamar, F., Abbas, M. H., & Abbas, H. (2019). Ameliorating A Sandy Soil Using Biochar and Compost Amendments and Their Implications as Slow-Release Fertilizers on Plant Growth”, Egyptian Journal of Soil Science.; 59:305-322.
10. Sisouvanh, P., Trelo-Ges, V., Isarangkool Na Ayutthaya, S., Pierret, A., Nunan, N. et al.(2021) Can Organic Amendments Improve Soil Physical Characteristics and Increase Maize Performances In Contrasting Soil Water Regimes?, Agriculture; 11(2): 132.
11. Kang, M.W., Yibeltal, M., Kim, Y.H., Oh, S. J., Lee, J.C., Kwon, E.E., & Lee, S.S.(2022) Enhancement of Soil Physical Properties and Soil Water Retention With Biochar-Based Soil Amendments, Science of The Total Environment; 836:155746.
12. Hale, L., Curtis, D., Azeem, M., Montgomery, J., Crowley, D. E., & McGiffen, M. E. Jr.(2021) Influence of compost and biochar on soil biological properties under turfgrass supplied deficit irrigation”, Applied Soil Ecology; 168:104134.
13. Konz, J., Cohen, B., & Van der Merwe, A. (2015). Assessment of The Potential to Produce Biochar And Its Application to South African Soils As a Mitigation Measure”, Environmental Affairs Department: Republic of South Africa: Pretoria, South Africa.
14. Abouhussien, E., Elbaalawy, A. M., & Hamad, M. (2019). Chemical Properties of Compost in Relation to Calcareous Soil Properties and Its Productivity of Wheat”, Egyptian Journal of Soil Science; 59: 85- 97.
15. Mohamed, M.S. & Rashad, R.T. (2020). Studying Some Characteristics of Sandy Soil Amended By Water Hyacinth, Bean Straw, And Compost”, Egyptian Journal of Soil Science; 60(1): 53-65.
16. Badar, R. Batool, B. Ansari, A. Mustafa, S. Ajmal, A. & Perveen, S.(2015). Amelioration of Salt Affected Soils for Cowpea Growth by Application of Organic Amendments, Journal of pharmacognosy and phytochemistry; 3(6):87-90.
17. Barthod, J., Rumpel, C., & Dignac, M. F.(2018). Composting With Additives to Improve Organic Amendments” A review. Agronomy for Sustainable Development; 38(2): 17.
18. Mansoor, S. Kour, N. Manhas, S. Zahid, S. Wani, O.A. Sharma, V. et al.(2021) Biochar as a tool for effective management of drought and heavy metal toxicity”, Chemosphere; 271: 129458.
19. Bolan, N. Hoang, S.A. Beiyuan, J. Gupta, S. Hou, D. Karakoti, A. et al.(2022) Multifunctional applications of biochar beyond carbon storage”, International Materials Reviews; 67: 150-200.
20. Jing, F., Sun, Y., Liu, Y., Wan, Z., Chen, J., & Tsang, D.C. (2022) Interactions Between Biochar And Clay Minerals In Changing Biochar Carbon Stability”, Science Of The Total Environment; 809:151124.
21. Rodrigues M.A., Garmus T., Arrobas M., Gonçalves A., Silva, E., Rocha, L. et al. (2019). Combined biochar and organic waste have little effect on chemical soil properties and plant growth”, Spanish Journal of Soil Science;. 9:199-211.
22. Xing Y., Wang J., Shaheen S.M., Feng X., Chen Z., Zhang H., Rinklebe J.(2020). Mitigation of mercury accumulation in rice using rice hull-derived biochar as soil amendment: A field investigation. Journal of Hazardous Materials; 388:121747.
23. Huang, H., Reddy, N. G., Huang, X., Chen, P., Wang, P., Zhang Y. et al. (2021). Effects of pyrolysis temperature, feedstock type and compaction on water retention of biochar amended soil”, Scientific Reports; 11: 7419.
24. Haider, F. U., Coulter, J. A., Liqun, C. A. I., Hussain, S., Cheema, S. A., Jun, W. U., & Zhang, R.(2022). An overview on biochar production, its implications, and mechanisms of biochar-induced amelioration of soil and plant characteristics. Pedosphere; 32: 107-130.

25. Yadav, N. K., Kumar, V., Sharma, K. R., Choudhary, R. S., Butter, T. S., Singh, et al.(2018). Biochar and their impacts on soil properties and crop productivity: a review”, *Journal of Pharmacognosy and Phytochemistry*; 7: 49-54.
26. T.M. Salem, K.M. Refaie, A.E.-H.E.-G.Abd, E-L. Sherif, M.A.M. Eid, “Biochar application in alkaline soil and its effect on soil and plant”,*Acta Agriculture Slovenica*,vol. 114, pp. 85-96, 2019.
27. Sánchez-Monedero, M.A., Cayuela, M.L., Sanchez-Garcia, M., Vandecasteele, B., D'Hose, T., Lopez, G., Martinez-Gaitan, C., Kuikman, P.J., Sinicco, T., Mondini, C. (2019). Agronomic evaluation of biochar, Compost and biochar-blended Compost across different cropping systems: Perspective from the European project FERTIPLUS. *Agronomy*; 9: 225.
28. Mahmoud, E., Ibrahim, M., AN, N., AN, H.(2020). Effect of biochar and Compost amendments on soil biochemical properties and dry weight of canola plant grown in soil contaminated with heavy metals. *Communications in Soil Science and Plant Analysis*; 51:1561-1571.
29. Mensah, A.K., Frimpong, K.A.(2018). Biochar and/or Compost applications improve soil properties, growth, and yield of maize grown in acidic rainforest and coastal savannah soils in Ghana. *International Journal of Agronomy*; 7: 1-8.
30. Mancy, A., Sheta, M.(2021). Evaluation of biochar and Compost ability to improve soil moisture content and nutrients retention. *Al-Azhar Journal of Agricultural Research*; 46:153-165.
31. Jindo, K., Olivares, F.L., Malcher, D.J.d.P., Sánchez-Monedero, M.A., Kempenaar, C., Canellas, L.P.(2020). From lab to field: Role of humic substances under open-field and greenhouse conditions as biostimulant and biocontrol agent. *Frontiers in plant science*; 11: 426.
32. Justi, M., Morais, E.G., Silva, C.A., 2019. Fulvic acid in foliar spray is more effective than humic acid via soil in improving coffee seedlings growth. *Archives of Agronomy and Soil Science*, 65:1969–1983.
33. Gerke, J.(2018). Concepts and misconceptions of humic substances as the stable part of soil organic matter: A review. *Agronomy*, 8:76.
34. Sary, D.H., Rashad, R.T. (2021). Nutrient use Efficiency in Calcareous Soil Amended by the Silicate, Humate and Compost. *Asian Journal of Soil Science and Plant Nutrition*, 7:1-12.
35. Ouni, Y., Ghnaya, T., Montemurro, F., Abdelly, C., Lakhdar, A., 2014. The role of humic substances in mitigating the harmful effects of soil salinity and improve plant productivity. *International Journal of Plant Production*; 8:353-374.
36. Khan, M.I., Jan, A.(2018). impact of Soil Conditioning and Irrigation Regimes on the Performance of MaizeCrop. *Sarhad Journal of Agriculture*; 34:173-187.
37. Karčauskienė, D., Repšienė, R., Ambrazaitienė, D., Mockevičienė, I., Šiaudinis, G., Skuodienė, R. Melatonin as A Key Factor for Regulating and Relieving Abiotic Stresses in Harmony With (2019). A complex assessment of mineral fertilizers with humic substances in an agroecosystem of acid soil. *Zemdirbyste-Agriculture*; 106(4): 307-314.
38. Rashad, R.T. 2020. Effect of Soaking Grains in Some Growth Regulators on Wheat Grown in Sandy Soil. *Egyptian Journal of Soil Science*; 90:99-108.
39. Hassan, W., El-Farghal, W., Khalil, F., EL-Etr, W.(2017). Effect of some soil amendments and irrigation treatments on wheat crop productivity in middle Egypt. *Journal of Soil Sciences and Agricultural Engineering*; 8:553-563.
40. El-Nwehy, S.S., Sary, D.H., Afify, R.R.M.(2020). Effect of potassium humate foliar application on yield and quality of soybean (*Glycine max L.*) grown on calcareous soil under irrigation water regime. *Plant Arch*; 20:1495-1502.
41. Sary, D.& Elsokkary, I.(2019). Effect of irrigation water regime in presence of organic or biological fertilizer on olive trees. *Egyptian journal of soil science*; 59: 67-48.
42. Attia, R.H., Sary, D.H.(2021). Effect of some soil conditioners on snap bean growth yield and sandy soil properties. *Journal of Soil Sciences and Agricultural Engineering*; 12:123-129.
43. Abdelraouf, R., Essay, E., Saleh, M., 2017. Sustainable management of deficit irrigation in sandy soils by producing biochar and adding it as a soil amendment. *Middle East Journal of Agriculture Research*; 6:1359-1375.
44. Islam, S., Haque, M., Hasan, M., Khan, A., Shanta, U.(2018). Effect of different irrigation levels on the performance of wheat. *Progressive Agriculture*; 29: 99-106.

45. Al-Shareef, A.R., Ismail, S.M., El-Nakhlawy, F.S.(2018). Effect of humic acid application rates on physicochemical and fertility properties of sandy loam soil grown with mung bean under different irrigation water regimes. *Current Science*; 115:1374-1379.
46. Singh, M., Singh, S., Deb, S., Ritchie, G.(2023). Root distribution, soil water depletion, and water productivity of sweet maize under deficit irrigation and biochar application. *Agricultural Water Management*, 279: 108192.
47. Khatab, A.K., Abd El-Latif, K.M., Osman, E., Abdou, S.(2015). Maize productivity and crop-water relations as affected by irrigation levels and Compost rates. *Journal of Soil Sciences and Agricultural Engineering*; 6:1545-1562.
48. Page, A., Miller, R., Keeney, D., Baker, D., Roseoc Ellis, J., Rhodes, J.(1982). *Methods of soil analysis Part 2:chemical and Microbiological Properties*, Agronomy Monograph No. 9. American society of Agronomy and Soil Science Society America Madison, Wisconsin, USA.
49. Doorenbos, J., Kassam, A.(1986). *Yield Response to Water*; FAO Irrigation and Drainage, Paper No. 33; Food and Agriculture Organization of the United Nations. Water Resources and Development Service: Rome, Italy.
50. Allen, R.G., Pereira, L.S., Raes, D. and Smith, M. (1998) *Crop evapotranspiration – Guidelines for computing crop water requirements*. FAO Irrigation and Drainage Paper, No. 56, FAO, Rome.
51. Fischer, K. S. & Wood, G. 1981 Breeding and selection for drought tolerance in tropical maize. In: *Proceedings of Symposium on principles and methods in crop improvement for drought resistance with emphasis on rice*, IRRI, Philippines.
52. Abdullaev, I., & Molden, D. (2004). "Spatial and temporal variability of water productivity in the Syr Darya Basin, central Asia", *Water Resources Research*; 40:6.
53. Estefan, G., Sommer, R., Ryan, J., 2013. Estefan, G., Sommer, R. & Ryan, J. *Methods of soil, plant and water analysis: A manual for the West Asia and North. Int. Cent. Agric. Res. Dry Areas 30*
54. Nelson, D.W., Sommers, L.E.(1996). Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 3 Chemical methods 5*, 961-1010.
55. Klute, A., Dirksen, C.(1986). Hydraulic conductivity and diffusivity: Laboratory methods. *Methods of soil analysis: Part 1 physical and mineralogical methods 5*, 687-734.
56. Stakman, W. P. (1966). *Determination of soil water retention Curves. Sand box method 11. Pressure membrane methods" ICW. Wageningen, the Netherlands.*
57. Snedecor, G.W. and W.G. Cochran (1990). *Statistical Methods*. 8th Ed. Iowa State Univ., Press, Ames, Iowa, USA.
58. El-Ngar, D.A., Zein El-Abdeen, H.A., Sary, D.H.(2022). Effect of Compost and Nano-Hydroxyapatite on Phosphorus Nutrition and some Properties of Calcareous Soil. *Journal of Soil Sciences and Agricultural Engineering*; 13:317-324.
59. Amin, A.E.E.A.Z., Eissa, M.A.(2017). Biochar effects on nitrogen and phosphorus use efficiencies of zucchini plants grown in a calcareous sandy soil. *Journal of soil science and plant nutrition*; 17:912-921.
60. Yang, W., Feng, G., Jia, Y., Yang, Y., Gao, X., Gao, L., Qu, Z.(2022). Impact of single biochar application on maize growth and water-fertilizer productivity under different irrigation regimes. *Frontiers in Plant Science* 13.
61. Awwad,M., El-Hedek, K., Bayoumi, M., Eid, T.(2015). Effect of potassium humate application and irrigation water levels on maize yield, crop water productivity and some soil properties. *Journal of Soil Sciences and Agricultural Engineering*; 6: 461-482.
62. Silva, I.C.B.d., Basilio, J.J.N., Fernandes, L.A., Colen, F., Sampaio, R.A., Frazao, L.A.(2017). Biochar from different residues on soil properties and common bean production. *Scientia Agricola*; 74:378-382.
63. Sary, D.H., Hamed, E.N.(2021). Effect of Salicylic, Humic and Fulvic Acids Application on The Growth,Productivity and Elements Contents of Two Wheat Varieties Grown Under Salt Stress. *Journal of Soil Sciences and Agricultural Engineering* ; 12: 657 -671.
64. Sharma, P., Abrol, V., Sharma, V., Chaddha, S., Rao, C.S., Ganie, A., Hefft, D.I., El-Sheikh, M.A., Mansoor, S.(2021). Effectiveness of biochar and Compost on improving soil hydro-

- physical properties, crop yield and monetary returns in inceptisol subtropics. *Saudi Journal of Biological Sciences*; 28:7539-7549.
65. De Jesus Duarte, S., Glaser, B., Pellegrino Cerri, C.E.(2019). Effect of biochar particle size on physical, hydrological and chemical properties of loamy and sandy tropical soils. *Agronomy*; 9(4): 165.
 66. Ibrahim, M., Mahmoud, E., Gad, L., & Khader, A. (2019). Effects of biochar and phosphorus fertilizer rates on soil physical properties and wheat yield on clay textured soil in middle Nile Delta of Egypt", *Communications in Soil Science and Plant Analysis*, 50: 2756-2766.
 67. Sary, D. H. Niel, E.M. & Abd El-All, A.E.A. (2021). Effect of Drip Irrigation Water Regime and Calcium Humate on the Growth Performance, Yield of Faba Bean (*Vicia Faba L.*) Grown in Sandy Soil", *American-Eurasian Journal of Scientific Research*, 16: 53-66, 2021.
 68. Al-Dulaimi, O.I. Al-Ani, M.H. Al-Rawi, A.R. & Seadh, S.E. (2020). Effect of water stress and organic fertilization sources on maize growth and yield", *Int. J. Agricult. Stat. Sci*, 2020.