

“Modeling and forecasting of potato production in Rajasthan and their yield sustainability”

Abstract

This study analyzed trends, sustainability, and forecasted potato area, production, and productivity in India and Rajasthan from 1970 to 2030. Annual potato production data was modeled with an autoregressive integrated moving average (ARIMA). The models were trained on 1970–2020 data and evaluated on a 2021–2023 validation set. On the training set, a box showed the best performance. The ARIMA models produced the lowest forecast errors. The top-performing models forecast potato production up to 2030. The production of potato in Rajasthan and India in 2023 was 260.50 and 61250.50 thousand tonnes. Rajasthan and India are predicted in 2030 to produce 301.31 and 63318.26 thousand tonnes thousand tons, The forecasts can assist food security planning and agricultural policymaking in the region. Rajasthan and India are highly sustainable in the yield of potato in period-2 (1998–2023).

Keywords: Trend, Sustainability, ARIMA, Time Series, Policy.

Introduction: - **Potato** (*Solanum tuberosum*) is an annual herbaceous tuber crop belonging to the family Solanaceae, one of the most horticulture crops that have the highest nutrition and dry matter production per unit area and time of any major food crop. It is the world’s most significant crop for food, with a 78% water content and a 22% dry matter content. 20% fat, 2.1% protein, 1.1% dietary fiber, 0.9% ash, and 0.3% carbs. Potato is another important source of raw materials for the extraction of starch, with starch extraction typically ranging from 9% to 25%. According to the total amount of mineral compounds accumulated in potatoes, they exceed many types of fruits and vegetables, 18 amino acids are found in potato proteins. Potatoes are characterized by high nutritional value and are a source of vitamins C, A, (carotene), group B, and K. By eating an average of 300 g (about 10.58 oz) of boiled potatoes every day, we provide ourselves with the necessary carbohydrates, phosphorus, vitamin 8 %, 10-12%, amino acids 30-40%, vitamin C 40-50%, vitamin B- 20-25%, potassium 55-60%, calcium and iron-25-30% and carotene 1-2%. Sustainable agriculture" is defined as "agriculture that, over the long term, improves the environment quality and the resource base on which agriculture depends; provides for the basic needs of human food and fiber; is economically viable; and enhances the quality of life for farmers and society as a whole. Farming has always been characterized by large and irregular

changes in location, demand, and yield; this makes forecasting beneficial to farmers, governments, and the agribusiness sector. Due to the critical role that food production plays in a nation's stability, governments are becoming major providers and users of agricultural prediction (Mishra *et al.*, 2015). The area of potato production in India during 2022-23 is 2332 Thousand ha, with production of 61250.50 MT (Metric ton) and a yield of 257 q/ha. The total area, production, and productivity of potato in Rajasthan in 2021 is about 13.59 Thousand ha, 256.00 thousand tonnes, and 188.33 q/ha (Ministry of Agriculture and Farmer Welfare, Govt. of India, 2023).

Mishra *et al.*, (2024) utilized ARIMA (Auto Regressive Integrated Moving Average) and ETS (Error-Trend-Seasonality) forecast models to predict potato production. Kumari *et al.*, (2024) This study focuses on the crucial requirement for precise prediction of potato price, which is essential for optimizing production, marketing tactics, and inventory management. Yadav *et al.*, (2024) utilized autoregressive integrated moving average (ARIMA) models, state space models, and gradient-boosting machine learning approaches to analyze the annual potato production data from 1967 to 2020. Sahu *et al.*, (2024) utilized time series data on area, production, productivity, market prices, export quantity, and export value in the primary cultivating states, markets, and importing nations, respectively. Devi *et al.*, (2021) study looked at the sustainability and instability of wheat production in Haryana.

MATERIAL AND METHODS

Source of data

In the present investigation, secondary data has been gathered. A glance at Agricultural Statistics (2022) provided the area, production, and yield data on potato for the years 1970–2023 for Rajasthan and Whole India.

Descriptive Statistics

Descriptive statistics are frequently used to make numerical data easily understandable. We have the option of using one or more criteria to assess a large sample of people. We can make sense of very large datasets by using descriptive statistics. The three categories of descriptive statistics are central tendency (CT), dispersion, and association measures.

Trend models

A (Mishra *et al.* 2015) model is a way to illustrate a system or process. Statistical models often follow the course of the process together with its statistical characteristics and implications.

- i. **Linear Model:** $Y_t = b_0 + b_1t$

- ii. **Quadratic Model:** $Y_t = b_0 + b_1t + b_2t^2$
- iii. **Compound Model:** $\ln(Y_t) = \ln(b_0) + t \ln(b_1)$
- iv. **Cubic Model:** $Y_t = b_0 + b_1t + b_2t^2 + b_3t^3$
- v. **Exponential Model:** $Y_t = b_0 e^{(b_1t)}$
- vi. **Logarithmic Model:** $Y_t = b_0 + b_1 \ln(t)$
- vii. **Growth Model:** $\ln(Y_t) = b_0 + b_1t$
- viii. **Inverse Model:** $Y_t = b_0 + \frac{b_1}{t}$
- ix. **Power Model:** $Y_t = b_0 t^{b_1} +$
- x. **S Type Model:** $Y_t = e^{(b_0 + \frac{b_1}{t})}$
- xi. **Logistic Model:** $Y_t = \frac{K}{(1 + e^{b_0 + b_1t})}$

Randomness Test

The current non-parametric test for randomization is based on the quantity of turning points. The method is to tally the series' peaks and troughs. A value that is higher than its two neighbors is referred to as a "peak," and a value that is lower than its two neighbors is referred to as a "trough." The series' highs and lows are regarded as pivotal moments. Therefore, one requires at least those data points to identify a turning moment. There are fewer turning points in the series than there are ups and downs.

Let us consider now a set of values $\mu_1, \mu_2, \dots, \mu_n$, and let us define a "marker" variable X_i by

$$X_i = \begin{cases} 1 & \text{if } \mu_i < \mu_{i+1} > \mu_{i+2} \\ & \text{if } \mu_i > \mu_{i+1} < \mu_{i+2} \end{cases} \quad \text{The number of turning points } p \text{ is then simply}$$

$$= 0 \quad \text{otherwise; } i = 1, 2, \dots, n-2$$

$$p = \sum_{i=1}^{n-2} X_i$$

Then,

$$E(p) = \sum E(X_i) = \frac{2}{3}(n-2)$$

$$E(p^2) = E\left(\sum_{i=1}^{n-2} X_i\right)^2 \text{ on simplification,}$$

$$E(p^2) = \frac{40n^2 - 144n + 131}{90} \text{ and}$$

$$V(p) = \frac{16n - 29}{90}$$

Measures of sustainability

According to Devi et al. (2021) and Vishwajith et al. (2018), sustainability is a complicated, multifaceted phenomenon that has been described in a variety of ways. While highly divisive it is widely accepted that it is complex and needs many evaluations. It can be evaluated by taking into account its biophysical, social, and economic characteristics in its most basic form.

Sustainability Index (SI)

For the sustainability index will divide the potato productivity data into two periods.

- Period -1 (1970 -1997)
- Period -2 (1998 - 2023)
- Whole Period (1970 – 2023)

(a) According to **Singh et al.**, (1990), the sustainability measure is as follows.

$$\text{Sustainability Index (SI)} = \frac{\bar{Y} - s}{Y_{max}}$$

(b) According to **Sahu et al.**, (2005) A sustainability index value that is closer to zero is a desirable value.

$$\text{SI} = \frac{Y_{max} - \bar{Y}}{\bar{Y}}$$

(c) According to **Pal and Sahu** (2007), sustainability increases when the sustainability index value decreases.

$$\text{SI} = \frac{S_i}{\bar{Y}_i} \cdot \frac{1}{S_{max}}$$

Modeling and forecasting

The plan looks like a flowchart for time series analysis-based potato production forecasts. The flow is described as follows:

1. Start The procedure's first action.
2. Data on potato production: This appears to be the initial step in which the pertinent data on potato production is recognized as a process input by using SPSS23.
3. Input Potato Production Data (India and Rajasthan): This stage indicates that data on potato production will originate from India and Rajasthan.
- 4- Training data: After the information is acquired, it is categorized as training data, which is what the forecasting models are fitted to.

5. The training data splits into two routes, signifying the application forecasting techniques: The statistical technique known as Autoregressive Integrated Moving Average (ARIMA) is widely used in time series forecasting.

6. Accuracy for training data: A number of accuracy metrics are used.

MAE: Mean Absolute Error, a measure of errors between paired observations expressing the same phenomenon;

RMSE: Root Mean Square Error, a measure of the differences between values predicted by a model and the values observed;

BIC: Bayesian Information Criterion, a measure of the relative quality of statistical models for a given set of data.

Mean Absolute Error (MAE): A measurement of inaccuracies between two observations that represent the same occurrence.


Root Mean Square Error (RMSE): a measurement of the discrepancies between data observed and values projected by a model.

Autocorrelation Function at Lag 1 (ACF1): A measure of the correlation between time series observations separated by a single time period.

Mean Absolute Percentage Error (MAPE): A measure of forecasting model prediction accuracy.

7. Projecting potato production: The more accurate forecasting model is probably chosen to project future potato production based on the evaluations.

- **Flow chart forecasting**



8- Finish: When the procedure is complete.

Autoregressive Moving Average (ARMA)

The model containing p autoregressive terms and q moving average terms is denoted by the notation ARMA (p, q). The AR(p) and MA(q) models are contained in this model.(Mishra et al 2023)

$$X_t = c + \varepsilon_t + \sum_{i=1}^p p_i X_{t-i} + \sum_{i=1}^q \theta_i \varepsilon_{t-i}$$

RESULT & DISCUSSION

According to the purpose of the analysis, the secondary source data was evaluated and interpreted.

Performance of Potato in India and Rajasthan during 1970-2023

In this section, descriptive statistics such as maximum, minimum, mean, median, skewness, and kurtosis are employed to elucidate the series' pattern and generate a consensus that is examined in Table-1. The data Platykurtic structure of potato under area, production, and yield in India and Rajasthan. Rajasthan and India have an average area of 5.59 and 1291.31 thousand ha of potato annually. Rajasthan and India produce an average of 74.69 and 24714.96 thousand tons of potato annually. Rajasthan and India have an average yield of 99.07 and 172.65 q/ha of potato annually. Skewness of 0.41 and 0.94 suggest that the area under potato in India and Rajasthan has been changing over time. positive skewness scores of 1.38 and 0.72 indicate that output increased at the start of the study and stayed almost constant for the duration of the investigation in Rajasthan and India. The positive skewness score was 0.03 and 0.27 in Rajasthan and India.

Table1: Performance of Potato area, production, and yield in India and Rajasthan during 1970-2023

India			
	Area ('000 ha)	Production ('000 t)	Yield (q/ha)
Mean	1291.319	24714.596	172.650
Standard Error	79.023	2277.440	6.483
Median	1206.900	20709.060	167.700
Kurtosis	-1.181	-0.737	-0.571

Skewness	0.414	0.729	0.273
Minimum	482.000	4451.000	88.200
Maximum	2345.400	61250.500	277.460
Rajasthan			
Mean	5.592	74.69557	99.077
Standard Error	0.649	12.812	8.040
Median	2.752	27.151	110.835
Kurtosis	-0.809	0.570	-1.083
Skewness	0.946	1.383	0.039
Minimum	1.3	3.1	13.3
Maximum	15.985	337.991	211.442

Trend Analysis of Potato in India and Rajasthan

The trend analysis shows (table-2) that there are cubic trends in potato area, yield, and production throughout India and Rajasthan. Suggesting that in the recent past, most likely, the series have reached their maximum values and then either remained constant or decreased, which is concerning. As can be seen from the data, the area dedicated to potato has grown annually in India and Rajasthan. Figures 1 to 6, show the non-linear patterns in India and Rajasthan.

Table-2: Trends of Potato in India and Rajasthan

India							
	Model summary			Parameter Estimates			
	equation	R ²	Sig.	constant	B ₁	B ₂	B ₃
Area	Cubic	0.980	0.000	519.639	10.237	0.588	-0.002
production	Cubic	0.976	0.000	4046.793	504.577	-5.371	0.298
Yield	Cubic	0.936	0.000	79.466	6.563	-0.188	0.002
Rajasthan							
Area	Cubic	0.902	0.000	3.708	-0.342	0.013	-4.859E-05
production	Cubic	0.902	0.000	8.613	-0.125	-0.047	0.003
Yield	Cubic	0.752	0.000	-6.880	6.541	-0.143	0.002

Figure-1: Trends of Area under Potato in India




Figure-2: Trends of Area under Potato in Rajasthan




Figure-3: Trends of production under Potato in India




Figure-4: Trends of production under Potato in Rajasthan




Figure-5: Trends of yield under Potato in India





Figure-6: Trends of yield under Potato in Rajasthan



Randomness Test: -In table-3 shows, that Rajasthan and India exhibit distinct trends in the area, production and yield of potato, as demonstrated by the randomization test.

Table-3 : That Rajasthan And India Exhibit Distinct Trends In The Area, Production And Yield Of Potato, As Demonstrated By The Randomization Test

Test of randomness	India			Rajasthan		
	Area	Prod	yield	Area	Prod	yield
No. of Observation	54	54	54	54	54	54
No. of Point (p)	21	27	29	24	17	27
E (P)	38.67	38.67	38.67	38.67	38.67	38.67
V(P)	10.34	10.34	10.34	10.34	10.34	10.34
t _{cal}	-5.49	-3.63	-3.01	-4.56	-6.74	-3.63
Inference	Trend	Trend	Trend	Trend	Trend	Trend
outliers Test	NO	NO	NO	NO	NO	NO

Sustainability analysis

Table-4 presents the results of measuring sustainability in the productivities of potato in Rajasthan and India using sustainability measurement.

India			
Sustainability Index	Period 1 (1970-1997)	Period 2 (1998-2023)	Whole Period (1970-2023)
SI 1	0.570	0.640	0.451
SI 2	0.414	0.316	0.607
SI 3	0.004	0.003	0.006
Rajasthan			
SI 1	0.092	0.459	0.189
SI 2	1.505	0.520	1.134
SI 3	0.013	0.005	0.010

Table-4: Sustainability potato yield measurement

According to indexes provided by SI-1 (Singh et al., 1990), SI-2 (Sahu et al., 2005), and SI-3 (Pal and Sahu, 2007), as measured by various formulas for accuracy, it is evident the entire country of India is highly sustainable in yield of potato in period-2 (1998-2023) and Rajasthan is highly sustainable in yield of potato in period-2 (1998–2023).

Modeling and forecasting

We used the Box-Jenkins methodology for this. The model was built using data from 1970 to 2020, and it was validated using data from 2020 to 2023. Future series predictions are made using the models that suit the data the best. None of them are stable in nature, as the ACF and PACF graphs from the original series make abundantly evident, and first-order differencing is enough to make them so. It was found that ARIMA models, starting with the model-building procedure given in the material and method section, ranging from (0, 1, 0) to (1, 1, 5) are appropriate for forecasting and predicting the production behavior of potato. using data from 1970 to 2023, the study then uses the differenced series to estimate ARIMA equations for all parameters. It then produces predictions up to 2030. However, ACF and PACF graphs are also used to perform diagnostic checks on residuals.

Figures-10 displays the ACF and PACF plot of the first difference, or the area under potato, for Rajasthan and India. These plots indicate that $p=1$ and $q=5$ for Rajasthan and $p=1$ and $q=5$ for India would be a reasonable range for the area under potato. Consequently, Table-5 demonstrates the ARIMA (1,1,5) areas under potato have the highest R^2 value and the lowest RMSE, MAPE, MaxAE, and MAE values in India. butthe ARIMA (1,1,5) areas under potato have the highest R^2 value and the lowest MAPE and MAE values in Rajasthan.The projected areas for potato in Table-6demonstrateRajasthan and India in 2023 were 13.51 thousand hectares and 2423 thousand hectares, respectively, while the actual areas were 13.60 thousand hectares and 2345.40 thousand hectares. India is predicted to have 2620 thousand hectares and Rajasthan will have 15.54 thousand hectares in 2030 respectively.Figure-11displays the ACF and PACF plot of the first difference, or the value of production under potato in Rajasthan and India.The ARIMA

(0,1,5) and ARIMA (0,1,5) were therefore determined to be appropriate for the Production of potato in Rajasthan and India. Table-5 further shows that India has the highest R^2 value along with the lowest values of MAPE, MaxAPE, and MAE, however, Rajasthan ARIMA (0,1,5) potato production has the highest value of R^2 along with the lowest values of RMSE, MAE, and MaxAE. In Table-7 contrast to the predicted 274.22 thousand tons and 63318.26 thousand tons, the production of potato in Rajasthan and India in 2023 was 260.50 thousand tons and 61250.50 thousand tons, respectively. Rajasthan is predicted to produce 301.31 thousand tons, while India is forecast to produce 69806.76 thousand tons in 2030. The yield under potato in Rajasthan and India, as indicated by the ACF and PACF plot of the first difference Figures-12. It may be concluded that ARIMA (1,1,3) and ARIMA (1,1,5) are the ARIMA models that are most suited for yield under potato in Rajasthan and India. Moreover, Table-5 examination showed that while Rajasthan had the highest R^2 value and the lowest MAE value, India yielded potato highest R^2 and lowest MAPE, MAE and Normalized BIC values across all of India. In Table-8 contrast to the predicted 264.13 q/ha and 191.33q/ha, the yields of India and Rajasthan in 2023 and yield in 2023 were 277.46 q/ha and 191.54q/ha, respectively. Rajasthan is anticipated to receive 209.25q/ha, while India is expected to receive 274.09 q/ha in 2030.

Table 5: Different ARIMA Model for area(“000”ha), production (“000”tons), and yield(q/ha) under potato in India and Rajasthan

India								
	ARIMA	R^2	RMSE	MAPE	MAE	MaxAPE	MaxAE	Normalized BIC
Area	(1,1,5)	0.990	62.696	3.577	42.235	16.938	189.079	8.876
Production	(0,1,5)	0.978	2600.236	9.818	1902.447	39.111	7002.540	16.176
Yield	(1,1,5)	0.922	13.916	5.962	9.633	27.145	40.554	5.790
Rajasthan								
Area	(1,1,5)	0.946	1.186	16.608	0.741	61.810	5.120	0.865
Production	(0,1,5)	0.930	26.366	55.422	14.353	276.300	91.400	6.994
Yield	(1,1,3)	0.852	23.616	24.279	15.962	106.590	71.750	6.698

Table-6: Model Validation and forecasting of Area (000*ha) under potato in India and Rajasthan

India			
year	Observed	Predicted	Prediction error percentage

2021	2225.75	2276.36	-2.27
2022	2332.05	2352.92	-0.89
2023	2345.41	2422.88	-3.30
2024		2413.99	
2025		2487.86	
2026		2474.06	
2027		2559.1	
2028		2547.16	
2029		2628.77	
2030		2620.14	
Rajasthan			
2021	14.10	14.71	-4.32
2022	14.50	14.45	0.34
2023	13.60	13.51	0.66
2024		14.32	
2025		14.62	
2026		14.59	
2027		14.86	
2028		15.08	
2029		15.31	
2030		15.54	

Table-7: Model Validation and forecasting of Production ('000' tons) of potato in India and Rajasthan

India			
year	Observed	Predicted	Prediction error percentage
2021	56175.75	57521.81	-2.39
2022	59160.34	60571.58	-2.38
2023	61250.50	63318.26	-3.37
2024		63527.71	


2025		65419.16	
2026		65502.66	
2027		67459.86	
2028		67654.86	
2029		69589.11	
2030		69806.76	
Rajasthan			
2021	258.20	264.65	-2.49
2022	262.50	270.85	-3.18
2023	260.50	274.22	-5.26
2024		281.81	
2025		280.31	
2026		285.88	
2027		286.75	
2028		291.61	
2029		296.46	
2030		301.31	

Table-8: Model Validation and forecasting of Yield (q/ha) under potato in India and Rajasthan

India			
year	Observed	Predicted	Prediction error percentage
2021	252.40	249.62	1.10
2022	273.56	265.24	3.04
2023	277.46	264.13	4.80
2024		267.81	
2025		261.24	
2026		262.94	
2027		264.48	
2028		268.45	

2029		270.55	
2030		274.09	
Rajasthan			
2021	183.12	185.62	-1.36
2022	181.03	183.85	-1.55
2023	191.54	191.33	0.10
2024		194.04	
2025		195.98	
2026		198.26	
2027		200.79	
2028		203.5	
2029		206.33	
2030		209.25	

Figure -7: Observed and Predicted Area (000'ha) under potato in India and Rajasthan






Figure-8: Observed and Predicted of Production ('000' tons)of potato in India and Rajasthan




Figure-9: Observed and Predicted of Yield (q/ha) under potato in India and Rajasthan





Figure -10: ACF and PACF graphs of residuals for the best-fitted models of Area under Potato in India and Rajasthan







Figure-11: ACF and PACF graphs of residuals for the best-fitted models of Production under potato in India and Rajasthan








Figure-12: ACF and PACF graphs of residuals for the best-fitted models of Area under potato in India and Rajasthan





Conclusion

Potato is the most important horticultural crops and has the maximum nutrition and dry matter production per unit area and time. It is the most important crop in the world for food. This study covering the period from 1970 to 2023, has analyzed potato area, production and productivity in the Rajasthan and India employing forecasting based box-Jenkins methodology after analyzing its trend for each individual series. Trend behavior of potato production there has been significant growth in area, production, and productivity. Analysis shows the cubic trend in potato area, production, and productivity in India and Rajasthan. Sustainability analysis is used for the future plan of crop improvement. The ARIMA models that are most suited for yield under potato Rajasthan would be anticipated to receive 209.25q/ha, while India would be expected to receive 274.09 q/ha in 2030.

References :-

1. Devi, M., Kumar, J., Malik, D.P. and Mishra, P. (2021). An Inter-District Analysis of Instability and Sustainability for Major Crops in Haryana. *Economic Affairs*, 66(2): 217-223.

2. Kumari, P., Vekariya, P., Kujur, S. N., Macwan, J., & Mishra, P. (2024). Predicting Potato Prices in Agra, UP, India: An H2O AutoML Approach. *Potato Research*, 1-16. <https://doi.org/10.1007/s11540-024-09726-z>
3. Mishra, P., Alhussan, A. A., Khafaga, D. S., Lal, P., Ray, S., Abotaleb, M., ... & El-Kenawy, E. S. M. (2024). Forecasting production of potato for a sustainable future: Global market analysis. *Potato Research*, 1-20: <https://doi.org/10.1007/s11540-024-09717-0>.
4. Mishra, P., Ghazi Al Khatib, A.M., Lal, P., Anwar, A., Nganvongpanit, K., Abotaleb, M., Ray, S. and Punyapornwithaya, Y. (2023). An Overview of Pulses Production in India: Retrospect and Prospects of the Future Food with an Application of Hybrid Models. *Natl. Acad. Sci*, 10 (2): 1267-1276.
5. Mishra, P., Sahu P.K., Dhekale, B.S. and Vishwajith, K.P. (2015). Modelling and forecasting of wheat in India and their yield sustainability, *Indian Journal of Economic Development*. 11(3): 637-647.
6. Mishra, P., Sahu, P.K., Padmanaban, K., Vishwajith, K.P. And Dhekale, B.S. (2015). Study of instability and forecasting of food grain production in India. *International Journal of Agriculture Sciences*. 7(3):474-481.
7. Pal, S. and Sahu. P.K. (2007) "On assessment of sustainability of crops and cropping system-some new measures". *Journal of sustainable agriculture*, 31(3):43-54.
8. Sahu, P. K., Das, M., Sarkar, B., VS, A., Dey, S., Narasimhaiah, L., ... & Raghav, Y. S. (2024). Potato production in India: a critical appraisal on sustainability, forecasting, price and export behaviour. *Potato Research*, 1-37: <https://doi.org/10.1007/s11540-023-09682-0>.
9. Sahu, P.K., Kundu, A.L., Mani, P.K. and Pramanicl, M. (2005). "Sustainability of different nutrient combination in a long-term rice-wheat cropping system". *Journal of New Seeds*, 7(3):91-101.
10. Singh, R. P., Das, S. K., Bhaskara, R.U. M. and Reddy, N. M. (1990). "Towards sustainable dryland Agricultural Practices". *Technical Bulletin, Central Institute for Dryland Agriculture, Hyderabad, India*, 1-81.
11. Vishwajith, K.P., Sahu, P.K., Mishra, P., Dhekale, B.S. and Upadhyay N. (2018). Crop yield sustainability: A few measures. *J. Pharmacogn. Phytochem.*, Sp(2): 247-253.

12. Yadav, S., Al khatib, A. M. G., Alshaib, B. M., Ranjan, S., Kumari, B., Alkader, N. A., ... & Kapoor, P. (2024). Decoding Potato Power: A Global Forecast of Production with Machine Learning and State-of-the-Art Techniques. *Potato Research*, 1-22. <https://doi.org/10.1007/s11540-024-09705-4>.

UNDER PEER REVIEW