

Evaluating the Protective Potential of Activated Charcoal Against Tannic Acid-Induced Hepatotoxicity in Rats

Abstract

Background: There is an increase in substance toxicity and abuse with tannic acid in developing countries in the past few years. Different substances, both chemical and traditional, have been used to neutralize or reduce the toxic effects of tannic acid poisoning. **Aim:** This study aimed to assess the protective potential of activated charcoal against tannic acid-induced liver poisoning in rats. **Methodology:** Twenty male Wistar rats were divided into five groups: negative control (NC), positive control (PC), and three activated charcoal treatment groups (ACG1, ACG2, and ACG3), each consisting of five rats. NC was neither induced with tannic acid nor treated with activated charcoal. PC was induced with 2260 mg/kg tannic acid but not treated. ACG1, ACG2, and ACG3 were induced with 2260 mg/kg tannic acid and treated with activated charcoal at concentrations of 0.01 mg/ml, 0.02 mg/ml, and 0.03 mg/ml, respectively, starting on the 6th day after induction. Blood samples were collected via cardiac puncture to assess liver function. **Results:** The results showed a significant increase ($p < 0.05$) in liver parameters (AST, ALT, ALP, and albumin) following tannic acid poisoning, while a significant reduction ($p < 0.05$) in the same parameters was observed after the administration of activated charcoal. **Conclusion:** The study demonstrates that tannic acid induces hepatotoxicity, while activated charcoal shows potential in mitigating the toxic effects.

Keywords: *biochemical parameters, enzymes, liver, toxicity*

Introduction

Tannic acid is a naturally occurring plant polyphenol and can be found in practically all aerial plant tissues. The tannins are applied widely, with uses ranging from tanning, known over millennia (Mediterranean since *ca.* 1500BC), through medicinal uses in the food industry. It is mainly used as a fixative of dyes and as a chemical intermediate and reagent in the manufacture of inks, rubber, wine, medicine, imitation horns and tortoise shells [1]. Ingestion of tannic acid can cause hardening of the gastrointestinal mucosa [2], which can result in reduced gastrointestinal absorption of nutrients as well as of xenobiotics.

Traditionally, tannins are considered to have anti-nutritional properties [3]. Tannins have greater tendency to form complexes with proteins than carbohydrates and other food polymers because of the strong hydrogen bond affinity of the carboxyl oxygen of the peptide group. Complexes formed by tannins and proteins have been reported to be responsible for growth depression, low protein digestibility, decreased availability of amino acid, increased fecal nitrogen and other toxic effects [4-7].

Activated charcoal is an odorless, fine powder which is black in color it has been known to be effectively used to cure overdose in emergency rooms[8]. Activated

charcoal possess lots of medicinal and other characteristics which shows their effectiveness in several fields in which includes cosmetics and many more [8]. One of the effective characteristics of activated charcoal showed toxin absorbing activities [9]. The formation of activated charcoal done after heating the common charcoal in the existence of gas which causes charcoal to produce pores and inside spaces so, with the help of these pores, the trapping of chemicals occurs through activated charcoal [10]. Reports have shown that it can effectively treat flatulence, poison, cholestasis, and reduce the level of cholesterol [11]. From research, it was concluded that people who used activated charcoal for long duration are prevented from many health-related problems. It not only prevented stomach and intestine problems, but also showed interaction and absorbent effects for drugs, toxins, bacteria, viruses, chemicals & fungus, which are present in water [12].

The benefits of tannin have increased the consumption of tannins especially as supplements. This may expose individuals to tannin toxicity. Activated charcoal had shown usefulness in health related problems.

There is an increase in substance toxicity and abuse with tannic acid in developing countries in the past few years. Different substances both chemical and traditional have been used by different individuals to neutralize or reduce the toxic effects from tannic acid poisoning. This study intends to evaluate the biochemical (hepatic) effects of activated charcoal on acute toxicity poisoning with tannic acid, thus establishing the level of its efficacy. This may be useful in the management of tannic acid poisoning.

Methodology

Study design and period

This was a randomized experimental design to evaluate the effects of activated charcoal on tannic acid-induced hepatotoxicity. The research spanned 21 days, including a 14-day acclimatization period and a 7-day experimental phase.

List 1-Grouping

Groups	Details
NC	Negative control: No tannic acid given and no activated charcoal given
NP	Positive control: 2260mg/Kg of tannic acid was given
ACG 1	Treatment group 1: Administered 2260mg/Kg and treated with 0.01mg/ml of activated charcoal
ACG 2	Treatment group 2: Administered 2260mg/Kg and treated with 0.02mg/ml of activated charcoal
ACG 3	Treatment group 3: Administered 2260mg/Kg and treated with 0.03mg/ml of activated charcoal

AC: Activated charcoal

After the acclimatization, the rats were administered with a single oral dose of 2260mg/Kg (sub-lethal dose which has been reported in some studies to induce hepatotoxicity [13]) of tannic acid as presented in Table 1 above. After the oral administration of tannic acid to the appropriate rats groups the rats were observed for 6

days. After which, activated charcoal was administered orally in a one-time dose-dependent fashion as described in the Table 1 above.

After the treatment period, the rats were sedated using chloroform and their blood samples were collected via cardiac puncture for biochemical analysis of liver function parameters [14].

Study Area and Population

The study was conducted at the Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Nigeria. The study population comprised 25 male albino rats weighing between 120g and 200g. The animals were housed in standard laboratory conditions, maintained with solid pellets and water, and allowed a 14-day acclimatization period before the experiment.

Ethical Consideration

Ethical approval for the study was obtained from the Animal Ethics Committee of Rivers State University, Port Harcourt (Approval No: RSU/REC/2024-018). The care and use of animals in this research adhered strictly to institutional and international guidelines for humane handling of laboratory animals.

Inclusion Criteria

Only male albino rats aged between 8 and 12 weeks were included, body weight ranging from 120g to 200g and Healthy rats confirmed by veterinary examination.

Exclusion Criteria

Female rats, male rats outside the specified age and weight range, rats showing signs of illness or abnormalities during acclimatization period and rats previously used in other experiment or exposed to medication were excluded

Sample collection and Analysis

By means of syringe and needle 5ml of blood samples were collected aseptically into lithium heparin bottles. The whole blood samples were spun using a centrifuge at 3000RPM for 5minutes to separate plasma which was stored at 2-6°C for further analysis. The plasma was collected into a plain bottle for analysis of protein, albumin, bilirubin, AST, ALT and ALP using biuret method [15] for protein estimation, Bromocresol green (BCG)[[16] method for albumin estimation, Jendrassik and grof method [17] for bilirubin estimation, Reitman-Frankel method [18] for AST and ALT, and King Amstrong method for ALP [14].

Statistical Analysis

Data were recorded using Microsoft Excel and analyzed using GraphPad Prism version 6.0. Descriptive statistics (mean \pm standard deviation) were used to summarize the data. Inferential analysis was performed using one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test to compare group means. A p-value of <0.05 was considered statistically significant.

RESULTS

The results of the mean blood biochemical parameters for the control groups are shown in Table 1. The result showed that there were significant differences in AST, ALP, ALT and albumin in the negative control group when compared with the positive control group ($p < 0.05$). However significant differences were not observed in protein and bilirubin parameters.

Table 1: Comparison of Biochemical Markers in Negative and Positive Control Groups

Parameter	Negative control	Positive control	P. Value	Inference
ALP (U/L)	20.33±1.53	40.67±3.786	0.0010	S
AST (U/L)	31.33±4.51	64.00±6.00	0.0017	S
ALT (U/L)	22.00±2.65	31.33±3.22	0.0178	S
Protein (g/L)	34.67±2.52	35.33±2.08	0.7415	NS
Albumin (g/L)	30.33±0.58	33.67±1.53	0.0241	S
T.BIL (umol/L)	2.280±0.99	3.990±0.99	0.1012	NS
C.BIL (umol/L)	1.710±0.00	2.850±0.99	0.1161	NS

The results of the mean blood biochemical parameters for the groups are shown in table 3. The result showed that there were significant differences in AST, ALP, ALT and albumin in the groups ($p < 0.05$). However significant differences were not observed in protein and bilirubin parameters.

Table 2 Comparison of Biochemical Parameters in The Study Groups

	NC	PC	ACG 1	ACG 2	ACG 3	F-value	p-value	Remark
DB	1.71±0.00	2.85±0.99	2.28±0.99	2.85±0.99	1.71±0.00	1.667	0.23	NS
TB	2.28±0.99	3.99±0.99	2.85±0.99	2.28±0.99	2.28±0.99	1.70	0.22	NS
ALB	30.33±0.57	33.67±1.53	30.33±0.58	32.00±1.00	31.67±1.53	4.53	0.024	S
PR	34.67±2.52	35.33±2.08	37.33±2.08	35.33±1.53	34.0±1.00	1.273	0.343	NS
ALT	22.0±2.65	31.3±3.22	25.33±2.51	25.67±3.51	25.67±0.57	4.66	0.02	S
ALP	20.33±1.53	40.67±3.8	32.33±3.77	33.67±0.58	28.67±1.53	24.67	<0.0001	S
AST	34.67±3.21	64.0±6.0	46.32±4.0	32.0±2.6	32.0±1.7	39.03	<0.0001	S

Key: S- Significant; NS-Non-significant; ACG: Activated Charcoal Group

Comparison of biochemical parameters as presented in Table 4, shows significant change ($p<0.05$) in AST, ALP, ALT and albumin when compared between negative control, positive control and treatment groups.

Table 3: Tukey's Multiple Comparisons for Biochemical Parameters

	AST (U/L)	ALT (U/L)	ALP (U/L)	Protein (G/L)	Albumin (G/L)	T. BIL (UMOL/L)	C. BIL (UMOL/L)
NC vs. PC	S	S	S	NS	S	NS	NS
NC vs. ACG1	S	NS	S	NS	NS	NS	NS
NC vs. ACG2	NS	NS	S	NS	NS	NS	NS
NC vs. ACG3	NS	NS	S	NS	NS	NS	NS
PC vs. ACG1	S	NS	S	NS	NS	NS	NS
PC vs. ACG2	S	NS	S	NS	NS	NS	NS
PC vs. ACG3	S	NS	S	NS	NS	NS	NS
S + 1 vs. ACG2	S	NS	NS	NS	S	NS	NS
S + 1 vs. ACG3	S	NS	NS	NS	NS	NS	NS
S + 2 vs. ACG3	NS	NS	NS	NS	NS	NS	NS

KEY: S- significant; NS-non-significant; NC-negative control; PC- positive control;

Discussion

The present study evaluated the effects of activated charcoal on hepatic parameters of tannic acid toxicity in albino rats. As presented in Table 1, there was a significant difference ($p<0.05$) in the concentrations of AST (aspartate aminotransferase), ALP (alkaline phosphates), ALT (alanine aminotransferase) and albumin between negative and positive control groups. Turkey's multiple comparison between these groups also presented significant increase in AST parameters. The significant increase in enzyme activities is an indication that tannic acid may be capable of inducing liver toxicity. Reports have shown that Tannic acid is capable of causing toxicity in rats as it is capable of causing hepatic necrosis, thus leading to the death of liver cells causing toxicity [19]. This result is in agreement with the work by Samanta and his colleagues in 2004 [19], who reported an increase in death of liver cells after exposure to tannic acid when administered to rats. However, there was no significant difference in protein and bilirubin concentrations. This may be due to the concentration of tannic acid and the duration of study.

AST (aspartate aminotransferase) is an enzyme that is found mostly in the liver, but also in the heart (cardiac muscle), skeletal muscle, kidneys, brain, and red blood cells.

When the liver is damaged, it releases AST into the bloodstream [20]. An AST blood test can help in the diagnosis of liver damage or disease. The significant increase in AST may indicate hepatitis, cirrhosis or other liver diseases which in this case may be caused by tannic toxicity. Although, high AST levels can also indicate heart problems or pancreatitis by Pratt & Kaplan 2000 [21]. However, the rise of all studied liver enzymes; AST, ALT and ALP is a clear implication of liver pathology.

Furthermore, the findings from this study in Table 3 showed a significant difference ($p < 0.05$) in the levels of almost all the liver function parameters among the different treatment groups signifying that activated charcoal may be capable of ameliorating the toxic effect which was induced by tannic acid as evident in the parameters (AST, ALT, ALP and albumin) on Table 3. This could suggest that activated charcoal has some protective effects over tannic acid toxicity.

However, Protein and bilirubin did not show significant different in the study groups in both toxicity induction and activated charcoal treatment. This may be due to administration of inadequate dose to elicit an effect in the parameters studied.

Conclusion

The study has shown that tannic acid caused changes in AST, ALP, ALP and albumins levels. The results obtained suggested that activated charcoal showed an effect in ameliorating the hepatotoxicity induced by tannic acid. This may be used in the management of tannic acid toxicity.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

1. Luckeneder P., Gavino J., Kuchernig R., Petutschnigg A. & Tondi G. (2016). Sustainable phenolic fractions as basis for furfuryl alcohol-based co-polymers and their use as wood adhesives. *Polymers*, 8, 396.
2. Robles S. (2014). Encyclopedia of Toxicology Reference Work. Mosby Elsevier. Third Edition • Pages 474-475.
3. Ojo MA. (2018). Changes in some antinutritional components and in vitro multienzymes protein digestibility during hydrothermal processing of *Cassia hirsutta*. *Preventive Nutrition and Food Science*, 23, 152–159.
4. Waghorn G. (2008). Beneficial and detrimental effects of dietary condensed tannins for sustainable sheep and goat production—progress and challenges. *Animal Feed Science Technology*, 147, 116–139.
5. Woodward S.L., Waghorn G.C., Watkins K.A. & Bryant M.A. (2009). Feeding birdsfoot trefoil (*Lotus corniculatus*) reduces the environmental impacts of dairy farming. *New Zealand Society of Animal Production*, 69, 179–183.
6. Dijkstra J, Oenema O, van Groenigen JW, Spek JW, van Vuuren AM, Bannink A. (2013). Diet effects on urine composition of cattle and N₂O emissions. *Animal*, 7, 292–302.
7. Grosse Brinkhaus A, Bee G, Silacci P, Kreuzer M, Dohme-Meier F. (2016). Effect of exchanging *Onobrychis viciifolia* and *Lotus corniculatus* for *Medicago sativa* on ruminal fermentation and nitrogen turnover in dairy cows. *J Dairy Science*. 99, 4384–4397.
8. Cooney, D. (2016). Activated charcoal: Antidote, remedy and health aid: TEACH Services, Inc. Carballa, M., Omil, F., Lema, J., Llompart, M. pharmaceuticals, cosmetics and hormones in a sewage treatment plant. *Water Research*, 38, 2918-2926.
9. Ramos, A., Fink, J., Hernandez, E. (1996). Prevention of toxic effects of mycotoxins by means of nonnutritive adsorbent compounds. *Journal of Food Protection*, 59, 631-641.
10. Luhmann, A., Kong, X., Tutolo, B., Ding, K., Saar, M. & Seyfried, J. (2013). Permeability reduction produced by grain reorganization and accumulation of exsolved CO₂ during geologic carbon sequestration: A new CO₂ trapping mechanism. *Environmental science & technology*, 47, 242-251.

11. Festi, D., Montagnani, M., Azzaroli, F., Lodato, F., Mazzella, G., Roda, A., Colecchia, A. (2007). Clinical efficacy and effectiveness of ursodeoxycholic acid in cholestatic liver diseases. *Current clinical pharmacology*, 2, 155-177.
12. Park, S., Ha, S., Yoon, J., Lee, S., Hong, K., Lee, E., Bae, D. (2009). Exposure to ethyl carbamate in alcohol-drinking and nondrinking adults and its reduction by simple charcoal filtration. *Food Control*, 20, 946-952.
13. Okolonkwo, B. N., Amadi, C. F., Chikwubike, U. O. and Nyenke, C. U. (2022). The Comparative Effect of Vitamin E + C on the Chronic Toxicity of Paraquat in Albino Rats (*Rattus norvegicus*). *European Journal of Medicinal Plants*, 33(6), 7-13.
14. Okolonkwo, B. N., Chukwubike, U. O. Amadi, C. F. and Nyenke, C. U. (2022). Therapeutic Effects of Vitamin E on Paraquat Induced Liver Toxicity in Male Albino Rats (*Rattus Norvegicus*). *Journal of Advances in Medicine and Medical Research*, 34(17),1-7.
15. Zheng K, Wu L, He Z, Yang B, Yang Y. Measurement of the total protein in serum by biuret method with uncertainty evaluation. *Measurement*. 2017 Dec 1;112:16-21.
16. Iwasaki T, Nakajima K, Hirowatari Y, Matsushita M. Evaluation of the two-point calibration bromocresol green method, showing reduced deviation from the bromocresol purple method in sera from patients with hypoalbuminemia. *Annals of Clinical Biochemistry*. 2023 Sep;60(5):320-7.
17. Kiuchi S, Ihara H, Osawa S. Realization of the total and direct reaction of ditau bilirubin via the use of a modified Jendrassik-Gróf diazo procedure. *Int J Anal Bio-Sci Vol*. 2019;7(1).
18. Giusti G, Ruggiero G, Cacciatore L. A comparative study of some spectrophotometric and colorimetric procedures for the determination of serum glutamic-oxaloacetic and glutamic-pyruvic transaminase in hepatic diseases. *Enzymologiabiologica et clinica*. 2017 Aug 11;10(1):17-38.
19. Samanta S, Giri S, Parua S. (2004). Impact of tannic acid on the gastrointestinal microflora. *MicrobEcol Health Disease*, 16(1), 32–34.
20. Zhang J, Song Q, Han X, Zhang Y, Zhang Y, Zhang X, Chu X, Zhang F, Chu L. Multi-targeted protection of acetaminophen-induced hepatotoxicity in mice by tannic acid. *International immunopharmacology*. 2017 Jun 1;47:95-105.
21. Pratt, D. S., & Kaplan, M. M. (2000). Evaluation of abnormal liver-enzymes results in asymptomatic patients. *New England Journal of Medicine*, 342(1), 1266–71.