

C-reactive protein in acute myocardial infarction: association with heart failure - about 118 cases

Comment [zs1]: C-reactive Protein Levels in Acute Myocardial Infarction and Their Association with Heart Failure: Analysis of 118 Cases

ABSTRACT

Introduction: Inflammation plays a key role in initiation and progression of atherosclerosis. A few inflammatory biomarkers have been associated with cardiovascular outcomes, both in healthy individuals and in patients with established coronary heart disease and high-sensitivity C-reactive protein (hsCRP) has been identified as a more important prognostic biomarker than low-density lipoprotein (LDL) cholesterol.

Aims: The study aimed to investigate whether variations in inflammatory markers at the time of an ACS event are linked to patient prognosis.

Materials and methods: An observational prospective study was conducted with 118 patients diagnosed with ACS, admitted to the intensive care unit at ERRAZI hospital from January 2024 to June 2024. The inclusion criteria were: patients over 18 years of age, with confirmed ACS (unstable angina, STEMI, or NSTEMI), and who provided informed consent. The outcome measure was the development of heart failure during follow-up. Statistical analyses were performed using Microsoft Excel and IBM SPSS software, with significance set at $p < 0.05$.

Results: The study involved 118 patients, with a median age of 61.5 years. The majority were male (94 men, sex ratio of 3.91). 94 patients had MI and 24 had unstable angina. Common risk factors included smoking (64 patients), hypertension (46 patients), and diabetes (58 patients). The study found the following outcomes: 50% of patients experienced pump failure. 22% had thrombolysis, with 9% experiencing failure. 84 patients received PCI, 15 had CABG, and 19 were treated medically due to complex coronary anatomy. Short-term cardiovascular mortality (<29 days) was 16%, with causes including cardiogenic shock, uncontrolled arrhythmias, and severe metabolic disorders. Platelet counts ranged from 54,000 to 710,000 but this was not significantly linked to ACS outcomes. However, hs-CRP levels were significantly elevated (median: 45.04), and elevated hs-CRP, white blood cell count, and serum cholesterol were associated with a higher probability of adverse cardiac outcomes.

Discussion: Cardiovascular disease (CVD), particularly coronary heart disease (CHD), is the leading cause of death globally. Inflammation plays a crucial role in the onset and progression of CHD, with CRP being a key inflammatory biomarker. Studies have shown that elevated hs-CRP levels at the time of an ACS event are associated with both short-term and long-term mortality, and CRP is an important predictor of adverse cardiovascular outcomes. Additionally, blood cell profiles (such as neutrophils and monocytes) have been shown to predict mortality.

While hs-CRP is linked to the severity of ACS events, its predictive power is modest, and its addition to traditional risk factors (age, smoking, cholesterol) provides only slight improvements in risk assessment.

Comment [zs2]: ACS: Acute Coronary Syndrome; you should write the abbreviation

Comment [zs3]: In the methodology section, you mentioned the age range as "from...to," but now you refer to "above 18." Please ensure consistency in your writing.

Comment [zs4]: you should write the abbreviation

Comment [zs5]: you should write the abbreviation

Those results suggest that anti-inflammatory therapies might reduce ischemic events after myocardial infarction.

Conclusion: CRP, particularly hs-CRP, has proven to be an important biomarker in ACS, with elevated levels correlating to a higher risk of heart failure both in the short and long term. While hs-CRP is not specific to heart failure, it provides valuable prognostic information and can help guide clinical decision-making. However, further research is needed to fully understand the role of CRP in post-myocardial infarction heart failure and to explore the potential benefits of CRP-targeted therapies in improving long-term cardiovascular outcomes.

Keywords: Acute coronary syndrome; new markers ; inflammation ; hs-CRP ; heart failure

1. INTRODUCTION :

Atherosclerosis is a slowly progressing condition that typically involves a prolonged, asymptomatic phase, which can eventually lead to acute coronary syndrome (ACS), presenting as either myocardial infarction (MI) or unstable angina (UA). ACS often represents a late-stage complication and is frequently the first clinical manifestation of underlying atherosclerotic disease.

A chronic, low-grade inflammatory process is widely recognized as a key factor in the development and progression of atherosclerosis, as well as in the acute thrombotic events that characterize ACS. Numerous studies have demonstrated a strong association between blood cell profiles, the level of inflammation at the time of admission, and the outcomes of ACS. These findings highlight the ability to differentiate between MI and UA outcomes and suggest that anti-inflammatory therapies may offer potential benefits in reducing ischemic events following myocardial infarction.

2. MATERIAL AND METHODS :

We performed an observational prospective study including 118 patients diagnosed with an ACS admitted to the intensive care unit with acute chest pain at ERRAZI hospital from January 2024 to June 2024

Inclusion criteria:

1. Patient above 18 years of age .
2. All patients gave their informed consent before taking part.
3. Having ACS including unstable angina, STEMI, NSTEMI enrolled as per ESC definition of MI 2023.

Exclusion criteria:

1. Not confirmed diagnosis of ACS.
2. Subjects not willing to give consent.
3. Patients suffering from CKD and cancer.
4. Patients lost in follow-up.

Comment [zs6]: The introduction must be no less than 400 words and should include in-text citations.

Comment [zs7]: Where are the references???????????

Comment [zs8]: The methodology section must include all the materials used, such as ELISA kits or the spectrophotometer.

Comment [zs9]: you should write the abbreviation at the first time at least

The development of a heart failure during the follow up period was our outcome measure.

Statistical analysis:

Statistical analysis was performed using the software microsoft excel office 2016 and IBM SPSS software for IOS version 22. The categorical variables were compared by chi square test.

Continuous variables were presented as mean (\pm SD) and were compared by unpaired test.

A probability value of <0.05 at 95% Confidence Interval (CI) was considered significant.

3. RESULTS AND DISCUSSION

RESULTS:

Results:

The study population comprised 118 patients with ACS. The age varies between 41 and 84yo with a median of 61.50 y.o \pm 8.89. 94 men were included in our study with a sex ratio= 3,91.94 were diagnosed with myocardial infarction and 24 with unstable angina. With regard to distribution of risk factors, smoking history was noted upon 64 patients, hypertension upon 46 patients, diabetes mellitus upon 58 patients.

Table 1: Clinical characteristics of patients at inclusion

Total (n=118)	HF (n=60)	Non HF(n=58)	P value
Age (years)>61 yo	42	28	0.016
Male	49	45	0.582
Myocardial infarction	56	45	
Unstable angina	4	13	0.015
Hypertension	24	26	0.596
Diabetes mellitus	33	25	0.196
Smoking (current)	36	29	0.275

Comment [zs10]: write years old

Comment [zs11]: write table1 at the end of this paragraph

Comment [zs12]: add chart for this table

In our study, the incidence of adverse events was observed as follows:

- **Pump failure:** 50 % of patients experienced pump failure.
- **Failed thrombolysis:** 22% of patients had thrombolysis of whom 09% had thrombolysis failure.

- 84 patients benefited of PCI, 15 benefited of CABG and 19 patients was referred to only medical treatment for several causes which the complexity of coronary's anatomy was the leading cause with a rate of 78%
- **Short term mortality (<29 days) due to cardiovascular causes:** 16% of patients died due to cardiovascular-related complications.

Short term total mortality was primarily due to cardiac disease mortality:10 patients due to cardiogenic shock, 6 patients due to arrhythmias non controlled and 3 patients due to severe metabolic disorders.

The platelet count variates between 54000 and 710000 with a median of 241000+/-104121. After analyzing the data on platelet, we found that platelet count was not significantly different between groups.

It was observed that hs-CRP level was significantly elevated among ACS cases. The Hs-CRP variates between 0.4 and 276 with a median of 45.04+/-65.76.

The median of serum cholesterol level is 1.58. The white cells count variates between a min 3340 and a max of 30110 with a median of 11350 +/-4758

Further doing multiple logistic regression analysis it was observed that probability of adverse cardiac outcome increases with increase in hs-CRP level, white blood cells count and high serum cholesterol.

Table 2: blood characteristics of patients at inclusion

TOTAL (N=118)	HF (N=60)	NON HF (N=58)	P VALUE
SERUM CHOLESTEROL >1.58	40	27	0.027
HS CRP>45.04	29	7	0.0001
PQ >241000	38	31	0.276
LEUCO >11350	31	15	0.004

Comment [zs13]: add chart

Discussion:

Cardiovascular disease (CVD) is the leading cause of death globally, with coronary atherosclerotic heart disease (CHD) being the primary contributor to these fatalities. Research has increasingly highlighted the significant role of inflammation in the onset and progression of CHD. In particular, C-reactive protein (CRP), a key biomarker of inflammation, has drawn considerable interest for its association with atherosclerosis (AS), CHD, and inflammatory processes.[1] Many studies were realized :

In an observational cohort sub-study CHAPS [2], which included a 10-year follow-up using registry data, we found that plasma inflammatory biomarkers (such as hsCRP, SAA, and fibrinogen) measured during the acute event were independently associated with long-term mortality, regardless of the initial ACS diagnosis. Additionally, blood cell profiles (e.g., neutrophils, monocytes) at the time of the acute event were predictive of both short- and long-term mortality. We also observed that recently identified inflammatory composite biomarkers (such as NLR and MLR) were linked to long-term mortality. These findings suggest that standard blood biomarkers of inflammation measured during the acute event can provide valuable prognostic information for assessing future risk.

Comment [zs14]: you must write the abbreviation

An other study, [3] proved that patients with high SYNTAX scores had significantly higher levels of age, glucose, fibrinogen, monocyte, and total cholesterol LDL, and lower levels of albumin and total protein. We found that FAR and monocyte levels were independent predictors of the high SYNTAX score.

Finally, a study published on JIMA[4], showed that the increase in hs-CRP levels is significantly higher in ACS patients, and it is strongly associated with an elevated risk of developing ACS events. Furthermore, it was observed that hs-CRP levels were directly correlated with the severity of the adverse outcome of ACS events otherwise the platelet count was not significant as a prognostic indicator of combined ACS events.

The relationship of Hs CRP levels to future vascular risk is not a new relation revealed. In the 1930s, CRP gained attention among cardiovascular researchers following reports in the mid-1990s that increased levels are associated with unstable angina and acute coronary ischemia.[5]

High-sensitivity CRP relates more closely to the risks of plaque rupture and vascular thrombosis than to the extent of underlying atherosclerotic burden. In terms of cardiovascular event-free survival, the risks associated with hsCRP are comparable with and independent of LDL-C. [6]

However, these effects are modest; addition of hsCRP testing to traditional factors, such as age, smoking, and blood pressure, has only modest impact on the C-statistic (a measure of discrimination) and on indexes of reclassification. Analyses from the Framingham Heart Study, Women's Health Study, EPIC-Norfolk, and MESA (Multi Ethnic Study of Atherosclerosis), reported net reclassification index values of 8% to 12%, but smaller effects were seen elsewhere [7].

The interpretation of hsCRP levels is relatively straightforward. Levels below 1 mg/L are considered optimal, indicating a low systemic inflammatory state and a lower risk of atherosclerosis. Levels ranging from 1 to 3 mg/L suggest a moderate vascular risk, while levels above 3 mg/L are associated with a higher vascular risk, especially when considered alongside other risk factors. If hsCRP levels exceed 10 mg/L, this may indicate a transient infectious process or an acute-phase response, in which case the test should be repeated after 2 to 3 weeks. For risk prediction, the lower value from repeated tests, rather than the average, should be used. Persistently elevated hsCRP levels, however, are not necessarily false positives and may indicate a significantly increased vascular risk[8–10].

To move forward, and propose interventions to reduce cardiovascular event rates; Initial interventions for patients with elevated hsCRP levels should focus on lifestyle changes, including a healthy diet, regular exercise, and smoking cessation. Additionally, evidence-based use of statins and aspirin can be considered for primary prevention in individuals with elevated hsCRP. However, it is important to note that while aspirin primarily exerts antiplatelet effects and statins primarily lower lipid levels, the available data do not directly support the notion that reducing inflammation alone will necessarily lead to a reduction in cardiovascular event rates[11,12].

Alternative anti-inflammatory strategies are also under investigation for chronic atherosclerosis or acute coronary syndrome. These include agents such as colchicine (commonly given for pericarditis), for which the nonblinded LoDoCo trial provided provocative preliminary data [13]; salsalate (an anti-inflammatory agent shown to have efficacy in diabetes) [14]; anakinra (an IL-1 receptor antagonist) [15]; and mitogen-activated protein kinase inhibitors (under investigation in the ongoing TIMI-60 trial of acute ischemia) [16]. Careful assessment of risks for infection are needed in anti-inflammatory trials, and it is important to recognize that prior studies of non-steroidal anti-inflammatory agents and tumor necrosis factor inhibitors have not shown vascular protection.

4.LIMITATIONS :

Our study has several limitations. First of all, the sample size was relatively small, which limits the statistical power and makes it difficult to draw definitive conclusions. Then , as an observational study, the strength of our findings may be subject to potential biases and limitations inherent in this design.

5.CONCLUSION

In conclusion, C-reactive protein (CRP) has emerged as an important biomarker in the context of acute coronary syndrom(ACS), with growing evidence linking elevated CRP levels to the development and progression of heart failure.

CRP, as a marker of systemic inflammation, reflects the underlying inflammatory processes that contribute to myocardial injury, tissue damage, and subsequent adverse cardiac remodeling. Elevated CRP levels in ACS patients are associated with an increased risk of heart failure, both in the short and long term, making it a valuable prognostic tool. Although CRP is not specific to heart failure, its measurement can help identify patients at higher risk for poor outcomes, guide clinical decision-making, and potentially inform the development of targeted therapies aimed at reducing inflammation.

However, further research is needed to fully elucidate the role of CRP in the pathophysiology of heart failure following AMI and to determine whether CRP-targeted interventions can improve long-term cardiovascular outcomes.

associated with an increased risk of heart failure, both in the short and long term, making it a valuable prognostic tool. Although CRP is not specific to heart failure, its measurement can help identify patients at higher risk for poor outcomes, guide clinical decision-making, and potentially inform the development of targeted therapies aimed at reducing inflammation.

However, further research is needed to fully elucidate the role of CRP in the pathophysiology of heart failure following AMI and to determine whether CRP-targeted interventions can improve long-term cardiovascular outcomes.

References:

1. Zhao X, Gao C, Chen H, Chen X, Liu T, Gu D. C-Reactive Protein: An Important Inflammatory Marker of Coronary Atherosclerotic Disease. *Angiology* [Internet]. 10 août 2024 [cité 18 nov 2024];00033197241273360. Disponible sur: <https://doi.org/10.1177/00033197241273360>
2. Odeberg J, Halling A, Ringborn M, Freitag M, Persson ML, Vaara I, et al. Markers of inflammation predicts long-term mortality in patients with acute coronary syndrome – a longitudinal cohort study [Internet]. 2024 [cité 2 nov 2024]. Disponible sur: <https://www.researchsquare.com/article/rs-4973727/v1>

Comment [zs15]: References must be written in APA format.

3. Seydel GS, Gunturk I, Akkaya H, Gunturk EE. The relationship between the new inflammatory markers and disease severity in patients with acute coronary syndrome. *Acta Cardiologica* [Internet]. 8 août 2024 [cité 18 nov 2024];79(7):778-86. Disponible sur: <https://doi.org/10.1080/00015385.2024.2403933>
4. Trivedi AS, Chakrabarti D, Khasnabish S, Bhattacharyya AK. High Sensitive CRP and Platelet Volume Indices in Acute Coronary Syndrome and their Impact in Prognostication. *Journal of the Indian Medical Association*. 2024;122(04).
5. Berk BC, Weintraub WS, Alexander RW. Elevation of C-reactive protein in "active" coronary artery disease. *The American Journal of Cardiology* [Internet]. 15 janv 1990 [cité 18 nov 2024];65(3):168-72. Disponible sur: <https://www.sciencedirect.com/science/article/pii/000291499090079G>
6. Ridker PM. A Test in Context. *Journal of the American College of Cardiology* [Internet]. févr 2016 [cité 18 nov 2024];67(6):712-23. Disponible sur: <https://linkinghub.elsevier.com/retrieve/pii/S073510971507549X>
7. Martin SS, Johnson AE, Blumenthal RS. 12 - Use of High-Sensitivity C-Reactive Protein for Risk Assessment. In: Ballantyne CM, éditeur. *Clinical Lipidology (Second Edition)* [Internet]. Philadelphia: W.B. Saunders; 2015 [cité 18 nov 2024]. p. 135-45. (Companion to Braunwald's Heart Disease). Disponible sur: <https://www.sciencedirect.com/science/article/pii/B9780323287869000128>
8. Vlachopoulos C, Xaplanteris P, Aboyans V, Brodmann M, Cífková R, Cosentino F, et al. The role of vascular biomarkers for primary and secondary prevention. A position paper from the European Society of Cardiology Working Group on peripheral circulation: Endorsed by the Association for Research into Arterial Structure and Physiology (ARTERY) Society. *Atherosclerosis* [Internet]. 1 août 2015 [cité 18 nov 2024];241(2):507-32. Disponible sur: <https://www.sciencedirect.com/science/article/pii/S0021915015013088>
9. Variability and Classification Accuracy of Serial High-Sensitivity C-Reactive Protein Measurements in Healthy Adults | *Clinical Chemistry | Oxford Academic* [Internet]. [cité 18 nov 2024]. Disponible sur: <https://academic.oup.com/clinchem/article/47/3/444/5639518>
10. Nasermoaddeli A, Sekine M, Kagamimori S. Intra-Individual Variability of High-Sensitivity C-Reactive Protein. *Circulation Journal*. 2006;70(5):559-63.
11. Rosuvastatin to Prevent Vascular Events in Men and Women with Elevated C-Reactive Protein | *New England Journal of Medicine* [Internet]. [cité 18 nov 2024]. Disponible sur: <https://www.nejm.org/doi/full/10.1056/NEJMoa0807646>
12. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, et al. Ezetimibe Added to Statin Therapy after Acute Coronary Syndromes. *New England Journal of Medicine* [Internet]. 18 juin 2015 [cité 18 nov 2024];372(25):2387-97. Disponible sur: <https://www.nejm.org/doi/full/10.1056/NEJMoa1410489>
13. Nidorf SM, Eikelboom JW, Budgeon CA, Thompson PL. Low-Dose Colchicine for Secondary Prevention of Cardiovascular Disease. *Journal of the American College of Cardiology* [Internet]. 29 janv 2013 [cité 18 nov 2024];61(4):404-10. Disponible sur: <https://www.sciencedirect.com/science/article/pii/S0735109712054782>
14. Salicylate (Salsalate) in Patients With Type 2 Diabetes: A Randomized Trial: *Annals of Internal Medicine*: Vol 159, No 1 [Internet]. [cité 18 nov 2024]. Disponible sur: <https://www.acpjournals.org/doi/10.7326/0003-4819-159-1-201307020-00003>
15. effect of interleukin-1 receptor antagonist therapy on markers of inflammation in non-ST elevation acute coronary syndromes: the MRC-ILA Heart Study | *European Heart Journal* | *Oxford Academic* [Internet]. [cité 18 nov 2024]. Disponible sur: <https://academic.oup.com/eurheartj/article/36/6/377/445879>
16. GlaxoSmithKline. A Clinical Outcomes Study to Compare the Incidence of Major Adverse Cardiovascular Events in Subjects Presenting With Acute Coronary Syndrome Treated With Losmapimod Compared to Placebo (PM1116197) LosmApimod To Inhibit p38 MAP Kinase as a Therapeutic Target and modify Outcomes After an Acute Coronary syndrome (LATITUDE)-TIMI 60. [Internet]. clinicaltrials.gov; 2017 avr [cité 18 nov 2024].

Report No.: NCT02145468. Disponible sur: <https://clinicaltrials.gov/study/NCT02145468>

Comment [zs16]: The link must be written correctly by pressing the spacebar on the keyboard so that it appears in blue.

UNDER PEER REVIEW