

Original Research Article

Mutation breeding for genetic improvement and the development of high yielding mutant variety of sesame (Binatil-3)

Abstract: Sesame plays an important role in human nutrition. Sesame seed contain fat, protein, carbohydrates, fiber and some minerals. The seeds of creamy seed-coat color of popular sesame variety Binatil-1 were irradiated with gamma rays to create genetic variability for earliness, higher seed yield and other desirable agronomic traits. The experiment was conducted at BINA headquarters farm, Mymensingh during March-May (Kharif-I) 2002 to 2007 and 47 mutants were primarily selected based on their better field performance including seed yield plant⁻¹ and other important agronomic traits. The mutant SM-10-04 (mutant of Binatil-1 with 500Gy of gamma rays) was found to be superior compared to other mutants and the mother variety. The evaluation trials were conducted during March-May (Kharif-I) season at different sesame growing areas of Bangladesh during 2008 to 2012. Significant variations were observed both in individual location and combined over locations for most of the traits. Reactions to major disease (stem rot) and insect-pests (hairy caterpillar, hawk moth and pod borer) infestations were also studied. BINA applied to the National Seed Board (NSB) of Bangladesh for registration of the superior mutant SM-10-04 as an important sesame variety. The NSB of Bangladesh registered the mutant SM-10-04 as an improved sesame variety naming Binatil-3 in 2013 for commercial cultivation in Bangladesh.

Key words: Binatil-3, Gamma irradiation, High yielding, Mutant, Sesame

1. Introduction

Sesame (*Sesamum indicum* L.) $2n=2x=26$ is a self-pollinated oilseed crop of the family Pedaliaceae. It is called the “Queen of oil seeds” because of its high oil content, delicious nutty aroma, and flavors (Johnson *et al.*, 1979) and is traditionally categorized as a healthy food in Asian countries (Miyake *et al.*, 2005). Sesame is grown for its seeds, which contain 45-55% high quality oil (Tepora and De la cruz 1991). Brown or black seeded are valued more for oil whereas, white seeded are rich in iron. According to ElKhier *et al.* (2008), the majority of sesame seeds are utilized to extract oil, with the remaining seeds being consumed for food.

The oil of sesame seed is renowned for its stability because it strongly resists oxidative rancidity even after long exposure to air (Global Agri Systems, 2010).

Sesame is a self-pollinated crop and often has lower genetic variability than out crossing crops (Charlesworth, 2003). Mutation is one of the popular methods to generate genetic variation (Acquaah, 2012, Malek *et al.*, 2014, 2016, 2020, 2022). Mutations occur randomly throughout the genome and within the locus or genes; thereby not only producing desired mutation but allowing the emergence of other mutations that can increase the diversity of a germplasm (De Vicente *et al.*, 2004). Induced mutations with gamma ray irradiation are often performed due to their wide availability and flexibility of use (Foster & Shu, 2012). According to Gafaar *et al.* (2016), gamma rays are a type of ionizing radiation that cause cells to produce free radicals, which can alter the components of the cell and affect various aspects of a plant's morphology, anatomy, biochemistry, and physiology depending on the dosage of gamma irradiation. Effect of gamma irradiation in diversification of genetic variation was recorded by Indriatama *et al.*, (2016) in wheat, Laskar & Khan (2017) in lentil and Malek *et al.*, (2014) in soybean.

The production of sesame is greatly depending on the cultivars, cultural techniques, and growing conditions. Sesame is a low yielder and worldwide average yield is only 535 kg ha⁻¹ (Brigham,

1985). Mutation breeding helps in creating variability not only in agro-morphological traits like yield and yield components, but also a better oil quality profile can be achieved in sesame (Kang *et al.*, 1998, Chowdhury *et al.*, 2009, Ong'injo and Ayiecho, 2009, Savant and Kothekar, 2011, Malek M.A. 2021). The major constraints in sesame production worldwide are lack of wider adapting cultivars, shattering of capsules at maturity, non-synchronous maturity, poor stand establishment, lack of fertilizer responses, profuse branching, and low harvest index (Ashri, 1994). Besides these, improvement of sesame has been slow mainly due to lack of adequate research and efficient breeding programs (Ashri, 1998). Therefore, the present investigation was undertaken with the objectives to determine the effects of gamma irradiation in sesame, the mutant-environment interaction on sesame yield and finally to develop a high yielding mutant variety of sesame.

2. Materials and methods

Binatil-3 (*Sesamum indicum* L.) has been developed from the sesame variety, Binatil-1. The seeds of Binatil-1 were exposed to different doses of gamma rays (500, 600, 700 and 800Gy) using Co⁶⁰ gamma source in mutation breeding laboratory at BINA, Mymensingh and grown as M₁ generation during March-May, 2002. During evaluation (March-May, 2003) of the irradiated population in M₂ generation, a large number of variants for different traits including seed yield were observed. Forty seven mutants were primarily selected in M₃ generation considering their better field performance including seed yield plant⁻¹ and other important agronomic traits as compared to the mother variety in March-May, 2004. These experimental materials were grown in M₄ and M₅ generations to study their breeding behavior in respect of seed yield and yield contributing traits to select elite mutants during 2005, 2006 and 2007.

Observation trial was conducted with a large number of M₅ mutant variants at BINA hqs. farm, Mymensingh and BINA sub-station farm at Ishurdi in 2008. After that, advanced yield trial was conducted in the BINA sub-station farms at Magura and Ishurdi during March-May, 2009 with five M₆ mutants (SM-03-04, SM-05-04, SM-08-04, SM-10-04 and SM-11-04) and two check varieties (Binatil-1 and BARI Til-2). Multi-location trials (on-station and on-farm) were conducted in various agro-ecological zones of the country in 2010, 2011 and 2012 in the experimental farms and also in the farmers' field following research management as well as farmers' management. On-station yield trial with advanced sesame mutants was conducted in the BINA sub-station farms at Magura and Ishurdi in 2010 with three M₇ mutants (SM-03-04, SM-10-04 and SM-11-04) and two checks (Binatil-1 and BARI Til-2).

On-station and on-farm yield trial with advanced sesame mutants was conducted in the farms of BINA sub-station at Ishurdi, Magura and Rangpur; and farmers' field at Jashore, Pabna and Rajshahi in 2011. Another on-station yield trial with advanced sesame mutants was conducted in the farms of BINA sub-station at Ishurdi and Magura in 2012. There were three advanced mutants (SM-10-04, SM-054 and SM-067) and two checks (Binatil-1 and BARI Til-2) were evaluated in 2011 and 2012. All these experiments were laid out in randomized complete block (RCB) design with three replications. Unit plot size was 20m² (4m × 5m) keeping 25cm spacing between rows and 6-8cm among the plants in a row. Seeds were sown in 1st week of March in 2011 and 2012. Recommended production packages like application of recommended doses of fertilizers, weeding, thinning, irrigation, application of pesticide etc. were properly followed to ensure normal growth and development of the plants. Data were taken for plant height, branches plant⁻¹, capsules plant⁻¹ and seeds capsule⁻¹ from 10 randomly selected plants from each plot. Maturity period was counted when 80% capsules were matured and most of the plants turned

into straw or yellowish color in each plot. Seed yield of each plot was recorded after proper drying and then converted into kg ha^{-1} . Reactions to major disease (stem rot) and insect-pests (hairy caterpillar, hawk moth and pod borer) infestations were also studied. Appropriate statistical analyses were performed for comparison of means of each character. Using the computer program Statistix 10, an analysis of variance was performed in accordance with the experimental design. 5% level of significance was used to compare mean differences among the treatments (Gomez and Gomez, 1984).

3. Results

Environmental effects on plant growth, development and yield is an important factor for crop improvement as well as development of new variety. Selection of desirable mutants was made from M_2 during 2003 and homogeneity was confirmed in M_5 generation during 2007. A large number of M_5 mutant variants were put into observation trial (non-replicated) at BINA headquarters farm, Mymensingh and BINA sub-station farm at Ishurdi in 2008. Five mutants, SM-03-04, SM-05-04, SM-08-04, SM-10-04 and SM-11-04 were primarily selected to put them into advanced yield trial on the basis of their desirable morphological characters and yield contributing traits compared to the mother variety.

The mean values for different characters of the mutants and checks for advanced yield trials with M_6 mutants during 2009 are presented in table 1. The results showed significant variation among the mutants and checks for most of the characters. Among the mutants and checks, mutant SM-10-04 took the shortest maturity period of 85 days and check BARI Til-2 took the highest maturity period of 95 days. Plant height ranged from 115cm (SM-10-04) to 133cm (BARI Til-2). The number of branches is one of the most important yield contributing traits in sesame. Number

of branches plant⁻¹ was highest in BARI Til-2 (2.6 per plant) followed by SM-10-04 (2.5), where Binatil-1 is unicum, i.e. non-branched. It is observed that, most of the mutants are branched but the mother variety Binatil-1 is unicum, i.e. non-branched. This result revealed that induced mutation can be changed the branching habit of sesame. The mutant SM-10-04 produced the highest seed yield (1671 kg ha⁻¹) followed by SM-11-04 (1584 kg ha⁻¹). These two mutants had also significantly higher number of capsules plant⁻¹. Number of seeds capsules⁻¹ ranged from 67 in BARI Til-2 to 93 in SM-03-04.

Table 1. Mean performance of M₆ sesame mutants and two checks for different quantitative characters at Magura in 2009

Mutants/ Varieties	Days to maturity	Plant height (cm)	No. of branches plant ⁻¹	No. of capsules plant ⁻¹	No. of seeds capsule ⁻¹	Seed yield (kg ha ⁻¹)
SM-03-04	88d	128b	0.7d	41c	93a	1411c
SM-05-04	90c	127b	1.0c	41c	83c	1062d
SM-08-04	92b	117c	1.3b	43bc	76d	1095d
SM-10-04	85e	115c	2.5a	53a	70e	1671a
SM-11-04	93b	129b	0.7d	48ab	89b	1548b
Binatil-1	88d	125b	0.0e	41c	90ab	1443c
BARI Til-2	95a	133a	2.6a	42bc	67f	1378c

*Same letter (s) in a column didn't differ significantly at 5% level of probability

The results of on-station yield trial during 2010 for each location and combined over locations are presented in table 2. The results showed significant variation among the mutants and two checks for all the characters at both the locations and combined over locations except plant height at Ishurdi. On an average, BARI Til-2 took the longest maturity period of 91 days and Binatil-1 took the shortest maturity period of 81 days. The mutants, SM-03-04, SM-10-04 and SM-11-04 took the maturity period of 87 days, 88 days and 88 days, respectively. The mutant SM-10-04 had the tallest plant (117cm) having non-significant difference with SM-11-04 (114cm) and SM-03-04 had the shortest plant (101cm). BARI Til-2 produced the highest number of branches (2.8) followed by SM-10-04 (2.5) while Binatil-1 was unicum. The mutant SM-10-04, Binatil-1 and BARI Til-2 produced the highest number of capsules plant⁻¹ (53) having non-significant difference with SM-03-04 (48). The mutant SM-10-04 produced the highest number of seeds capsule⁻¹ (73) which showed non-significant difference with Binatil-1 and SM-03-04. The lowest number of capsules plant⁻¹ was observed in BARI Til-2. Finally, the mutant SM-10-04 produced the highest seed yield of 1423 kgha⁻¹ followed by the check Binatil-1 (1322 kgha⁻¹) and another check BARI Til-2 produced the seed yield of 1236 kgha⁻¹.

Table 2: Mean performance of sesame mutants and two checks for different quantitative traits in 2010

Locations	Mutants/ Varieties	Days to maturity	Plant height (cm)	No. of branches plant ⁻¹	No. of capsules plant ⁻¹	No. of seeds capsule ⁻¹	Seed yield (kgha ⁻¹)
Magura	SM-03-04	91bc	92b	1.9c	63a	74b	1185b
	SM-10-04	89cd	109a	2.8b	58ab	73b	1328a

	SM-11-04	91b	104a	2.2c	57b	73b	1240b
	Binatil-1	88d	101ab	0.0d	49c	82a	1243b
	BARI Til-2	93a	101ab	3.2a	59ab	67c	1178b
Ishurdi	SM-03-04	84b	110a	0.0b	33c	69ab	1250c
	SM-10-04	88a	124a	2.2a	48b	64bc	1517a
	SM-11-04	85b	124a	0.0b	34c	74a	1150c
	Binatil-1	74c	116a	0.0b	58a	60c	1400ab
	BARI Til-2	89a	113a	2.4a	47b	59c	1293bc
Combined	SM-03-04	87b	101c	1.0c	48ab	71ab	1218c
(means	SM-10-04	88b	117a	2.5b	53a	73a	1423a
over two	SM-11-04	88b	114ab	1.1c	45b	69b	1195c
locations)	Binatil-1	81c	108bc	0.0d	53a	71ab	1322b
	BARI Til-2	91a	107bc	2.8a	53a	63c	1236c

*Same letter (s) in a column didn't differ significantly at 5% level of probability

On-station and on-farm yield trial conducted during 2011 showed significant variations for all the traits among the mutant and checks in each location and combined over locations except days to maturity. The average plant height in all locations revealed that the mutants SM-10-04, SM-058 and SM-067 were taller in height than Binatil-1 and only SM-067 was taller than BARI Til-2. The mutant SM-10-04 had higher number of branches plant⁻¹ as compared with variety Binatil-1 (unicum) and BARI Til-2. The higher number of capsules as well as higher number of seeds capsules⁻¹ in SM-10-04 resulted in significantly higher seed yield in all the locations. The

mutant SM-10-04 produced the highest seed yield of 1825 kg ha⁻¹ at Jashore and 1763 kg ha⁻¹ at Magura (Figure 1).




Figure 1. Mean seed yield performance of sesame mutants at different locations during 2011

The results of on-station yield trial during 2012 showed the significant variations for all the traits among the mutants and two checks in both locations except for days to maturity (Table 3). The tallest plant height was found in the mutants SM-058 and SM-067 at Ishurdi and Magura, respectively. The average number of branches ranged from 2.1-3.5, where the mutant SM-10-04 produced the highest number of branches. The number of capsules plant⁻¹ was significantly higher 78 in the mutant SM-10-04 followed by 66 in SM-058 and SM-067 in combined means over two locations. Bearing of three capsules axils⁻¹ is one of the important features of the mutant SM-10-04. The mutant SM-10-04 produced significantly higher number of capsules as well as higher number of seeds capsules⁻¹ which resulted in significantly higher seed yield in both locations (average yield 1505 kg ha⁻¹). The mutant SM-10-04 beard hairy stems which protect them from aphid infestation naturally.

Table 3. Mean performance of sesame mutants along with two checks for different quantitative traits during 2012

Locations	Mutant/ Varieties	Days to maturity	Plant height (cm)	No. of branches plant ⁻¹	No. of capsules plant ⁻¹	No. of seeds capsule ⁻¹	Seed yield (kg ha ⁻¹)
Ishurdi	SM-10-04	90c	93b	3.5a	88a	67a	1450a
	SM-058	95a	95a	3.3ab	73b	63c	1360b
	SM-067	92bc	90c	2.9b	70b	65b	1260c
	Binatil-2	91bc	87d	3.4a	61c	62c	1270c
	BARI Til-2	93b	90c	3.5a	72b	58d	1280c
Magura	SM-10-04	86c	107c	2.3b	69a	76a	1560a
	SM-058	92a	108c	2.5b	64b	69b	1460b
	SM-067	91ab	113a	2.1b	61b	66c	1420b
	Binatil-2	90b	104d	3.0a	56bc	69b	1275c
	BARI Til-2	89b	111b	3.1a	54c	63d	1325c
Combine d means over two locations	SM-10-04	88c	101a	2.9b	78a	72a	1505a
	SM-058	93a	102a	2.9b	66b	66b	1410b
	SM-067	91b	102a	2.5c	66b	66b	1340bc
	Binatil-2	90b	96b	3.2a	58d	65b	1273c
	BARI Til-2	91b	101a	3.3a	63c	61c	1302c

*Same letter (s) in a column didn't differ significantly at 5% level of probability

Overall performance for yield and yield related attributes of the new variety, Binatil-3 has been analyzed considering the mean seed yield of all trials conducting during 2009-2012. Four years seed yield results show that the new variety Binatil-3 produced the higher mean seed yield (1518 kg ha^{-1}) as well as year-wise seed yield as compared with two check varieties. Two checks, Binatil-1 and BARI Til-2 produced the seed yield (average of four years) of 1372 kg ha^{-1} and 1310 kg ha^{-1} , respectively (Figure 2).




Figure 2. Year-wise mean seed yield (kg ha^{-1}) of the new variety Binatil-3 as compared to checks

4. Discussion

Many crops have benefited greatly from induced mutation, including oilseed crops like sesame (Das and Haque 1997, Li and Chen 1998, Mehta and Singh 1998, Sorour et al. 1999, Govindarasu and Ramamoorthi 2000, Sheeba *et al.*, 2003, 2005, Chowdhury *et al.*, 2009, Diouf *et al.*, 2010, Begum and Dasgupta 2010, 2011, 2014). The increases in the number of capsule was due to an increase of flowers number and yield characters in some other variety developed using gamma rays were reported in mungbean (Tickoo and Chandra, 1999) soybean (Mehetre *et al.*, 1999, Malek *et al.*, 2014, 2022) and grass pea (Waghmare and Mehra, 2000). The increase in

the number of capsule plant⁻¹ among sesame mutants was reported by Hoballah (1999). Arslan et al. (2007) reported that the gamma-ray-induced mutants having improved plant architecture in sesame, including uniform growth habits, closed capsules, and resistance to Fusarium blight. The higher number of capsules as well as higher number of seeds capsules⁻¹ in SM-10-04 resulted in significantly higher seed yield in all the locations [54,55]. These results agreed with previous observations of Pathak and Dixit (1992) that reported a positive relationship between plant height, capsules plant⁻¹ and seed yield. Positively increase in seed yield due to the effect of gamma-rays induction was also reported earlier by many researchers likes in sesame and rapeseed-mustard (Malek *et al.*, 2012, Begum and Dasgupta, 2015, Malek *et al.*, 2020). Sureja and Sharma (2000), Reddy and Haripriya (1992) also reported a positive and highly significant relationship between number of seed capsule⁻¹, number of capsule plant⁻¹ and seed yield plant⁻¹. Seed yield is the most important trait for development of a promising crop variety. Mutants of oilseed crops having higher seed yield over mother variety also reported in sesame (Das *et al.*, 1999, Malek & Munshi, 2010, Aristya *et al.*, 2018, Malek, 2020), in rapeseed-mustard (Zhao et al., 2009, Ali et al., 2020, Mondal et al., 2020). Throughout the world including Bangladesh, the utilization of gamma ray induced mutation techniques for varietal improvement of sesame (Bhuiyan *et al.*, 2019, Malek, 2020) and other oilseed crops; particularly in soybean (Malek *et al.*, 2022), rapeseed-mustard (Malek *et al.*, 2016, Malek *et al.*, 2020, Bhuiyan *et al.*, 2021) is still underutilized. Moreover, the development of mutant variety Binatil-3 indicated the potentiality of evolving higher seed yielding variant through induce mutation.

5. Conclusion

The newly release mutant variety “Binatil-3” has been developed from Binatil-1. The seed coat colour of Binatil-3 is brownish but the seed coat colour of Binatil-1 is creamy and Binatil-1 is unicum but Binatil-3 is branched. This result revealed that induced mutations can be extensively and successfully used for the improvement of sesame in respect of changing branching habit and different in seed coat colour along with higher seed yield. The mutant also showed lower incidence of insect-pests infestation. There is an opportunity to increase cultivable land under sesame cultivation which can mitigate the challenges of increasing demand of edible oil in Bangladesh.

References

1. Acquaah G (2012). Principles of Plant Genetics and Breeding, 2nd ed. United Kingdom: WileyBlackwell Publishing. 740 p.
2. Ali MA, Bhuiyan SH, Rashid MH, Parveen S, Robbani M et al. (2016). Breeding for an ideal plant type in *Brassica rapa* L. Plant Knowledge Journal 5(1) 2016: 36-43.
3. Arslan Ç, Uzun B, Ülger S, Çağırman Mİ (2007). Determination of oil content and fatty acid composition of sesame mutants suited for intensive management conditions. Journal of the American Oil Chemists' Society 84: 917–920.
4. Ashri A (1994). Genetic resources of sesame: present and future perspectives. In: Arora R. K. & Riley K. W. (eds.) Sesame Biodiversity in Asia: Conservation, Evaluation and Improvement. IPGRI Office for South Asia, New Delhi, India: 25-39.
5. Ashri A (1998). Sesame breeding. Plant Breeding Review 16, 179-228.

6. Begum T, Dasgupta T (2010). A comparison on the effect of physical and chemical mutagenic treatments in sesame (*Sesamum indicum L.*). Genetics and Molecular Biology 33: 761-766.
7. Begum T, Dasgupta T (2011). Effect of mutagens on character association in sesame (*Sesamum indicum L.*). Pakistan Journal of Botany 43: 243-251.
8. Begum T, Dasgupta T (2014). Induced genetic variability, heritability and genetic advance in sesame (*Sesamum indicum L.*). SABRAO Journal of Breeding and Genetics 46: 21-33.
9. Begum, T, Dasgupta (2015). Amelioration of Seed Yield, Oil Content and Oil Quality Through Induced Mutagenesis in Sesame (*Sesamum Indicum L.*) Bangladesh Journal of Botany 44 (1), 15-22.
10. Bhuiyan MSH, M Malek, MMA Sarkar, M Islam, MW Akram (2019). Genetic variance and performance of sesame mutants for yield contributing characters. Malaysian journal of sustainable agriculture, 3 (2): 27-30.
11. Brigham RD (1987). Status of sesame (*Sesamum indicum L.*) breeding in the USA. Agronomy abstracts. Annual Meetings, p. 57. American Society of Agronomy. Madison, Wisconsin, (abstract).
12. Charlesworth D (2003). Effects of inbreeding on the genetic diversity of populations. Philosophical transactions of the Royal Society of London 358:1051–1070.

13. Chowdhury S, Datta AK, Saha A, Maity S (2009). Radiation-induced two oil rich mutants in sesame (*Sesamum indicum* L.). Indian Journal of Science and Technology 2: 51-52.
14. Das ML, Haque MM (1997). Response of sesame seeds to gamma ray and EMS in M₁ generation. Bangladesh Journal of Botany 26: 75-78.
15. De Vicente, MC, Lopez C, Fulton T (2004). Genetic Diversity Analysis with Molecular Marker Data: Learning Module. International Plant Genetic Resources Institute (IPGRI), Rome, Italy.
16. Diouf M, Boureima S, Diop T, Çağırkan M (2010). Gamma rays-induced mutant spectrum and frequency in sesame. Turkish Journal of Field Crops 15: 99-105.
17. El Khier MKS, Ishag KEA, Yagoub AEA (2008). Chemical Composition and Oil Characteristics of Sesame Seed Cultivars Grown in Sudan. Research Journal of Agriculture and Biological Sciences, 4(6): 761-766.
18. Forster BP, Shu QY (2012). Plant mutagenesis in crop improvement: Basic terms and applications. In: Shu, Q. Y., Forster, B. P. & Nakagawa, H. (Eds.). Plant Mutation Breeding and Biotechnology. CAB International & FAO. 9-20.
19. Gaafar RM, Hamouda M, Badr A (2016). Seed coat color, weight and eye pattern inheritance in gamma-rays induced cowpea M₂-mutant line. Journal of Genetic Engineering and Biotechnology 14 (1): 61-68.
20. Global Agri Systems (2010). Dehulled and roasted sesame seed oil processing unit. 18/08/11. [7] Hoballah, A.A. 1999. Selection and agronomic evaluation of induced

mutant lines of sesame. In induced mutations for sesame improvement. IAEA-TECDOC, IAEA, Vienna, pp.71-84.

21. Gomez KA, Gomez AA (1984). Statistical procedures for agricultural research. John Wiley & Sons. USA.
22. Govindarasu R, Ramamoorthi N (2000). Increased genetic variability following hybridization and mutagenesis in sesame. *Indian Journal of Genetics* 60: 251-253.
23. Hoballah AA (1999). Selection and agronomic evaluation of induced mutant lines of sesame. In induced mutations for sesame improvement. IAEA-TECDOC, IAEA, Vienna, pp.71-84.
24. Indriatama WM, Trikoesoemaningtyas SI Aisyah, Human S (2016). Pendugaan ragam genetik dan heritabilitas karakter agronomi gandum (*Triticum aestivum* L.) hasil berbagai perlakuan teknik iradiasi sinar gamma. *Jurnal Ilmiah Aplikasi Isotop dan Radiasi* 12(2): 79– 88.
25. Johnson LA, Suleiman TM, Lusas EW (1979). Sesame Protein: A Review and Prospectus. *Journal of the American Oil Chemists' Society* 56, 463–468.
26. Kamala T, Sasikala S (1985). Gamma-ray and colchicine induced mutants of TMV5' and 'IS 103' *Sesamum*. *The Indian Journal of Agricultural Sciences* 55: 151-155.
27. Kang CW, Rho JH, Lee SW, Park CB, Lee J et al. (1998). A high oil, linoleic fatty acid content and yielding new sesame mutant variety "Jangbaeckkae". *RDA Journal of Industrial Crop Science* 40: 83-88.

28. Laskar RA, Khan S (2017). Assessment on induced genetic variability and divergence in the mutagenized lentil populations of micro sperma and macrosperma cultivars developed using physical and chemical mutagenesis. PLoS ONE 12 (9): 1–18.
29. Lee JI, Kang CW, Lee ST (1984). Breeding of sesame for oil quality improved. III. Variation of fatty acid composition in gamma ray irradiated M₂ mutant populations. Research Report of the Office of Rural Development, South Korea Crop 26: 134-143.
30. Li Y, Chen QM (1998). Induction of male sterility mutation in sesame (*Sesamum indicum* L.) and a preliminary study of the genetics of the mutation. Chinese Journal of Oil Crop Science 20: 24 -27.
31. Majumdar SK, Barik KC, Bera PS, Ghosh DC (1987). Path coefficient analysis in sesame (*Sesamum indicum* L.) with varying levels of nitrogen and potassium. Indian Agriculturist 31: 165-169.
32. Malek M, Rafii M, Khatun M (2016). Selection of rapeseed mutants for higher yield and yield contributing traits. Journal of the Bangladesh agricultural university 14(2): 141-146.
33. Malek MA (2020). Binatil-2: a gamma rays induced mutant variety of sesame. Bangladesh Journal of Nuclear Agriculture 33 & 34: 151-158.
34. Malek MA, Begum HA, Begum M, Sattar MA, Ismail MR et al. (2012a). Development of two high yielding mutant varieties of mustard (*Brassica juncea* (L) Czern) through gamma rays irradiation. Australian Journal of Crop Science 6(5): 922-927.

35. Malek MA, Emon RM, Khatun MK, Bhuiyan MSH (2020). Yield performance of rapeseed mutant at M_8 generation. *Bangladesh Journal of Nuclear Agriculture* 33 & 34:73-82.
36. Malek MA, Emon RM, Khatun MK, Bhuiyan MSH, Nevame AYM et al. (2022). Binasoybean-6: A High Yielding Mutant Soybean Variety Developed through Sustainable Mutation Breeding. *Legume Research* DOI:10.18805/LRF-651.
37. Malek MA, Ismail MR, Monshi FI, Mondal MMA, Alam MN (2012b). Selection of promising rapeseed mutants through multi-location trials. *Bangladesh Journal of Botany* 41 (1), 111–114.
38. Malek MA, Rafii MY, Afroz MSS, Nath UK, Mondal MMA (2014). Morphological characterization and assessment of genetic variability, character association and divergence in soybean mutants. *The Scientific World Journal* 2014: 1-12.
39. Mehetre SS, Mahajan CR, Shinde RB, Ghatge RD (1999). Assessment of gamma induced genetic divergence in M_2 generation of Soybean. *Indian Journal of Genetics* 56: 186-190.
40. Mehta AK, Singh CB (1998). A gamma irradiated dichotomously branched plant type in *Sesamum*. *Plant Science* 11: 233-235.
41. Miyake Y, Fukumoto S, Okada M, Sakaida K, Nakamura Y et al. (2005). Antioxidative catechol lignans converted from Sesamin and Sesamino ITriglucoside by culturing with *Aspergillus*. *Journal of Agriculture and Food Chemistry* 53, 22–27.
42. Ong'injo EO, Ayiecho PO (2009). Genotypic variability in sesame mutant lines in Kenya. *African Crop Science Journal* 17: 101-107.

43. Pathak HC, Dixit SK (1992). Genetic variability and inter-relationship studies in black seeded sesame (*Sesamum indicum* L.). Madras Agricultural Journal 79: 94-100.
44. Reddy CDR, Haripriya S (1992). Genetic character association and path coefficient analysis in parents and their F1 sesame. Journal of Maharashtra Agricultural University 17: 55-57.
45. Savant KD, Kothekar VS (2011). Induction of variability in fatty acid profile in sesame (*Sesamum indicum* L.). Journal of Phytology 3: 1-3.
46. Sheeba A, Anbumalarmathi J, Babu S, Ibrahim SM (2005). Mutagenic effect of gamma rays and EMS in M₁ generation in sesame (*Sesamum indicum* L.). Research on Crops 6: 303-306.
47. Sheeba A, Ibrahim SM, Yogameenakshi P (2003). Induced chlorophyll mutation studies in sesame (*Sesamum indicum* L.). Sesame and Safflower Newsletter 18: 33-38.
48. Singh RK, Chaudhary BD (1979). Biometrical methods in quantitative genetic analysis. New delhi, India: Kalyani publishers.
49. Sorour WAI, Hussein MH, Ei-imam MA (1999). Gamma-ray induced mutations in sesame (*Sesamum indicum* L.) I. Selection of useful mutants. Bull Faculty Agric University of Cairo 50: 516-531.
50. Sureja AK, Sharma RR (2000). Genetic variability and heritability studies in garden pea (*Pisum sativum* L.). Indian Journal of Horticulture 57: 243 - 47.

51. Tepora NM, De La Cruz QD (1991). Mutation induction in sesame. Proc. International Symposium on the Contribution of Plant Mutation Breeding for Crop Improvement. IAEA, Vienna. Vol. 1, 371-374.
52. Tickoo JL, Chandra N (1999). Mutagen induced polygenic variability in Mungbean (*Vigna radiata* (L.) Wilczek). The Indian Journal of Genetics and plant Breeding 59(2): 193-201.
53. Waghmare VN, Mehra RB (2000). Induced genetic variability for quantitative characters in grass pea (*Lathyrus sativus* L.). Indian Journal of Genetics 60(1): 81-87.
54. Ali, Shahzeb, and Talekar Nilesh Suryakant. 2024. "Mutation Breeding and Its Importance in Modern Plant Breeding: A Review". Journal of Experimental Agriculture International 46 (7):264-75. <https://doi.org/10.9734/jeai/2024/v46i72581>.
55. Raina, Aamir, Rafiul Amin Laskar, Shahnawaz Khursheed, Ruhul Amin, Younas Rasheed Tantray, Kouser Parveen, and Samiullah Khan. 2016. "Role of Mutation Breeding in Crop Improvement- Past, Present and Future". Asian Research Journal of Agriculture 2 (2):1-13. <https://doi.org/10.9734/ARJA/2016/29334>.