

Impact of Biofertilizers on yield and Economics of *Rabi* sorghum (*Sorghum bicolor*L.) under rainfed condition.

Abstract: A field experiment was conducted at All India Coordinated Research Project on sorghum, Main Agricultural Research Station, University of Agricultural Sciences, Dharwad-580050, Karnataka for two consecutive *rabi* season (2018 and 2019) to study the Impact of Biofertilizers on growth, yield and Economics of *Rabi* sorghum (*Sorghum bicolor* L.) under rainfed condition in northern transition tract of Karnataka. The pooled results of the experiment indicated that, application of RDF (50:25:0 kg N:P₂O₅:K₂O/ha) +seed treatment with *Azospirillum* @4ml +PSB @ 4ml/kg seeds recorded significantly higher grain (42.3 q/ha) and fodder yield (8.4 t/ha) of *rabi* sorghum compared to rest of the treatments. The higher grain yield was due to significantly higher values of yield components viz., grain yield g/plant (40.0) and 1000-grain weight (33.3 g). Further, significantly higher gross returns (Rs.102133/ha), net returns (Rs.68232/ha) and BC ratio (3.0) were recorded due to application of RDF (50:25:0 kg N:P₂O₅:K₂O/ha) +seed treatment with *Azospirillum* @4ml +PSB @ 4ml/kg seeds compared to rest of the treatments.

Key words: Biofertilizers, seed treatment, *rabi* sorghum, Economics.

Introduction

Sorghum (*Sorghum bicolor* L. Moench) is one of the most important cereal crop in the world, popularly known as king of millets and great millet as an account of its large grain size among millet and vast area under it. It is grown in tropics and subtropics climates. Sorghum is called as camel crop because of drought, water logging and saline-alkaline tolerance. It is one of the most efficient C₄ plant in terms of photosynthesis. It is gaining importance as health food now a days because of its higher dietary fiber (2.56%). Apart from grains it provides valuable fodder which is one of the principle roughage feed in India.

Biofertilizer: The need of the hour in agriculture at present times, there is a growing concern about environmental hazards and threats to sustainable agriculture. In view of the above stated facts, the long term use of bio-fertilizers proves to be economical, ecofriendly, more efficient, productive and accessible to marginal and small farmers over chemical fertilizers. Biofertilizers have an advantage over chemical fertilizers, as they provide nutrients

Commented [GN1]: Symbol not matched and more over if we assume it is closing of important point/ head where is the opening symbol or vice versa. Please use appropriate punctuation

Commented [GN2]: Ha⁻¹ units should be expressed like this only.

Commented [GN3]: Remove space

Commented [GN4]: Remove space

Commented [GN5]: 4 ml kg⁻¹ al the units write like this please

Commented [GN6]: Remove space

Commented [GN7]: Units impproper

Commented [GN8]: Space

Commented [GN9]: Remove space

Commented [GN10]: Remove space

Commented [GN11]: Remove space

Commented [GN12]: Remove space

Commented [GN13]: Mention the units properly

Commented [GN14]: New Paragraph opening, hence give space and start

Commented [GN15]: Remove space

Commented [GN16]: healthy

Commented [GN17]: It also

Commented [GN18]: Remove space

Commented [GN19]: Remove space

Commented [GN20]: Remove space

Commented [GN21]: Remove space

Commented [GN22]: Remove space

Commented [GN23]: Remove space

Commented [GN24]: Remove space

Commented [GN25]: Remove space

Commented [GN26]: Remove space

Commented [GN27]: Remove space

Commented [GN28]: Remove space

Commented [GN29]: Remove space

Commented [GN30]: Remove space

Commented [GN31]: Remove space

Commented [GN32]: Remove space

Commented [GN33]: Remove space

in addition to plant growth promoting substances like hormones, vitamins, amino acids etc. (Shivankar *et. al*, 2000)

Material and methods

A field experiment was conducted at All India Coordinated Research Project on sorghum, Main Agricultural Research Station, University of Agricultural Sciences, Dharwad-580005, Karnataka, for two consecutive *rabi* season (2018 and 2019) to study the ² Impact of Biofertilizers on growth, yield and Economics of *Rabi* sorghum (*Sorghum bicolor* L.) under rainfed condition in northern transition tract of Karnataka. The soil of the experimental site was clayey soils. The experiment consisted of six treatments viz. T₁: Recommended dose of fertilizer (RDF) alone (Control), T₂: RDF + Seed treatment with Azospirillum @ 50g+PSB @ 50g/kg seed, T₃: RDF + Seed treatment with Azospirillum @ 2ml+PSB @ 2ml/kg seed, T₄: RDF + Seed treatment with Azospirillum @ 2ml+PSB @ 4ml/kg seed, T₅: RDF + Seed treatment with Azospirillum @ 4ml+PSB @ 2ml/kg seed and T₆: RDF + Seed treatment with Azospirillum @ 4ml+PSB @ 4ml/kg seed. The treatments were evaluated in Randomized Block Design (RBD) with four replications. The seeds of *rabi* sorghum (var. CSV-29R) was sown with a row spacing of 45 cm x 15 cm and recommended dose of fertilizers was applied to soil in each treatment and seeds were treated with both powder and liquid form of biofertilizers (*Azospirillum* and PSB) at the time of sowing. Sorghum yield attributes, grain and stover yields were recorded and economics were worked. The data obtained on *rabi* sorghum was subjected to statistical analysis by using MSTAT-programme according to Gomez and Gomez (1984).

Results and discussion

The data on grain yield (g/plant), 1000-grain weight (g), grain yield (q/ha) and stover yield (t/ha) of *rabi* sorghum were differed significantly due to various treatment. The results of pooled data indicated that, significantly higher grain yield (40 g/plant), 1000-grain weight (33.3g), grain yield (42.3q/ha), stover yield (8.4 t/ha) were recorded in seed treatment with *Azospirillum*@ 4ml +PSB @ 4ml /kg seeds along with RDF (50:25:0 kg N:P₂O₅:K₂O/ha) (T₆) as compared to control and powder form of biofertilizers. However, lower levels of liquid form of biofertilizers were on par with each other but significantly superior over control and powder form of biofertilizers. The per cent increase in grain yield of *rabi* sorghum with liquid form of biofertilizers over powder form of biofertilizers was an extent of 3 to 11. Further, The per cent increase in grain yield of *rabi* sorghum was 5.3 and 16.4 with powder and liquid form of biofertilizers respectively over RDF alone. The use of biofertilizers may lead to

Commented [GN34]: Starting word of the paragraph space with tab please

Commented [GN35]: Symbol not matched and more over if we assume it is closing of important point/ head where is the opening symbol or vice versa. Please use appropriate punctuation

Commented [GN36]: Remove 2 spaces

Commented [GN37]: Give space

Commented [GN38]: Mention the quantity of NPK used under RDF. You have mentioned in the abstract. Write or include whether any FYM quantity under RDF in clear

Commented [GN39]: Spacings are not proper and writing of the units follow as mentioned in Abstract.

Commented [GN40]: Space between two words

Commented [GN41]: Remove space

Commented [GN42]: space

Commented [GN43]: convert the units

Commented [GN44]: space

higher availability of nitrogen and phosphorus that promoted growth and development and ultimately resulting in higher yields (Raja and Takankharb, 2017) and (Jat *et al.*, 2013).

Application of RDF (50:25:0 kg N:P₂O₅:K₂O/ha) + seed treatment with *Azospirillum* @ 4ml +PSB @ 4ml /kg seeds recorded significantly higher gross returns (Rs.102123/ha), net returns (Rs.68232/ha) and BC ratio (3.0) compared to rest of the treatments followed by RDF (50:25:0 kg N:P₂O₅:K₂O/ha) + seed treatment with *Azospirillum* @ 4ml +PSB @ 2ml /kg seeds. This might be due to higher grain and stover yields of sorghum with these treatments. These results were in conformity with the findings of Patel, 2015.

Commented [GN45]: Units are improper and check spacings

Conclusion

Based on the experimental results it could be concluded that, all the yield and yield parameters of *rabi* sorghum were significantly superior with liquid form of biofertilizers over powder form of biofertilizers and gave higher gross and net returns.

References

- Gomez, K. A. and Gomez, A. A., 1984, Statistical procedures for agricultural research, 2nd ed. John Wiley and Sons. New York, P. 639.
- Jat, M.K., Purohit, H. S., Bahadur Singh, Garhwal, R.S and Mukesh Choudhary 2013. Effect of integrated nutrient management on yield and nutrient uptake in sorghum (*Sorghum bicolor*). Indian Journal of Agronomy. 58 (4): 543-547.
- Patel, B.K., 2015. Effect of land configuration and integrated nutrient management on productivity of different varieties of sorghum (Rabi) grown on fluventicustochets. Ph.D. thesis submitted to NAU, Navsari. (Gujarat). 2015.
- Raja, D. and Takankharb, V.G., 2017. Effect of Liquid biofertilizers (Bradyrhizobium and PSB) on availability of nutrients and soil chemical properties of Soybean (*Glycine max* L.). International Journal of Pure and Applied Bioscience. 5 (5): 88-96.
- Shivankar, S.K., Joshi, R.P. and Shivankar, R.S., 2000. Effect of biofertilizers and levels of nitrogen and phosphorus on yield and uptake by wheat under irrigated condition. *J. Soils & Crops*, 10(2): 292-294

Table 1 : Effect of powder Vs. liquid form of biofertilizer *onrabi* sorghum (CSV-29R) (Pooled of 2018 and 2019).

Commented [GN46]: Spacings for different words

Treatments	Grain yield (g/panicle)	Test weight (1000-grain) (g)	Grain yield (q/ha)	Fodder yield (t/ha)	Gross returns (Rs/ha)	Net returns (Rs/ha)	B:C
T ₁ : RDF(50:25:0 kg NPK/ha) alone	29.6	29.7	35.4	7.6	86023	52123	2.5
T ₂ :RDF+ seed treatment with Azospirillum @ 50g+ PSB @ 50g /kg seeds	33.8	29.4	37.4	8.1	90913	56983	2.7
T ₃ : RDF+ seed treatment with Azospirillum @ 2 ml+ PSB @ 2ml/kg seed	36.1	31.9	38.7	8.2	93900	60019	2.8
T ₄ :RDF+ seed treatment with Azospirillum @ 2 ml+ PSB @ 4 ml/kg seed	36.2	31.4	38.5	8.3	93369	59984	2.8
T ₅ : RDF+ seed treatment with Azospirillum @ 4 ml+ PSB @ 2 ml/kg seed	31.5	31.1	40.3	7.9	96953	63066	2.9
T ₆ : RDF+ seed treatment with Azospirillum @ 4 ml+ PSB @ 4 ml/kg seed	40.0	33.3	42.3	8.4	102123	68232	3.0
S.Em+ /	1.57	0.86	0.35	0.13	746	739	0.02
CD at 5%	4.73	2.60	1.07	0.40	2248	2229	0.07

Commented [GN47]: Not included cost of cultivation.

UNDER PEER REVIEW

