

Field Screening for Late Leaf Spot resistance in the BC₁F₄ population in Groundnut (*Arachis hypogaea* L.)

Abstract

Globally, late leaf spot (LLS), a foliar fungal disease is one of the most important biotic constraint in groundnut production. LLS is a major cause of yield loss accounting for over 70% in groundnut production. This study aimed to evaluate 115 BC₁F₄ groundnut lines for resistance to LLS long with parents was done at ICRISAT in Rainy 2024 (*Kharif*). The field screening was taken in Single plant progeny rows with 2m x 1 row and disease scoring was done at 75 Days After Planting (DAP), 90 DAP and 105 DAP in that progeny rows under natural disease conditions based on visual observation. Two lines BC₁F₄-1 and BC₁F₄-2 with score 4 at 105 Days after planting (DAP) was identified as resistant and four lines BC₁F₄-3, BC₁F₄-4, BC₁F₄-63 and BC₁F₄-65 with score 5 and 6 at 105 DAP exhibited moderate resistance were identified. In the present study out of 115 BC₁F₄ lines evaluated two lines showed resistance, and four lines showed moderate resistance. These lines will be advanced to the next generation and tested in multi-location trials to further assess their resistance and potential for use in breeding programs

Key words: Late Leaf Spot, Biotic constraint, Groundnut production and Multi-location

INTRODUCTION

Groundnut (*Arachis hypogaea* L.), is often hailed as the "king of oilseeds." Globally, cultivated in 112 countries with a total area of 30.5 million hectares (m ha) and total production of 54.2 million tonnes (m t) having productivity 1776 kg/ha and in India the total area of 5.7 (m ha) and total production of 10.1 (m t) with productivity 1776 kg/ha. (FAOSTAT, 2023). Groundnut fosters soil health through symbiotic nitrogen fixation with Rhizobia bacteria, promoting sustainable agricultural practices. Groundnut cultivation faces a formidable challenge from fungal diseases like late leaf spot (LLS), rust, Alternaria blight, stem rot, dry root rot and collar rot (Pal *et al.*, 2014, Joshi *et al.*, 2020.). LLS disease caused by *Phaeoisariopsis personata* pose the most significant threats. These widespread fungal pathogens inflict severe damage on groundnut crops. In most cases, these two pathogens occur together and hamper groundnut production (Kishore *et al.*, 2005). Globally, late leaf spot (LLS), a foliar fungal disease is one of the most important biotic constraint in groundnut production (Wankhade *et al.*, 2023). In India, these diseases have been documented to cause yield losses exceeding 70%, significantly impacting groundnut productivity (Subrahmanyam *et al.*, 1995, Ibrahim 2010). At ICRISAT, breeding for foliar fungal disease has resulted in development of several genotypes with high level resistance to rust and moderate resistance to LLS (Singh *et al.*, 2003). An estimated global yield loss of US\$600 million due to LLS was reported (Dwivedi *et al.*, 2003). The detrimental effects extend beyond yield reduction, as the quality of groundnut seeds is also compromised by these fungal infections.

Late leaf spot disease LLS is caused by the fungus known a *Nothopassalora personata* (Berk. & M.A. Curtis) poses a significant challenge in groundnut cultivation. While chemical treatments have been recommended for the treatment of LLS, they are often costly

and unsafe in certain situations and locations (Chandra *et al.*, 2005; Muhammad & Bdliya, 2011, Khan *et al.*, 2014). The frequency of infection is notably influenced by temperature and relative humidity (Shew *et al.*, 1989, Waliyar *et al.*, 1995). While latent period, lesion size and spore production have been the components most commonly associated with genetic resistance (Chiyembekeza *et al.*, 1993, Walls *et al.*, 1985). In contrast, host plant resistance, offers a safe, efficient and environmentally friendly way of controlling LLS (Ibrahim, 2010). application of host plant resistance further improves the yield and quality of groundnuts produced by farmers (Shoba *et al.*, 2012). However, the nature of the inheritance of LLS resistance is complex anpolygenic (Dwivedi *et al.*, 2002), thus making the identification of resistant and susceptible lines unreliable through conventional screening techniques (Leal-Bertioli *et al.*, 2009). With good polymorphic molecular markers, a breeder can easily identify LLS-resistant and susceptible groundnut plants (Mac *et al.*, 2006).

2.0 Materials and methods

2.1 Experimental Site and Plant Material

In the present study, the field evaluation of the backcrossed lines and parents was carried out during the rainy season 2024 at red precision 2C field-block (17.51° North latitude and 78.27° East longitude). The soil type of the experimental block was Alfisols (Patancheru Soil Series; UdicRhodustolf) with a pH ranging from 7.0 to 7.5. The air temperature ranged between 22.6° C to 30.6° C, bright sunshine 4.1 Hrs, relative humidity varied between 60 to 90 % and rainfall 201.1mm during *kharif* 2024. The details of the material used and the techniques adopted in the present investigation for recording of observations and analysis of data are briefly presented in this chapter. The experimental material for the present study comprised of BC₁F₄ lines were derived from the donor parent (ICGV 201009) which has score 4 at 105 DAP (unpublished data) which is resistant and recipient parent (Narayani) which has score 9 at 105 DAP, which is susceptible (Subrahmanyam *et al.* 1995).

2.2 Late Leaf Spot Disease Screening

The BC₁F₄ lines, their parents and checks were raised in single plant progeny rows and evaluated for late leaf spot disease during *kharif* 2024 (Fig. 1). Screening of Late leaf spot disease was carried out by visual screening method Fig. 2 using modified 9-point scale for late leaf spot (Table 1) given by Subrahmanyam *et al.* 1995.

2.3 Disease Scoring for Late Leaf Spot

For late leaf spot disease screening, visual screening (Fig. 2) and modified 9-point scale as given by Subrahmanyam *et al.* 1995 (Table 1) was used. The visual scores (1-9) and the extent of leaf area destroyed (0-100 %) are linearly related to each other. Disease scoring was done at 75, 90 and 105 days after sowing (pod filling stage). The disease scores were mainly based on the extent of leaf area damage.



Fig.1. Field view of BC₁F₄ population




Fig. 2. Standard reference for scoring late leaf spot disease resistance (Subrahmanyam *et al.*, 1995)

Table 1. Modified 9-Point scale used for field-screening of groundnut genotypes for Late Leaf Spot resistance (Subrahmanyam *et al.*, 1995)

Disease score	Phenotype description	Disease severity (%)
1	No disease	0
2	Lesions present largely on lower leaves; no defoliation	1-5%
3	Lesions present largely on lower leaves, very few on middle leaves; defoliation of some leaflets evident on lower leaves	6-10%
4	Lesions present on lower and middle leaves but severe on lower leaves; defoliation of some leaflets evident on lower leaves	11-20%
5	Lesions present on lower and middle leaves, over 50 % of defoliation of lower leaves	21-30%
6	Severe lesions on lower and middle leaves; lesions present but less severe on top leaves; extensive defoliation of lower leaves; some defoliation on middle leaves	31-40%
7	Lesions on all leaves but less severe on top leaves; defoliation of all lower and middle leaves	41-60%
8	Defoliation of all lower and middle leaves; severe lesions on top leaves evident	61-80%
9	Almost all leaves defoliated, leaving bare stem; some leaflets may remain but show severe leaf spot	81-100%

3.Results and discussion

3.1. Superior Selected lines for LLS Disease in the BC₁F₄ Generation

Among the 115 BC₁F₄ Single plant progeny lines, two lines BC₁F₄-1 and BC₁F₄-2 with score 4 at 105 DAP were identified as resistant to LLS disease and four lines BC₁F₄-3, BC₁F₄-4, BC₁F₄-63 and BC₁F₄-65 with score 5 and 6 at 105 DAP showed moderate resistance to LLS disease. Remaining 109 lines were with score 7 to 9 at 105 DAP showed highly susceptible to LLS disease.

Table 2. Resistance lines and their scores at different days after planting(DAP)

S.No	Genotypes	75 DAP	90 DAP	105 DAP	% LAD	R/MR/S *
1	BC ₁ F ₄ -1	2	3	4	20	R
2	BC ₁ F ₄ -2	2	3	4	20	R
3	BC ₁ F ₄ -3	3	4	5	30	MR
4	BC ₁ F ₄ -4	3	4	5	30	MR
5	BC ₁ F ₄ -63	3	5	6	40	MR
6	BC ₁ F ₄ -65	3	5	6	40	MR
7	ICGV 201009	2	3	4	20	R
8	Narayani	4	6	8	80	S

DAP-Days after planting, **% LAD**- Percentage of leaf area damage

* **R**- Resistance, **MR**- Moderate resistance and **S**- susceptible

3.2 Discussion

The results of this study provide valuable insights into the symptomatology, disease severity under natural conditions of the late leaf spot (LLS). The BC₁F₄ lines exhibited varying responses to LLS under natural conditions. The low disease score is an important indicator of disease dynamics, exhibited significant variability among different groundnut cultivars in field screening. These observations highlight the complexity of host-pathogen interactions in LLS and reinforces the significance of resistant cultivars in managing LLS outbreaks.

Conclusion

In the present investigation, 115 BC₁F₄ Single plant progeny lines were evaluated along with parents and one check to identify superior lines for LLS disease. Results depicted that among the 115 lines, 2 lines showed consistence performance both at 75th, at 90th and at 105th days after planting normal and disease plots. Therefore, these identified lines will be forwarded for next generation and then after multi-location trails for disease resistance.

Disclaimer (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies, such as large Language Models (chatGPT, COPILOT, etc) and text-to-image generations have been used during writing or editing of this manuscript.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

Option 1

Reference:

- Chandra, B., Viswavidyalaya, K., & Bengal, W. (2005). Management of leafspot (*Cercospora arachidicola* and *Phaeoisariopsis personata*). 75, 3–4.
- Dwivedi, S. L., Pande, S., Rao, J. N., & Nigam, S. N. (2002). Components of resistance to late leaf spot and rust among interspecific derivatives and their significance in a foliar disease resistance breeding in groundnut (*Arachis hypogaea* L.). *Euphytica*, 125(1), 81–88.
- Dwivedi, S.L., Crouch, J.H., Nigam, S.N., Ferguson, M.E., Paterson, A.H. (2003). Molecular breeding of groundnut for enhanced productivity and food security in the semi-arid tropics: opportunities and challenges. *Adv Agron* 80:153-222
- FAOSTAT (2023). FAO Statistical Database. Available online at: <http://faostat.fao.org/> (accessed Jan 9, 2024).
- Ibrahim, Y. J. (2010). Screening of groundnuts (*Arachis hypogaea* L.) for resistance to early and late leaf spot [MSc. thesis]. Kwame Nkrumah University of Science and Technology.
- Joshi E, Sasode DS, Singh N, Chouhan N. (2020). Diseases of groundnut and their control measures. *Biotica Res. Today*. 2(5): 232–237.

- Khan, A. R., Ijaz, M., Haq, I. U., Farzand, A., & Tariqjaved, M. (2014). Management of *Cercospora* leaf spot of groundnut (*Cercospora arachidicola* & use of systemic fungicides. *Cercetari Agronimice in Moldova*, XLVII(2), 98–102.
- Kishore GK, Pande S. (2005). Integrated management of late leaf spot and rust diseases of groundnut (*Arachis hypogaea* L.) with *Prosopis juliflora* leaf extract and chlorothalonil. *Int. J. Pest Manag.*; 51(4):325–332.
- Leal-Bertioli, S. C. M., José, A. C. V. F., Alves-Freitas, D. M. Moretzsohn, M. C., Guimarães, P. M., Nielen, S., Vidigal, B S., Pereira, R. W., Pike, J., Fávero, A. P., Parniske Varshney, R. K., & Bertioli, D. J. (2009). Identification o candidate genome regions controlling disease resistance in *Arachis*. *BMC Plant Biology*, 9(1), 112.
- Mace, E. S., Phong, D. T., Upadhyaya, H. D., Chandra, S., & Crouch, J. H. (2006). SSR analysis of cultivated groundnu (*Arachis hypogaea* L.) germplasm resistant to rust and late leaf spot diseases. *Euphytica*, 152(3), 317–330.
- Muhammad, A., & Bdliya, B. (2011). Effects of variety and fungicidal rate on *Cercospora* leaf spot disease of groundnut in the Sudan Savanna. *Nigerian Journal of Basic and Applied Sciences*, 19(1), 135–141.
- Pal KK, Dey R, Tilak KV. (2014). Fungal diseases of groundnut: Control and future challenges. *Future challenges in crop protection against fungal pathogens*. 1–29. DOI: 10.1007/978-1-4939-1188-2
- Shoba, D., Manivannan, N., Vindhivarman, P., & Nigam, N. (2012). SSR markers associated for late leaf spo disease resistance by bulked segregant analysis in groundnut (*Arachis hypogaea* L.). *Euphytica*, 188(2), 265 272.
- Subrahmanyam, P., McDonald, D., Waliyar, F., Reddy, L.J., Nigam, S.N., Gibbons, R.W., Rao, V.R., Singh, A.K., Pande, S., Reddy, P.M., Rao, P.S. (1995) Screening methods and sources of resistance to rust and late leaf spot of groundnut. *Information Bulletin no.47*.
- Wankhade, A.P., Chimote, V.P., Viswanatha, K.P., Yadaru, S., Deshmukh, D.B., Gattu, S., Sudini, H.K., Deshmukh, M.P., Shinde, V.S., Vemula, A.K., Pasupuleti, J. (2023). Genome-wide association mapping for LLS resistance in a MAGIC population of groundnut (*Arachis hypogaea* L.). *Theor. Appl. Genet.*; 136(3):43.
- Shew, B. B., Sommartya, T., Beute, M.K. (1989). Components of partial resistance in peanut genotypes to isolates of *Cercosporidium personatum* from the United States and Thailand. *Phytopathology*.79:136142.
- Waliyar, F., Shew, B. B., Sidahmed, R., Beute, M.K. (1995). Effects of host resistance on germination of *Cercospora arachidicola* on peanut leaf surfaces. *Peanut Science*.22:154-157.

Chiyembekeza, A.J., Knauff, D.A., Gorbet, D.W. (1993). Comparison of components of resistance in peanut to late leaf spot in different environments. *Crop Science*.33: 994-997.

Walls SB, Wynne JC, Beute MK. (1985). Resistance to late leaf spot of peanut of progenies selected for resistance to early leaf spot. *Peanut Science*.12:22-27.

UNDER PEER REVIEW