

## **Study of soil properties in major cropping systems in SPSR Nellore district, Andhra Pradesh, India.**

### **Abstract**

A study was carried out during 2023-24 in SPSR Nellore district of Andhra Pradesh, India to know the impact of major cropping systems on soil chemical and biological properties. Totally 100 soil samples were collected randomly from five major cropping systems *viz.*, paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses and groundnut-paddy (20 samples from each cropping systems) in SPSR Nellore district and processed, analyzed for available nitrogen, phosphorus, potassium and sulphur and dehydrogenase enzyme. The results revealed that available nitrogen in soils varied from 88 kg ha<sup>-1</sup> in paddy-cotton cropping system to 289 kg ha<sup>-1</sup> in paddy-pulses cropping system with a mean value of 130 and 242 kg ha<sup>-1</sup> respectively. 95% of samples showed deficient in available nitrogen. The available phosphorous in soils varied from 19 kg ha<sup>-1</sup> in paddy-cotton cropping system to 58 kg ha<sup>-1</sup> in paddy-pulses cropping system. The mean available phosphorus is 37.03 and 42.8 kg ha<sup>-1</sup> respectively. The available potassium in soils varied from 177 kg ha<sup>-1</sup> in paddy-paddy cropping system to 601 kg ha<sup>-1</sup> in paddy-pulses cropping system. The available potassium of soils of different cropping systems was medium to high in range. The available sulphur of soils varied from 7.56 mg kg<sup>-1</sup> in paddy-paddy cropping system to 33.73 mg kg<sup>-1</sup> in paddy-pulses cropping system. The soils had deficient to sufficient in available sulphur in various cropping systems at SPSR Nellore district of Andhra Pradesh. The soils in paddy-pulses cropping system showed the highest dehydrogenase activity (116.47 µg of TPF g<sup>-1</sup> soil day<sup>-1</sup>) whereas the lowest dehydrogenase activity (18.08 µg of TPF g<sup>-1</sup>day<sup>-1</sup>) was observed in paddy-paddy cropping system.

**Keywords:** Cropping systems, available nutrients and soil dehydrogenase activity and SPSR Nellore district.

### **INTRODUCTION**

Soil is a vital natural resource that plays a crucial role in the sustenance of life on Earth. It serves as the foundation for agricultural systems, supporting plant growth by providing essential nutrients, water and a medium for roots to anchor. Healthy soil directly

affects the quality and yield of crops. Cropping systems have a significant impact on soil health, influencing its chemical properties. Chemical properties refer to the composition and behaviour of various chemical elements and compounds within the soil. These properties play a crucial role in determining soil fertility, influencing plant growth, and supporting ecosystem functions. The chemical properties of soil are integral to its function as a medium for plant growth and a critical component of broader environmental systems. Proper management and monitoring of these chemical properties such as available nitrogen, phosphorus, potassium, sulphur help maintain soil fertility, prevent degradation, and support sustainable agricultural practices. Dehydrogenase activity in soil is a key indicator of microbial activity and overall soil health. Maintaining or enhancing dehydrogenase activity through practices that support soil microbial health is essential for promoting sustainable agricultural productivity and ecosystem resilience. The choice of crops, the diversity of plants, and the management practices all play crucial roles in determining the condition and sustainability of soil. Keeping this view, a study was taken up to know the effect of major cropping systems on soil chemical and biological properties in SPSR Nellore district, Andhra Pradesh, India.

## **MATERIALS AND METHODS**

The study area SPSR Nellore district lies in between  $14^{\circ} 4' 12.9''$  and  $14^{\circ} 57' 56.8''$  North latitudes and  $79^{\circ} 30' 29.6''$  and  $80^{\circ} 4' 24.7''$  East longitudes and the investigation was carried in the year 2023-24. One hundred soil samples were collected from five major cropping systems. From each cropping system 20 samples were collected randomly. The collected soil samples were shade dried, sieved with 2mm sieve and the sieved samples used for analysing chemical and biological properties. All the samples were analyzed for available nitrogen, phosphorus, potassium and sulphur and dehydrogenase enzyme. Available nitrogen in the soil samples was determined by alkaline potassium permanganate method (Subbiah and Asija, 1956) and expressed the results in  $\text{kg ha}^{-1}$ . Available phosphorus in the soil samples were extracted as per procedure described by Olsen *et al.* (1954) and phosphorus in the extract was determined by using ascorbic acid as reducing agent as described by Watanabe and Olsen (1965) using spectrophotometer (Systronics UV-VIS spectrophotometer 119) at 660 nm wavelength and expressed the results in  $\text{kg ha}^{-1}$ . Available potassium in the soil samples was extracted with neutral normal ammonium acetate (Jackson, 1973) and

determined by using flame photometer(Systronics flame photometer 128) and expressed the results in  $\text{kg ha}^{-1}$ . Available sulphur was determined by extracting with 0.15 per cent calcium chloride followed by development of turbidity with barium chloride(Hesse, 1971). The turbidity is measured using spectrophotometer (Systronics UV-VIS spectrophotometer 119) at 420 nm wavelength and expressed the results in  $\text{mg kg}^{-1}$ . Dehydrogenase activity in the soil sample was determined by the procedure described by Casida *et al.* (1964) and expressed in micro gram of triphenyl formazon (TPF) formed per gram of soil per day ( $\mu\text{g of TPF g}^{-1} \text{ soil day}^{-1}$ ).

## RESULTS AND DISCUSSION

### Available Nitrogen

The data in Table 1 showed that the available nitrogen in soils varied from 88  $\text{kg ha}^{-1}$  in paddy-cotton cropping system to 289  $\text{kg ha}^{-1}$  in paddy-pulses cropping system. The available nitrogen in soils of paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses, groundnut-paddy cropping systems was ranged from 100-163, 113-176, 88-213, 201-289 and 113-213  $\text{kg ha}^{-1}$ , respectively(Fig 1) with a mean value of 133, 140, 130, 242, and 159  $\text{kg ha}^{-1}$  respectively The mean available nitrogen in different cropping systems was in the order of paddy-cotton (130  $\text{kg ha}^{-1}$ ) followed by paddy-paddy (133  $\text{kg ha}^{-1}$ ), fallow-paddy (140  $\text{kg ha}^{-1}$ ), groundnut-paddy (159  $\text{kg ha}^{-1}$ ) and Paddy-Pulses (242  $\text{kg ha}^{-1}$ ).The CV was 30.29 per cent for available nitrogen in the soils of different cropping systems. Available nitrogen in soils of different cropping system in SPSR Nellore district was low to medium in range. The maximum value of available nitrogen (289  $\text{kg ha}^{-1}$ ) was observed in soils of paddy-pulses cropping system. The minimum value (88  $\text{kg ha}^{-1}$ ) of available nitrogen was observed in paddy-cotton cropping system.

The available nitrogen under paddy-pulses cropping system was higher than that of other cropping systems. Because of their symbiotic relationship with rhizobium bacteria, legumes have the capacity to fix and store more atmospheric nitrogen, increasing the amount of nitrogen that is readily available in the soil of paddy-pulses cropping systems (Rajpoot *et al.*, 2021).

### Available Phosphorus ( $\text{P}_2\text{O}_5$ )

The available phosphorus in soils varied from 19  $\text{kg ha}^{-1}$  in paddy-cotton cropping system to 57.5  $\text{kg ha}^{-1}$  in paddy-pulses cropping system (Table 1). The range

of available phosphorus varied from 20.2-30.1, 22.6-35.6, 19-30, 29.7-57.5 and 29.7-45.3 kg ha<sup>-1</sup> in soils of paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses and groundnut-paddy cropping systems respectively with a mean value of 25, 29.5, 24.5, 42.8 and 37.03 kg ha<sup>-1</sup> respectively (Fig 2). The mean available phosphorus in different cropping systems was in the order of paddy-cotton (24.5 kg ha<sup>-1</sup>) followed by paddy-paddy (25 kg ha<sup>-1</sup>), fallow-paddy (29.5 kg ha<sup>-1</sup>), groundnut-paddy (37.03 kg ha<sup>-1</sup>), paddy-pulses (42.8 kg ha<sup>-1</sup>) cropping system. The CV was 26.65 per cent for available phosphorus in the soils of different cropping systems. Available phosphorus in soils of different cropping system in SPSR Nellore district was low to medium in range. The maximum value of available phosphorus (57.5 kg ha<sup>-1</sup>) was observed in soils of paddy-pulses cropping system. The minimum value of available phosphorus (19 kg ha<sup>-1</sup>) was observed in paddy-cotton cropping system.

Compared to other cropping systems, the paddy-pulses cropping system has more accessible phosphorus. This might be due to the result of a higher available N in the paddy-pulses cropping system, where a positive interaction between available N and P is caused by BNF. Inclusion of legumes in crop rotation can promote mycorrhizal root colonization. Through root colonization, mycorrhizal connections have the biggest effect on raising P availability in soils. These outcomes agree with the research findings of Smith *et al.* (2011) and Newton *et al.* (2011).

### **Available Potassium (K<sub>2</sub>O)**

The data is presented in Table 1 indicated that the available potassium in soils varied from 177 kg ha<sup>-1</sup> in paddy-paddy cropping system to 601 kg ha<sup>-1</sup> in paddy-pulses cropping system. The available potassium in soils of paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses and groundnut-paddy cropping systems was ranged from 177-556, 244-401, 235-336, 249-601 and 215-367 kg ha<sup>-1</sup>, respectively with a mean value of 281, 322, 282, 418 and 308 kg ha<sup>-1</sup> respectively (Fig 3). The mean available potassium in different cropping systems were in the order of paddy-paddy (281 kg ha<sup>-1</sup>) followed by paddy-cotton (282 kg ha<sup>-1</sup>), groundnut-paddy (308 kg ha<sup>-1</sup>), fallow-paddy (322 kg ha<sup>-1</sup>), paddy-pulses (417 kg ha<sup>-1</sup>) cropping system. The CV was 24.80 per cent for available potassium of different cropping systems.

Available potassium in soils of different cropping system in SPSR Nellore district was medium to high in range. The maximum value of available potassium (601 kg ha<sup>-1</sup>) was observed in soils of paddy-pulses cropping system. The minimum value of

available potassium ( $177 \text{ kg ha}^{-1}$ ) was observed in paddy-paddy cropping system. The highest available potassium was observed in soils of paddy-pulses cropping system. This might be attributed to addition of large amount of K through crop residues. The potassium requirement for pulse crops is also less. Farmers may not applied the required quantity of K. This might be the reason for lower available N in paddy-paddy cropping system. The results are in agreement with the findings of Malecka *et al.* (2012) and Jat *et al.* (2018).

### **Available sulphur**

The available sulphur of soils varied from  $7.56 \text{ mg kg}^{-1}$  in paddy-paddy cropping system to  $33.73 \text{ mg kg}^{-1}$  in paddy-pulses cropping system. The range of available sulphur varied from 7.56-19.97, 9.68-19.04, 7.71-21.88, 11.74-33.73 and 7.86-25.2  $\text{mg kg}^{-1}$ , respectively in soils of paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses and groundnut-paddy cropping systems with a mean value of 12.69, 13.25, 13.41, 21.94 and  $17.31 \text{ mg kg}^{-1}$ , respectively. The mean available sulphur in different cropping systems were in the order of paddy-paddy ( $12.69 \text{ mg kg}^{-1}$ ) followed by fallow-paddy ( $13.25 \text{ mg kg}^{-1}$ ), paddy-cotton ( $13.41 \text{ mg kg}^{-1}$ ), groundnut-paddy ( $17.31 \text{ mg kg}^{-1}$ ), paddy-pulses ( $21.94 \text{ mg kg}^{-1}$ ). The CV was 36.45 per cent for available sulphur in the soils of different cropping systems. (Table 1 and Fig 4).

Available sulphur in soils of different cropping system in SPSR Nellore district was deficient to sufficient range. The maximum value of available sulphur ( $33.73 \text{ mg kg}^{-1}$ ) was observed in soils of paddy-pulses cropping system. The minimum value of available sulphur ( $7.56 \text{ mg kg}^{-1}$ ) was observed in paddy-paddy cropping system. The higher available sulphur in these soils might be due to the continuous application of fertilizers like single super phosphate which contains sulphur also. These results are in confirmation with findings of Kavitha *et al.* (2019).

### **Dehydrogenase Activity**

Data in Table 2 and Fig 5 showed that a wide variation was noticed from  $18.08 \mu\text{g of TPF g}^{-1}\text{day}^{-1}$  in paddy-paddy cropping system to  $116.47 \mu\text{g of TPF g}^{-1}\text{day}^{-1}$  in paddy-pulses cropping system. The dehydrogenase activity of soils in paddy-paddy, fallow-paddy, paddy-cotton, paddy-pulses and groundnut-paddy cropping systems was ranged from 18.08-43.47, 26.55-57.23, 20.20-45.59, 49.82-116.47 and 48.76-94.25

$\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ , respectively with a mean value of 31, 38.34, 31.99, 73.8 and 70.66  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ , respectively. The mean dehydrogenase activity in different cropping systems were in the order of paddy-paddy (31.25  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ) followed by paddy-cotton (31.99  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ), fallow-paddy (38.34  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ), groundnut-paddy (70.66  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ) and paddy-pulses cropping system (73.84  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ). The CV was 44.45 per cent for dehydrogenase activity in soils of different cropping systems.

The higher dehydrogenase activity (116.47  $\mu\text{g}$  of TPF  $\text{g}^{-1}\text{day}^{-1}$ ) was observed in soils of paddy-pulses cropping system. Higher dehydrogenase activity was discovered in paddy-legume cropping systems, that might be due to addition of more biomass to soil which is converted to organic matter and acted as feed or energy source to microbes led to more microbial proliferation. This indicates more microbial respiration and dehydrogenase activity related to increased microbial biomass and microbial activity, which promotes the degradation of organic materials which is strongly connected with dehydrogenase activity. Dehydrogenase activity is directly correlated with microbial population and also with moisture content in soil. (Zhang *et al.*, 2010). Similar findings were reported by Raviteja *et al.* (2024). Lesser dehydrogenase activity was observed in paddy-paddy cropping system.

#### References :

- Casida Jr, L.E., Klein, D.A and Santoro, T. 1964. Soil dehydrogenase activity. *Soil science*. 98 (6): 371-376.
- Jackson, M.L. 1973. *Soil Chemical Analysis*. Oxford IBH Publishing House, Bombay. 38.
- Jat, H.S., Datta, A., Sharma, P.C., Kumar, V., Yadav, A.K., Choudhary, M., Choudhary, V., Gathala, M.K., Sharma, D.K., Jat, M.L and Yaduvanshi, N.P.S. 2018. Assessing soil properties and nutrient availability under conservation agriculture practices in a reclaimed sodic soil in cereal-based systems of North-West India. *Archives of Agronomy and Soil Science*. 64(4): 531-545.
- Kavitha, M., Vajantha, B., Naidu, M.V.S and Reddi Ramu, Y. 2019. Spatial Distribution and Mapping of DTPA Extractable Micronutrients in Sugarcane Growing Tracts of Prudential Sugar Factory Zone in Chittoor District, Andhra Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*. 8(10): 968-976.
- Malecka, I., Blecharczyk, A., Sawinska, Z and Dobrzeniecki, T. 2012. The effect of various long-term tillage systems on soil properties and spring barley yield. *Turkish Journal*

*of Agriculture and Forestry*. 36: 217-226.

- Newton, Z.L., Kennedy, A.N and Chirwa, R.M. 2011. Grain legume impacts on soil biological processes in sub-Saharan Africa. *African Journal of Plant Science*. 5(1): 1-7.
- Olsen, S.R., Cole, C.V., Watanabe, F.S and Dean, L.A. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *Circular of United States Department of Agriculture*. 939.
- Rajpoot, R.S., Bajpai, R.K., Shrivastava, L.K., Kumar, U., Tedia, K and Mishra, V.N. 2021. Evaluation of soil physical and chemical properties under rice-based cropping system in Alfisols of Northern Hill Region of Chhattisgarh. *International Journal of Current Microbiology and Applied Sciences*. 10(1): 2748-2761.
- Raviteja,G., Vajantha,B., Krishnaveni, D., Prasanthi, A., Raveendra Reddy., M and Naidu., M.V.S.2024. Soil enzymatic activity and yield of sugarcane short crop as influenced by fertilizers and microbial inoculants. *International Journal of Advanced Biochemistry Research*. 8(5): 97-101.
- Smith, S.E., Jakobsen, I., Gronlund, M and Smith, F.A. 2011. Roles of arbuscular mycorrhiza as in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiology*. 156(3): 1050-1057.
- Subbiah, B.V and Asija, C.L. 1956. A rapid procedure for the estimation of available nitrogen in soils. *Current Science*. 25: 32.
- Watanabe, F.S and Olsen, S.R. 1965. Test of ascorbic acid method for determining phosphorous in water and sodium bicarbonate extracts of soils. *Soil Science Society of America Proceedings*. 29: 677-678.
- Zhang, W.J., Wang, X.J., Xu, M.G., Huang, S.M., Liu, H and Peng, C. 2010. Soil organic carbon dynamics under long term fertilizations in arable land of northern China. *Biogeosciences*. 7(2): 4.

**Table 1: Macronutrients status under major cropping systems of SPSR Nellore District, Andhra Pradesh**

Cropping Systems		Avail. N (kg ha <sup>-1</sup> )	Avail. P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	Avail K <sub>2</sub> O (kg ha <sup>-1</sup> )	Sulphur (mg kg <sup>-1</sup> )
Paddy-Paddy	Range	100-163	20.2-30.1	177-556	7.56-19.97
	Mean	133	25	281	12.69
Fallow-Paddy	Range	113-176	22.6-35.6	244-401	9.68-19.04
	Mean	140	29.5	322	13.25
Paddy-Cotton	Range	88-213	19-30	235-336	7.71-21.88
	Mean	130	24.5	282	13.41
Paddy-Pulses	Range	201-289	29.7-57.5	249-601	11.74-33.73
	Mean	242	42.8	418	21.94
Groundnut-Paddy	Range	113-213	29.7-45.3	215-367	7.86-25.2
	Mean	159	37.03	308	17.31
<b>Overall range</b>		88-289	19-57.5	177-601	7.56-33.73
<b>Overall mean</b>		161	32	322	16
<b>CV (%)</b>		30.29	26.65	24.80	36.45

**Table 2: Dehydrogenase activity under major cropping systems of SPSR Nellore District, Andhra Pradesh**

Cropping Systems		Dehydrogenase activity (µg of TPF g <sup>-1</sup> day <sup>-1</sup> )
Paddy-Paddy	Range	18.08- 43.47
	Mean	31
Fallow-Paddy	Range	26.55- 57.23
	Mean	38.34
Paddy-Cotton	Range	20.20- 45.59
	Mean	31.99
Paddy-Pulses	Range	49.82- 116.47
	Mean	73.8
Groundnut-Paddy	Range	48.76- 94.25
	Mean	70.66
<b>Overall range</b>		18.08-116.47
<b>Overall mean</b>		49
<b>CV (%)</b>		44.45




Fig 1: Range of available nitrogen under major cropping systems







Fig 2: Range of available phosphorus under major cropping systems



**Fig 3: Range of available potassium under major cropping systems**



**Fig 4: Range of available Sulphur under major cropping systems**



**Fig 5: Range of dehydrogenase activity under major cropping systems**

UNDER PEER REVIEW