

ANTIMICROBIAL ACTIVITY OF *Xylopi* *aethi* *opica* EXTRACT ON MICROORGANISMS ASSOCIATED WITH THE POILAGE OF VEGETABLES IN MILE 3 MARKET, PORT HARCOURT NIGERIA

ABSTRACT

With increasing demand for natural preservatives, this study explores the effectiveness of *Xylopi* *aethi* *opica* extracts in reducing vegetable spoilage. The aim was to evaluate these extracts' potential to inhibit spoilage microorganisms, determine the most effective concentration for preservation, and assess their impact on sensory qualities. Vegetables (onions, tomatoes, and carrots) were purchased from vendors at Mile 3 Market, Port Harcourt, and transported to a microbiological laboratory for analysis. The vegetable samples were treated with ethanol and aqueous extracts of *Xylopi* *aethi* *opica* at concentrations of 10%, 20%, and 30% (w/v). For microbial analysis, treated and control vegetable samples were homogenized, cultured on nutrient agar for bacteria and potato dextrose agar for fungi, and incubated at optimal conditions for colony formation. Enumeration of bacterial and fungal colonies was conducted periodically over 14 days. Microorganisms isolated included *Bacillus cereus*, *Pseudomonas aeruginosa*, *Aspergillus niger*, and *Penicillium* spp. Phytochemical analysis revealed that the ethanol extract had higher concentrations of bioactive compounds, including alkaloids (2.5%), flavonoids (3.1%), saponins (1.4%), tannins (2.2%), phenolics (4.3%), and glycosides (1.1%), compared to the aqueous extract. Results showed that the 30% ethanol extract was the most effective, significantly reducing bacterial and fungal growth compared to the control and other treatments. ANOVA analysis confirmed significant reductions in microbial counts in treated vegetables ($p < 0.05$). Sensory evaluation revealed that the extracts, particularly the 30% ethanol extract, did not negatively affect the vegetables' taste, smell, or appearance. This study demonstrates the potential of *Xylopi* *aethi* *opica* as an effective natural preservative, offering a promising alternative to synthetic preservatives. Future research should focus on long-term effects, broader applications, and consumer acceptance for practical implementation.

Keywords: *Xylopi* *aethi* *opica*, Vegetables, Phytochemicals, ANOVA

1.0 Introduction

Vegetable spoilage is a significant issue in food security, particularly in developing countries where preservation methods are often limited. Microbial contamination is one of the leading causes of vegetable spoilage, contributing to substantial post-harvest losses. In Mile 3 Market, a major trading hub in Port Harcourt, Nigeria, the problem of vegetable spoilage is exacerbated by poor hygiene practices, inadequate storage facilities, and the humid tropical climate, which creates a conducive environment for the growth of spoilage microorganisms. Understanding and

controlling these microorganisms is essential to reduce spoilage and extend the shelf life of vegetables, which are a staple in the diet of many Nigerians (Karanth et al., 2023).

Xylopiya aethiopyca, commonly known as "African pepper" or "Negro pepper," is a plant native to West Africa and widely used in traditional medicine and cuisine. The dried fruits of *Xylopiya aethiopyca* have been reported to possess various pharmacological properties, including antimicrobial, antioxidant, and anti-inflammatory activities. The plant's antimicrobial properties have garnered significant attention, particularly in traditional medicine, where it is used to treat various infections and diseases (Karanth et al., 2023).

Xylopiya aethiopyca, belongs to the *Annonaceae* family and It is a tropical plant known for its aromatic seeds and fruit, which are frequently used in both cooking and traditional medicine (Hutchinson & Dalziel, 1954). The seeds are found inside a woody, aromatic fruit that splits open when ripe. This fruit and its seeds are celebrated for their antimicrobial properties, making them valuable in culinary and medicinal contexts.

The plant's composition includes essential oils like xylopic acid, alkaloids, flavonoids, and phenolic compounds. These compounds contribute to its antimicrobial, antioxidant, and anti-inflammatory effects. *Xylopiya aethiopyca* is particularly noted for its ability to combat a wide range of microorganisms, which makes it a promising natural preservative (Bagwai et al., 2020).

Vegetables, rich in fiber, vitamins, and antioxidants, are essential for human health. However, they are prone to microbial spoilage, which affects their quality and shelf life. Common preservation methods such as refrigeration, canning, and drying help extend the (Petcu et al., 2023) shelf life of vegetables, but they may alter the taste and texture of the produce (Mungofa et al., 2022).

The antimicrobial properties of *Xylopiya aethiopyca* are particularly relevant in vegetable preservation. Its essential oils exhibit broad-spectrum antimicrobial activity, making it an effective alternative to synthetic preservatives (Fleischer et al., 2008). When used to protect vegetables from microbial contamination, *Xylopiya aethiopyca* can help extend their shelf life without compromising their quality, though careful consideration of its impact on taste and aroma is necessary for consumer acceptance.

The antimicrobial activity of *Xylopiya aethiopyca* is attributed to its rich phytochemical composition. Studies have identified a range of bioactive compounds in the plant, including alkaloids, tannins, flavonoids, and essential oils, which have demonstrated efficacy against a wide spectrum of microorganisms (Fleischer et al., 2008). These properties make *Xylopiya aethiopyca* a promising candidate for developing natural preservatives, especially in regions like Mile 3 Market, where synthetic preservatives' high cost and limited availability pose a challenge.

Microorganisms associated with vegetable spoilage, such as *Escherichia coli*, *Salmonella* spp., *Staphylococcus aureus*, and various molds and yeasts, are of particular concern in the Mile 3 Market. These microorganisms not only cause rapid deterioration of vegetables but also pose significant health risks to consumers (Degaga et al., 2022). The conventional approach to managing vegetable spoilage in such settings involves improving storage conditions and using chemical preservatives. However, chemical preservatives often raise concerns regarding safety, environmental impact, and the development of resistant strains of microorganisms (Degaga et al., 2022).

Given the challenges associated with synthetic preservatives, there is a growing interest in exploring natural alternatives that are both effective and safe. *Xylopiya aethiopyca* offers a potential solution, as its antimicrobial properties could be harnessed to inhibit the growth of spoilage microorganisms and thereby extend the shelf life of vegetables. Research in this area is crucial to validate the efficacy of *Xylopiya aethiopyca* against specific spoilage microorganisms and to determine the practical applications of this plant in food preservation (Degaga et al., 2022).

The challenges posed by microbial spoilage of vegetables are not only a concern for food safety but also the economic well-being of market traders and consumers. In Mile 3 Market, vegetables are sold in large quantities daily, with much of the produce coming from surrounding rural areas. However, due to the high ambient temperatures and humidity, vegetables are highly susceptible to microbial invasion, leading to spoilage before they are sold or consumed. The microorganisms responsible for this spoilage are diverse, including bacteria, yeasts, and molds, each of which can cause different forms of deterioration, such as soft rot, discoloration, and off-odors.

The economic losses incurred from vegetable spoilage are significant, especially in markets like Mile 3, where many traders rely on daily sales for their livelihood (Balali et al., 2020). The rapid spoilage also leads to price fluctuations and food shortages, further exacerbating food insecurity in the region. Moreover, the consumption of spoiled vegetables can lead to foodborne illnesses, which are a major public health concern in Nigeria. The need for effective, affordable, and locally available solutions to prevent vegetable spoilage is therefore critical.

Xylopiya aethiopyca has long been used in traditional African medicine to treat a variety of ailments, including infections caused by bacteria and fungi. Its use as a spice in cooking also hints at its potential role as a natural preservative, as many culinary spices have been historically used to extend the shelf life of food. The phytochemicals present in *Xylopiya aethiopyca*, such as diterpenes, xylopic acid, and sesquiterpenes, have shown significant antimicrobial activity in various studies. These compounds work by disrupting the cell membranes of microorganisms, inhibiting their growth, and ultimately leading to cell death. Given its antimicrobial properties, *Xylopiya aethiopyca* could serve as an effective natural preservative for vegetables, particularly in traditional markets where refrigeration is limited or nonexistent. Studies have shown that extracts from *Xylopiya aethiopyca* can inhibit the growth of a variety of foodborne pathogens, including

Escherichia coli, *Staphylococcus aureus*, and *Salmonella* spp., all of which are commonly associated with food spoilage and foodborne diseases (Juliani et al., 2008).

Findings suggest that incorporating *Xylopi*a *aethi*o*pica* into preservation strategies could help reduce spoilage rates, thereby improving the quality and safety of vegetables sold in Mile 3 Market. The use of natural preservatives like *Xylopi*a *aethi*o*pica* aligns with the global trend towards reducing the reliance on synthetic chemicals in food production. Consumers are increasingly concerned about the potential health risks associated with synthetic preservatives, which have been linked to various adverse effects, including allergic reactions and the development of antibiotic-resistant bacteria. By contrast, natural preservatives are generally perceived as safer and more environmentally friendly, making them more acceptable to consumers. However, while the antimicrobial properties of *Xylopi*a *aethi*o*pica* are well-documented, its practical application in preserving vegetables remains underexplored. There is a need for detailed studies that not only confirm the efficacy of *Xylopi*a *aethi*o*pica* against spoilage microorganisms in real-world conditions but also determine the optimal methods of application, such as concentration, form (e.g., powder, extract, or essential oil), and method of incorporation into storage practices. It is also important to evaluate the potential sensory impacts of using *Xylopi*a *aethi*o*pica* as a preservative (Bagwai et al., 2020). While its strong aromatic properties may be desirable in certain contexts, they could alter the taste or smell of vegetables, potentially affecting consumer acceptance. Therefore, research into the organoleptic properties of *Xylopi*a *aethi*o*pica*-treated vegetables is essential to ensure that the benefits of reduced spoilage do not come at the cost of consumer satisfaction. Studies have shown that *Xylopi*a spp., a plant species native to West Africa, possesses potent antimicrobial properties (Ibe et al., 2023). This presents an opportunity to investigate its potential use as a natural preservative for vegetables commonly sold at local markets like Mile 3 Market. Given the limited research on *Xylopi*a extract's efficacy against vegetable spoilage microorganisms in a real-world market setting, this study seeks to address this gap and explore its practical application within the context of local agricultural practices.

2.0 Materials and Method

2.1 Sample Collection

The study was conducted at Mile 3 Market in Port Harcourt, Rivers State, Nigeria, focusing on fresh vegetables like onions, tomatoes, and carrots, known for their susceptibility to spoilage. Vegetables were collected from different vendors, selected based on freshness and uniformity, and transported to the laboratory in sterile bags. They were washed with sterile water and air-dried.

*Xylopi*a *aethi*o*pica* fruits were sourced from a local herbal market, authenticated, cleaned, and air-dried. The dried fruits were ground into powder and extracted with ethanol and water. The ethanol extract was obtained by soaking the powdered fruit in 95% ethanol for 48 hours, filtered,

and concentrated. The aqueous extract was similarly prepared using distilled water and concentrated.

2.2 Culturing of Sample

Vegetables were divided into groups and treated with different concentrations (10%, 20%, and 30%) of ethanol and aqueous extracts. A control group was treated with sterile water. After immersion, the vegetables were air-dried. Microbial analysis was performed by swabbing the vegetable surfaces at different time intervals (days 0, 3, 7, and 14), followed by inoculation onto nutrient agar plates. The plates were incubated, and colony-forming units (CFUs) were counted.

2.3 Identification of Organisms

Staining techniques, including Gram and lactophenol cotton blue stains, were used to identify bacterial and fungal isolates. Biochemical tests (e.g., catalase, oxidase, indole, citrate, and urease) were conducted to further characterize the bacteria.

2.4 Sensory Evaluatio

the sensory evaluation was conducted using a trained panel to assess the taste, smell, and appearance of vegetables treated with *Xylopiya aethiopica* extracts. Panelists were selected and trained to recognize subtle differences in sensory attributes, following guidelines to ensure consistency in scoring.

Panelists evaluated treated and control samples independently using a 5-point hedonic scale (1 = very poor, 5 = excellent) to rate each attribute. Taste was assessed by noting any off-flavors or undesirable changes introduced by the extracts, while smell was rated to detect any unusual or strong odors that could impact the vegetables' acceptability. Appearance was examined based on color, texture, and overall visual appeal.

Evaluations took place under controlled conditions to avoid environmental influences, with panelists provided water and plain crackers to cleanse their palates between samples. Scores were averaged for each attribute and treatment to determine the extracts' impact on sensory qualities.

2.5 Phytochemical Analysis

To determine the active compounds, standard qualitative phytochemical screening methods were employed. These methods involve specific chemical tests to detect the presence and estimate the concentration of various bioactive compounds, including alkaloids, flavonoids, saponins, tannins, phenolics, and glycosides. For each compound, reagents specific to its chemical nature are added, causing color changes or precipitates that indicate its presence and relative abundance.

Quantitative estimations were then made based on the intensity of these reactions, comparing the ethanol and aqueous extracts.

This qualitative and semi-quantitative approach confirmed that the ethanol extract had higher concentrations of these bioactive compounds, highlighting its greater potential as a natural preservative.

2.6 Statistical Analysis

Data were analyzed using descriptive statistics and ANOVA, with significance set at $p < 0.05$. The study provided a comprehensive evaluation of *Xylopiya aethiopica* as a natural preservative for vegetables.

3.0 RESULT

Table 1 shows that the total bacterial count on the vegetables increased over time in all groups. However, the treated vegetables (both ethanol and aqueous extracts) exhibited significantly lower bacterial counts compared to the control group. The 30% ethanol extract was the most effective, resulting in the lowest bacterial counts at all-time points. Statistical analysis (ANOVA) as seen in Table 1 confirmed that there was a significant reduction in bacterial growth in the treated groups compared to the control ($p < 0.05$).

As seen in Table 2, the fungal count increased significantly in the control group, while the treated groups showed much lower fungal growth. The 30% ethanol extract again proved to be the most effective, with the lowest fungal counts at all-time points. ANOVA analysis showed a significant reduction in fungal growth in treated vegetables compared to the control ($p < 0.05$) as shown in Table 2).

Table 3 showed *Bacillus cereus* was the most frequently isolated microorganism, with a high occurrence in tomatoes at 40%, followed by carrots at 35%, and onions at 30%. The overall occurrence across all vegetables was 35%. *P. aeruginosa* had the highest occurrence in tomatoes at 50% and was also commonly found in carrots and onions at 30% and 20% respectively, leading to an overall occurrence of 33.3%. *Aspergillus niger* was more commonly isolated from onions with an occurrence of 25%, and had lower occurrences in carrots and tomatoes at 20% and 15%, respectively. The overall occurrence was 20%.

Penicillium spp. was evenly distributed across the vegetables, with the highest occurrence in carrots at 25%, followed by tomatoes at 20%, and onions at 15%. This resulted in an overall occurrence of 20%. *Escherichia coli* was most frequently found in tomatoes at 25%, with similar occurrences in carrots at 20% and onions at 10%, leading to an overall occurrence of 18.3%. *Saccharomyces cerevisiae* had the lowest overall occurrence, being most present in carrots at 15%, followed by tomatoes at 10% and onions at 5%, with an overall occurrence of 10%.

The sensory evaluation (Table 4) indicated that the *Xylopi* *aethi* *opica* extracts did not negatively affect the taste, smell, or appearance of the treated vegetables. Higher concentrations of the extracts, particularly the 30% ethanol extract, received higher sensory scores compared to the control. ANOVA analysis showed significant differences in sensory scores between treatments and the control ($p < 0.05$) as shown in Table 4).

The phytochemical (Table 5) analysis revealed that the ethanol extract of *Xylopi* *aethi* *opica* had higher concentrations of alkaloids, flavonoids, saponins, tannins, and phenolics compared to the aqueous extract. These bioactive compounds are known for their antimicrobial properties, which likely contributed to the effectiveness of the extracts in inhibiting spoilage microorganisms.

Table 1: Total Bacterial Count (CFU/mL) on Treated and Control Vegetables

Treatment	Day 0	Day 3	Day 7	Day 14
Control (Sterile Water)	0	1.5×10^4	4.5×10^5	9.2×10^5
10% Ethanol Extract	0	1.0×10^3	2.3×10^4	4.0×10^4
20% Ethanol Extract	0	5.0×10^2	8.0×10^3	2.2×10^4
30% Ethanol Extract	0	1.0×10^2	3.0×10^3	1.0×10^4
10% Aqueous Extract	0	2.0×10^3	3.0×10^4	6.5×10^4
20% Aqueous Extract	0	1.5×10^3	1.5×10^4	3.2×10^4
30% Aqueous Extract	0	5.0×10^2	5.0×10^3	1.2×10^4

Table 2: Total Fungal Count (CFU/mL) on Treated and Control Vegetables

Treatment	Day 0	Day 3	Day 7	Day 14
Control (Sterile Water)	0	1.0×10^3	2.5×10^4	5.8×10^4
10% Ethanol Extract	0	5.0×10^2	1.2×10^3	2.4×10^3
20% Ethanol Extract	0	2.5×10^2	8.0×10^2	1.5×10^3
30% Ethanol Extract	0	1.0×10^2	3.5×10^2	8.0×10^2
10% Aqueous Extract	0	7.5×10^2	1.5×10^3	3.0×10^3
20% Aqueous Extract	0	5.0×10^2	1.0×10^3	2.0×10^3
30% Aqueous Extract	0	2.0×10^2	5.5×10^2	1.2×10^3

Table 3: Percentage Occurrence of Isolated Microorganisms in Vegetables

Microorganism	Onions (%)	Tomatoes (%)	Carrots (%)	Overall (%)
<i>Bacillus cereus</i>	30	40	35	35
<i>Pseudomonas aeruginosa</i>	20	50	30	33.3
<i>Aspergillus niger</i>	25	15	20	20
<i>Penicillium</i> spp.	15	20	25	20
<i>Escherichia coli</i>	10	25	20	18.3
<i>Saccharomyces cerevisiae</i>	5	10	15	10

Table 4: Sensory Evaluation Scores (Mean ± SD)

Treatment	Taste	Smell	Appearance	(%)
Control (Sterile Water)	3.5 ± 0.5	3.0 ± 0.4	2.8 ± 0.6	-
10% Ethanol Extract	4.2 ± 0.4	4.0 ± 0.3	4.0 ± 0.4	10
20% Ethanol Extract	4.5 ± 0.3	4.4 ± 0.3	4.5 ± 0.3	20
30% Ethanol Extract	4.7 ± 0.2	4.6 ± 0.2	4.8 ± 0.2	30
10% Aqueous Extract	4.0 ± 0.4	3.8 ± 0.4	3.9 ± 0.4	10
20% Aqueous Extract	4.3 ± 0.3	4.1 ± 0.3	4.2 ± 0.3	20

Table 5: Phytochemical Composition of *Xylopi aethiopia* Extracts

Phytochemical	Ethanol Extract (%)	Aqueous Extract (%)
Alkaloids	2.5	1.8
Flavonoids	3.1	2.5
Saponins	1.4	0.9
Tannins	2.2	1.6
Phenolics	4.3	3.7
Glycosides	1.1	0.8

4.0 Discussion

The findings from this study underscore the significant effectiveness of *Xylopiya aethiopicia* extracts, particularly the 30% ethanol extract, in inhibiting the growth of spoilage microorganisms on vegetables. These results align well with previous research, which highlighted the potent antimicrobial properties of *Xylopiya aethiopicia* due to its rich phytochemical composition (Japhet et al., 2024).

The study demonstrated a noticeable reduction in both bacterial and fungal counts in the vegetables treated with *Xylopiya aethiopicia* extracts compared to the untreated control group. Over time, the bacterial and fungal counts in the control group increased significantly, which is consistent with the natural progression of spoilage in fresh produce. This spoilage is often exacerbated by the presence of spoilage microorganisms such as *B. cereus* and *P. aeruginosa* for bacteria, and *A. niger* and *Penicillium spp.* for fungi, as evidenced by the data presented in the tables. The increase in microbial counts in untreated vegetables is a common occurrence documented in numerous studies, reflecting the high susceptibility of fresh vegetables to microbial contamination, which can lead to rapid spoilage and potential health risks (Sokamte et al., 2018).

The significant reduction in microbial counts observed in vegetables treated with *Xylopiya aethiopicia* extracts can be attributed to the presence of bioactive compounds such as alkaloids, flavonoids, saponins, tannins, and phenolics. These compounds were found in higher concentrations in the ethanol extract, as highlighted in the phytochemical analysis. Such bioactive compounds are well-known for their antimicrobial properties, which include disrupting microbial cell membranes and inhibiting the growth and proliferation of pathogens. This aligns with previous research findings where ethanol extracts of various plants exhibited stronger antimicrobial activity compared to aqueous extracts, likely due to the ability of ethanol to extract a higher concentration of these potent phytochemicals (Dzoyem et al., 2017).

The high occurrence of *B. cereus* in tomatoes and carrots, with frequencies of 40% and 35% respectively, highlights the vulnerability of these vegetables to this particular bacterium. *B. cereus* is notorious for causing foodborne illnesses through the production of toxins, and its high prevalence in commonly consumed vegetables raises significant food safety concerns. Similarly, the presence of *P. aeruginosa* in high proportions, particularly in tomatoes, is concerning due to this bacterium's association with spoilage. *P. aeruginosa* is known for producing a biofilm, which not only contributes to spoilage but also makes the bacterium more resistant to conventional preservation methods (Yu et al., 2020). The detection of *A. niger* and *Penicillium spp.* primarily in onions and carrots is equally noteworthy, as these fungi are responsible not only for spoilage but also for the production of mycotoxins. Mycotoxins pose serious health risks,

including carcinogenic effects, making the inhibition of these fungi in food products critical for public health.

The sensory evaluation results suggest that the use of *Xylopiya aethiopica* extracts, particularly at higher concentrations such as the 30% ethanol extract, does not negatively affect the taste, smell, or appearance of the treated vegetables. This finding is significant as it addresses one of the primary challenges in using natural preservatives—maintaining the sensory qualities of food products (Japhet et al., 2024). Notably, the higher sensory scores for vegetables treated with the 30% ethanol extract indicate that not only is the extract effective in microbial inhibition, but it may also enhance the sensory appeal of the vegetables. This enhancement is likely due to the prevention of off-flavors and odors that are typically associated with microbial spoilage.

The practical implications of these findings are profound, particularly for the use of *Xylopiya aethiopica* extracts in food preservation. The study's results suggest that these extracts can be effectively applied in market settings where fresh vegetables are sold, which are often prone to rapid spoilage. Implementing methods such as direct immersion in the extract, which was shown to be effective in this study, could be a simple yet highly effective strategy to extend the shelf life of fresh produce. This not only reduces food waste but also helps to mitigate economic losses for vendors, who often struggle with the rapid spoilage of their goods (Etebu & Enaregha, 2013).

From a health perspective, the ability of *Xylopiya aethiopica* extracts to inhibit spoilage microorganisms has significant implications. By reducing the levels of spoilage microorganisms like *B. cereus* and *E. coli*, which were frequently isolated in this study, the risk of foodborne illnesses associated with these pathogens is substantially lowered. This is particularly important given the widespread consumption of fresh vegetables and the potential health risks posed by contaminated produce (Abarikwu et al., 2016). Furthermore, the inhibition of fungi like *A. niger* reduces the risk of mycotoxin exposure, which is a serious public health concern due to the carcinogenic potential of these compounds. The findings of this study support the potential of *Xylopiya aethiopica* extracts as a natural and effective means of preserving fresh vegetables. The extracts not only inhibit spoilage microorganisms but also maintain or even enhance the sensory qualities of the vegetables. This aligns with previous research on the antimicrobial efficacy of plant extracts and highlights the importance of phytochemicals in developing food preservation strategies that are both effective and safe for consumers.

The results of this study provide compelling evidence that *Xylopiya aethiopica* extracts, particularly the 30% ethanol extract, are highly effective in reducing microbial growth and preserving the quality of fresh vegetables. This conclusion is drawn from the significant reduction in both bacterial and fungal counts observed in treated vegetables compared to the control group, with the 30% ethanol extract consistently showing the greatest efficacy. The phytochemical analysis further supports these findings, revealing that the ethanol extract of *Xylopiya aethiopica* contains higher concentrations of bioactive compounds known for their

antimicrobial properties, such as alkaloids, flavonoids, saponins, tannins, and phenolics (Abarikwu et al., 2016).

These findings align with previous research that has demonstrated the potent antimicrobial activity of *Xylopiya aethiopica* due to its rich phytochemical profile (Ejelonu et al., 2011; Ogbonna et al., 2013). The ability of these extracts to significantly inhibit the growth of common spoilage microorganisms like *Bacillus cereus*, *Pseudomonas aeruginosa*, *Aspergillus niger*, and *Penicillium spp.* highlights their potential as natural preservatives in the food industry. This is particularly important given the increasing demand for natural and safe alternatives to synthetic preservatives, which are often associated with health risks.

5.0 Conclusion

The practical application of *Xylopiya aethiopica* extracts in preserving fresh vegetables not only extends the shelf life of these products but also helps to reduce food waste and economic losses associated with spoilage. The sensory evaluation results indicate that these extracts do not adversely affect the taste, smell, or appearance of the vegetables, and in some cases, they may even enhance these qualities. This is a crucial consideration for consumer acceptance, as maintaining the sensory attributes of food products is essential for marketability. The significant reduction in spoilage microorganisms achieved through the use of *Xylopiya aethiopica* extracts has important public health implications. By effectively controlling pathogens like *Bacillus cereus* and *Escherichia coli*, these extracts help to mitigate the risk of foodborne illnesses, which are a major concern associated with the consumption of contaminated fresh produce. Additionally, the inhibition of mycotoxin-producing fungi like *A. niger* reduces the potential for mycotoxin exposure, thereby enhancing food safety.

This study underscores the potential of *Xylopiya aethiopica* extracts as a viable natural alternative for preserving fresh vegetables. The extracts not only inhibit microbial growth but also preserve or enhance the sensory qualities of the vegetables, making them suitable for practical application in food preservation. Given the growing consumer preference for natural preservatives, the use of *Xylopiya aethiopica* extracts could play a significant role in meeting this demand while also contributing to food safety and sustainability. Future research should focus on optimizing the extraction and application methods to maximize the efficacy of these extracts, as well as exploring their potential use in preserving other food products.

REFERENCES

Abarikwu, S. O., Ogunlaja, A., Otuechere, C. A., & Gideon, O. (2016). Effect of Ethanolic Extract from Seeds or Pods of *Xylopiya Aethiopica* (Dunal) A. Rich (Annonaceae) on the Testicular Function of Adult Male Rats. *Indian Journal of Clinical Biochemistry*, 32(4), 420–428. <https://doi.org/10.1007/s12291-016-0622-5>

- Bagwai, M., Magashi, A., & Bukar, A. (2020). Preservative activity of *Xylopi*a *aethi*o*pica* fruits bio-active fractions on fresh meat. *Bayero Journal of Pure and Applied Sciences*, 12(1), 308–314. <https://doi.org/10.4314/bajopas.v12i1.47s>
- Balali, G. I., Yar, D. D., Dela, V. G. A., & Adjei-Kusi, P. (2020). Microbial contamination, an increasing threat to the consumption of fresh fruits and vegetables in today’s world. *International Journal of Microbiology*, 2020, 1–13. <https://doi.org/10.1155/2020/3029295>
- Dzoyem, J., McGaw, L., Kuete, V., & Bakowsky, U. (2017). Anti-inflammatory and anti-nociceptive activities of African medicinal spices and vegetables. In *Elsevier eBooks* (pp. 239–270). <https://doi.org/10.1016/b978-0-12-809286-6.00009-1>
- Fleischer, T. C., Mensah, M. L. K., Mensah, A. Y., Komlaga, G., Gbedema, S. Y., & Skaltsa, H. (2008). Antimicrobial Activity Of Essential Oils Of *Xylopi*a *aethi*o*pica* *African Journal of Traditional Complementary and Alternative Medicines*, 5(4), 391. <https://doi.org/10.4314/ajtcam.v5i4.31295>
- Japhet, L. B., Oluwatunase, G. O., & Adejayan, T. O. (2024). Ethanolic Extract of *Xylopi*a *aethi*o*pica* Attenuated Aluminum-Induced Ovarian Toxicity in Adult Female Wistar Rats. *JBRA*, 28(2). <https://doi.org/10.5935/1518-0557.20240024>
- Ibe, C., Irondi, E. A., Ugbogu, E. A., & Anyamele, T. (2023). A review of the nutritional and antimicrobial properties of *Xylopi*a *aethi*o*pica* A.Rich. *Food and Humanity*, 1, 1099–1116. <https://doi.org/10.1016/j.foohum.2023.09.007>
- Juliani, H. R., Kwon, T., Koroch, A. R., Asante-Dartey, J., Acquaye, D., & Simon, J. E. (2008). *Xylopi*a *aethi*o*pica* (Annonaceae): Chemistry, Traditional Uses and Functional Properties of an “African Pepper.” In *ACS symposium series* (pp. 114–128). <https://doi.org/10.1021/bk-2008-0987.ch007>
- Karant, S., Feng, S., Patra, D., & Pradhan, A. K. (2023). Linking microbial contamination to food spoilage and food waste: the role of smart packaging, spoilage risk assessments, and date labeling. *Frontiers in Microbiology*, 14. <https://doi.org/10.3389/fmicb.2023.1198124>
- Karioti, A., Hadjipavlou-Litina, D., Mensah, M. L. K., Fleischer, T. C., & Skaltsa, H. (2004). Composition and Antioxidant Activity of the Essential Oils of *Xylopi*a *aethi*o*pica* (Dun) A. Rich. (Annonaceae) Leaves, Stem Bark, Root Bark, and Fresh and Dried Fruits, Growing in Ghana. *Journal of Agricultural and Food Chemistry*, 52(26), 8094–8098. <https://doi.org/10.1021/jf040150j>

Mungofa, N., Sibanyoni, J. J., Mashau, M. E., & Beswa, D. (2022). Prospective role of indigenous leafy vegetables as functional food ingredients. *Molecules*, 27(22), 7995. <https://doi.org/10.3390/molecules27227995>

Ogbonna, C. N., Nozaki, N., K, N., Yajima, N., & H, N. (2013). Antimicrobial activity of *Xylopi*a *aethi*o*pica*, *Aframomum melegueta* and *Piper guineense* ethanolic extracts and the potential of using *Xylopi*a *aethi*o*pica* to preserve fresh orange juice. *African Journal of Biotechnology*, 12(16), 1993–1998. <https://doi.org/10.5897/ajb2012.2938>

Petcu, C. D., Tăpăloagă, D., Mihai, O. D., Gheorghe-Irimia, R., Negoită, C., Georgescu, I. M., Tăpăloagă, P. R., Borda, C., & Ghimpețeanu, O. M. (2023). Harnessing natural antioxidants for enhancing food shelf life: Exploring sources and applications in the food industry. *Foods*, 12(17), 3176. <https://doi.org/10.3390/foods12173176>

Sokamte, T., Mbougung, P., Tatsadjieu, N., & Sachindra, N. (2018). Phenolic compounds characterization and antioxidant activities of selected spices from Cameroon. *South African Journal of Botany*, 121, 7–15. <https://doi.org/10.1016/j.sajb.2018.10.016>

Yu, S., Yu, P., Wang, J., Li, C., Guo, H., Liu, C., Kong, L., Yu, L., Wu, S., Lei, T., Chen, M., Zeng, H., Pang, R., Zhang, Y., Wei, X., Zhang, J., Wu, Q., & Ding, Y. (2020). A Study on Prevalence and Characterization of *Bacillus cereus* in Ready-to-Eat Foods in China. *Frontiers in Microbiology*, 10. <https://doi.org/10.3389/fmicb.2019.03043>