

Evaluation of Physico-chemical Characteristics of Sponge Cake with Liquid Egg Yolk

ABSTRACT

Aims: This research aims to determine the quality of sponge cake with using liquid eggs based on expandability, texture, protein content, fat content, moisture content and color and produce products with high nutritional value.

Study design: This research are an laboratory experimental with a completely randomized design.

Place and Duration of Study: This research are located in Laboratory of Animal Product Technology, Faculty of Animal Science, Universitas Brawijaya. This research was conducted from July - August 2024.

Methodology: This research are using a 5 treatments and 5 replication. The main focus are a different substitutions. The 5 treatments are SPC1 (without using liquid egg yolk), SPC2 (25%), SPC3 (50%), SPC4 (75%), SPC5 (100%) of the total egg content.

Results: The use of different liquid eggs in making sponge cake gave a very significant effect ($P < 0.01$) on the expandability, texture, protein content, fat content, moisture content and color values. Each variable increased as the use of liquid eggs added increased.

Conclusion: The addition of liquid egg in the preparation of sponge cake can improve the expandability, texture, protein content, fat content, moisture content. The addition of liquid egg also affects the color of sponge cake to be more intense yellow. The use of 100% liquid egg in making sponge cake can improve physicochemical quality.

Keywords: Fresh egg; liquid egg; sponge cake; texture.

1. INTRODUCTION

Liquid egg yolk is an egg product that has been separated from the shell and made of egg yolk, and has gone through processes such as homogenization and pasteurization which are useful for ensuring food safety (Ho, Ciou, Qiu, Hsieh, Shih, Chen, et al., 2021). There are several advantages of liquid eggs based on the food industry, namely ease of use, reducing the time and effort required to prepare eggs such as breaking and beating eggs. Liquid egg also has the advantage of having a stable consistency that allows it to produce a more consistent end result in bakery products, with the advantages possessed by this liquid egg which makes liquid egg an alternative to fresh eggs used in food products such as sponge and angel cake (Yang, Yang, Dou and Chi, 2024).

Sponge cake is one of the foods made from simple ingredients such as eggs, flour and sugar. Sponge cake is known as a cake that has a soft texture like a sponge and tastes sweet (Yan, Liu, Cao, Wang, Li, Ren, et al., 2023). There is an advantage that sponge cake has, which is not using baking powder as a developer. Instead, sponge cake is made from eggs beaten together with sugar, until it expands perfectly. Eggs are the main ingredient used in the process of making sponge cake, and eggs are very influential in determining the sensory characteristics of sponge cake. There is previous research on making angel cake with the use of pasteurized egg white which results in maximum cake volume increase (Singh, Geveke, Jones and Tilman, 2019). The making of angel cake with the use of fermented eggs produces a more attractive texture and aroma compared to the use of fresh eggs (Jia, Xiong, Bai, Yuan, Song, Lan, et al., 2022). Therefore, the use of liquid egg in making sponge cake is useful to maximize the texture produced by sponge cake. The use of liquid egg in the preparation of sponge cake can provide a soft texture and high enough

chewiness due to the presence of albumin in eggs which functions as a binding agent to form a smooth and soft texture. In addition, liquid egg is also useful for increasing the volume of sponge cake due to its effective emulsification properties, which maximizes the entry of air into the dough (Shidara, Arimitsu, Miyashita, Uematsu, Akiniwa, Nishijima, et al., 2023).

2. METHODOLOGY

2.1 Materials

This research focused on sponge cakes made with liquid egg yolk. The ingredients used to make sponge cake are fresh chicken eggs obtained from Batu chicken farm, liquid egg yolk, flour, sugar, and butter obtained from Supermarket. The tools used in the research are mixer, container, baking pan and oven.

2.2 Methods

The research was conducted using a laboratory experimental method with a completely randomized design (CRD), involving 5 treatments and 5 repetitions. The five treatments are SPC1 (without using liquid egg), SPC2 (25%), SPC3 (50%), SPC4 (75%), SPC5 (100%) of the total egg content.

2.3 Data Analysis

The experimental data collected was analyzed using Microsoft Excel. A series of statistical analyses such as determining the mean, standard deviation, and performing an analysis of variance test were performed on the numerical results. ANOVA indicated the presence of significant differences among the treatments. Therefore, Duncan's Multiple Range Test (DMRT) was conducted for pairwise comparisons to strictly distinguish means that were significantly different from each other.

2.4 Production of Sponge Cake using Liquid Egg

Put the fresh egg or liquid egg along with sugar into the mixer and mixer on high speed for 3 minutes until the egg is fluffy and white like foam, then add the flour slowly on low speed and make sure the dough is homogeneous. Then melt the butter and put it into the cake batter by stirring slowly until smooth. Put the dough on a baking sheet lined with parchment paper and bake in the oven at 145°C for 45 minutes. Sponge cake using liquid egg is ready for further analysis.

2.5 Quality of Sponge Cake

2.5.1 Expandability

Expandability is one of the quality indicators of sponge cake to show the ability of the dough to expand optimally during the baking process. The procedure for measuring the height of the dough before and after baking is then determined using the formula:

$$\text{Expandability} = \frac{B - A}{A} \times 100\%$$

A : Height of sample before oven

B : Height of sample after oven

2.5.2 Texture

Texture is a parameter that includes softness, elasticity, and density of the pore structure. Texture test measurements use a texture analyzer with units of N.

2.5.3 Protein Content

Protein content can affect the structure and quality of sponge cakes. Proteins in flour such as glutenin and gliadin are important in the elasticity and the ability of the dough to resist

pressure during the dough's expansion. The procedure for testing protein content is by measuring the weight of the sample and then putting it in a kjeldahl flask, then adding distilled water and titrating it and then calculating it with the formula:

$$\text{Protein} = \frac{(VA - VB) \text{ Hcl} \times N \text{ Hcl} \times 14,007}{W \times 1000} \times 100\%$$

Notes:

VA: mL HCl for sample titration

VB: mL HCl for blank titration

N: normality of standard HCl used

14.007: atomic weight of nitrogen

W: sample weight in grams

2.5.4 Fat Content

Fat content acts as a tenderizer, which is to soften the texture of the cake by inhibiting the formation of excessive gluten, resulting in a more elastic texture. The procedure for measuring fat content is by putting the sample into a woolen cloth and placing it into a soxhlet and heating it to 105°C and cooling it in a desiccator for 30 minutes and then calculated by the formula:

$$\text{Fat Content} = \frac{\text{Fat Weight}}{\text{Sample Weight}} \times 100\%$$

2.5.5 Moisture Content

Moisture content was measured using the gravimetric method. The petri dish was first weighed without the sample (A). Then, 2 g of sausage sample was placed on the petri dish, which had been preheated in a 105°C oven for 12 hours, and weighed again (B). Dry the sample for 12 hours, then placed it on desiccator for 10 minutes before weighed it (C). The final weight can be calculated using the content formulation =

$$\text{Moisture Content} = \frac{(B - C)}{(B - A)} \times 100\%$$

2.5.6 Color Values

Color is determined using a color reader. Before analyzing the sample with a color reader, the color reader must be calibrated using black and white plates. Cut the sample about 2 cm x 2 cm x 2 cm and place it on the tray. Place the color reader lens over the sample and click the power button. The color value can be seen on the screen with 3 types of colors: redness, yellowness, and brightness.

3. RESULTS AND DISCUSSION

3.1 Expandability

The addition of liquid egg yolk in making sponge cake has a highly significant effect ($P < 0,01$) on the expandability. This can occur because liquid eggs have a fairly high moisture content, which can increase the moisture of the dough. In addition, liquid eggs are easy to beat into a stable foam, so they can maintain the dough structure during the baking process (Ho, Ciou, Qiu, Hsieh, Shih, Chen, et al., 2021). The protein in eggs also coagulate when exposed to heat, thus helping to form a firm yet light cake structure. This is comparable to previous research which states that the maturity level of a good sponge cake is in the range of 6.91-7.47% (Setyaningsih, Suraya and Salsabila, 2021).

Table 1. Average Expandability of Sponge Cake

Treatments	Expandability (%) ± SD
------------	------------------------

SPC1	5.95 ± 0.58 ^a
SPC2	6.95 ± 0.29 ^b
SPC3	7.05 ± 0.31 ^b
SPC4	7.35 ± 0.13 ^{bc}
SPC5	8.00 ± 0.26 ^c

*a, b, c, d shows a highly significant effect

3.2 Texture

The addition of liquid egg in making sponge cake gives no significant effect on texture produced. This can occur because the addition of liquid eggs to the sponge has the ability to bind water and more optimal emulsification power and also the presence of moisture content in liquid eggs helps increase the viscosity of the dough, which affects the stability of the froth to the resulting texture (Inanlar and Altay, 2024). In addition, the use of liquid egg yolk, which is high in fat, serves as an emulsifier to improve the texture and softness of the sponge cake (Arbowati, Mulyani and Hintono, 2021). Previous research stated that the texture of sponge cake ranging from 5.9 - 10.5 N is a good texture on sponge cake (Renzetti and Sman, 2022).

Table 2. Average Texture of Sponge Cake

Treatments	Texture (N) ± SD
SPC1	4.05 ± 0.53 ^a
SPC2	4.14 ± 0.28 ^a
SPC3	4.33 ± 0.60 ^a
SPC4	5.23 ± 0.22 ^a
SPC5	5.83 ± 0.18 ^a

3.3 Protein Content

The addition of liquid egg yolk in making sponge cake has a highly significant effect ($P < 0,01$) on the protein content of the sponge cake. This can occur because liquid egg is one of the ingredients that has a high enough protein content and functions to form the dough with the interaction between water and fat in the dough. In addition, the protein in liquid eggs undergoes denaturation and clumping when heated to form a hard and voluminous sponge cake structure (Wang, Yang, Li, Jin, Shu, Jin, et al., 2024). This is comparable to previous research which states that the protein content in sponge cake with the addition of moringa flour has increased, which ranges from 8.56-10.80% (Anurani and Evanuarini, 2022).

Table 3. Average Protein Content of Sponge Cake

Treatments	Protein Content (%) ± SD
SPC1	9.25 ± 0.25 ^a
SPC2	10.29 ± 0.18 ^b
SPC3	12.87 ± 0.40 ^c
SPC4	14.36 ± 0.27 ^d
SPC5	15.96 ± 0.26 ^e

*a, b, c, d, e shows a highly significant effect

3.4 Fat Content

The use of liquid egg in making sponge cake gives a highly significant effect ($P < 0,01$). This can occur because liquid egg contains egg yolk which is rich in fat and phospholipids, especially lecithin, which functions as an emulsifier (Zhao, Li, Liu and Chen 2023). The fat not only increases the softness of the cake, but also helps stabilize the air bubbles formed during whisking, maintaining the volume of the batter and resulting in a smoother texture. The emulsification process supported by lecithin also allows the fat to be well integrated, giving the sponge cake a moist and fluffy effect.

Table 4. Average Fat Content of Sponge Cake

Treatments	Fat Content (%) ± SD
SPC1	9.38 ± 0.56 ^a

SPC2	10.23 ± 0.30 ^{ab}
SPC3	10.96 ± 0.45 ^{bc}
SPC4	11.80 ± 0.58 ^{cd}
SPC5	12.43 ± 0.53 ^d

*a, b, c, d shows a highly significant effect

3.5 Moisture Content

The use of liquid egg in making sponge cake gives a highly significant effect ($P < 0,01$). This can occur because liquid egg that has gone through the pasteurization process has a higher moisture content, but with a high enough moisture content in liquid egg that can function as an emulsifier, resulting in a dough that is resistant to air and the texture of the sponge cake becomes light and soft. In making low fat sponge cake with the addition of pumpkin has a moisture content of 27.49-31.75% (Kurniawan and Puspita, 2019).

Table 5. Average Moisture Content of Sponge Cake

Treatments	Moisture Content (%) ± SD
SPC1	32.30 ± 0.68 ^a
SPC2	33.23 ± 0.57 ^{ab}
SPC3	33.37 ± 0.65 ^{ab}
SPC4	33.91 ± 0.38 ^b
SPC5	34.47 ± 0.74 ^b

*a, b shows a highly significant effect

3.6 Color Values

The use of liquid egg in making sponge cake gives a highly significant effect ($P < 0,01$). This can occur because egg yolk contains natural pigments such as carotenoids (lutein and zeaxanthin) which give a yellowish color. The use of liquid egg yolk can increase the pigment content which contributes to the increase of L (brightness), a* (bluish-reddish), and b* values in sponge cake. In addition, liquid egg yolk can influence the Maillard reaction, which plays a role in color formation during baking. This reaction will run more evenly, resulting in a more intense cake color. This is in accordance with previous research stating that a* color has a value ranging from 7-15 (Slade, Kweon and Levine, 2021), while b* color has a value ranging from 66.57 -74.00 (Nguyen and Tran, 2021).

Table 6. Average color values of sponge cake

Treatments	Color L ± SD	Color a ± SD	Color b* ± SD
SPC1	79.09 ± 0.89 ^a	2.75 ± 0.27 ^a	59.36 ± 0.82 ^a
SPC2	81.23 ± 0.73 ^{ab}	4.04 ± 0.26 ^{ab}	62.56 ± 0.76 ^{ab}
SPC3	84.55 ± 0.97 ^{bc}	5.51 ± 0.70 ^{bc}	65.09 ± 0.59 ^c
SPC4	87.46 ± 0.95 ^c	7.32 ± 0.85 ^c	69.25 ± 0.68 ^c
SPC5	89.92 ± 0.77 ^d	10.30 ± 0.89 ^d	70.50 ± 0.94 ^d

*a, b, c, d shows a highly significant effect

4. CONCLUSION

The addition of liquid egg in making sponge cake can improve the texture. The use of 100% liquid egg in sponge cake gives very significant results in expandability, texture, protein content, fat content, moisture content and color values. Further research needs to be done on the safety of sponge cake with using of liquid eggs.

ACKNOWLEDGEMENTS

This research was supported by the Faculty of Animal Science at Universitas Brawijaya.

Authors' Contributions

This research was conducted with contributions from all authors, who reviewed and approved the final version of the manuscript.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Ho, H. Y., J. Y. Ciou, Y. T. Qiu, S. L. Hsieh, M. K. Shih, M. H. Chen, and C. Y. Hou. 2021. Improvement of foaming characteristics and stability of sterilized liquid egg with egg white hydrolysate (EWH). *Foods*, 10(6),1326. <https://doi.org/10.3390/foods10061326>
2. Yang, S., F. Yang, W. Dou and Y. Chi. 2024. Testing adulterated Liquid-egg: Developing rapid detection techniques based on Colorimetry, Electrochemistry, and interfacial fingerprinting. *Food Chemistry*, 138674. <https://doi.org/10.1016/j.foodchem.2024.138674>
3. Yan, Z., J. Liu, S. Cao, Z. Wang, C. Li, J. Ren, and X. Liu. 2023. Substitution of sucrose by erythritol in sponge cake: Effect on protein foaming, baking performance and digestion properties. *International Journal of Biological Macromolecules*, 253, 126759. <https://doi.org/10.1016/j.ijbiomac.2023.126759>
4. Singh, A., D. J. Geveke, D. R. Jones, and E. D. Tilman. 2019. Can acceptable quality angel food cakes be made using pasteurized shell eggs? The effects of mixing factors on functional properties of angel food cakes. *Food Science & Nutrition*, 7(3), 987-996. <https://doi.org/10.1002/fsn3.911>
5. Jia, J., D. Xiong, J. Bai, Y. Yuan, Q. Song, T. Lan, and X. Duan. 2022. Investigation on flavor and physicochemical properties of angel food cakes prepared by lactic acid fermented egg white. *LWT*, 164, 113659. <https://doi.org/10.1016/j.lwt.2022.113659>
6. Shidara, H., K. Arimitsu, T. Miyashita, G. Uematsu, S. Akiniwa, N. Nishijima and T. Kojima. 2023. Application of Eggs in Food Product Development. In *Handbook of Egg Science and Technology* (pp. 237-258). CRC Press.
7. Ho, H. Y., J. Y. Ciou, Y. T. Qiu, S. L. Hsieh, M. K. Shih, M. H. Chen, and C. Y. Hou. 2021. Improvement of foaming characteristics and stability of sterilized liquid egg with egg white hydrolysate (EWH). *Foods*, 10(6), 1326. <https://doi.org/10.3390/foods10061326>

8. Setyaningsih, D., J. Suraya, and S. Salsabila. 2021. Effect of mono-acylglycerol (mag) addition as emulsifier for bakery products. *Journal of Agricultural Industry Technology*, 31(2), 198-210. <https://doi.org/10.24961/j.tek.ind.pert.2021.31.2.198>
9. Inanlar, B., and F. Altay. 2024. Effects of Egg on Cake Batter Rheology and Sponge Cake Texture. *Journal of Food Process Engineering*, 47(11), e14765. <https://doi.org/10.1111/jfpe.14765>
10. Arbowati, J. L., Mulyani, S., & Hintono, A. (2021). Effect of egg quality on physical properties and organoleptic properties of sponge cake. *Journal of Food Technology and Nutrition*, 20(1), 27-34. <http://dx.doi.org/10.33508/jtpg.v20i1.2680>
11. Renzetti, S., and R. G. van der Sman. 2022. Food texture design in sugar reduced cakes: Predicting batters rheology and physical properties of cakes from physicochemical principles. *Food Hydrocolloids*, 131, 107795. <https://doi.org/10.1016/j.foodhyd.2022.107795>
12. Wang, Q., Y. Yang, Z. Li, H. Jin, D. Shu, Y. Jin, and L. Sheng. 2024. Research advances on the effects of thermal and non-thermal processing techniques on the physicochemical properties and microbiological control of liquid eggs. *Food Control*, 155, 110106. <https://doi.org/10.1016/j.foodcont.2023.110106>
13. Anurani, R., and H. Evanuarini. 2022. Quality of Sponge Cake with Moringa Leaf Flour Addition in terms of Yield, Fat Content and Protein Content. *Thesis*.
14. Zhao, F., R. Li, Y. Liu, and H. Chen. 2023. Perspectives on lecithin from egg yolk: Extraction, physicochemical properties, modification, and applications. *Frontiers in Nutrition*, 9, 1082671. <https://doi.org/10.3389/fnut.2022.1082671>
15. Kurniawan, T., and J. Puspita. 2019. Effects Of Ripe Pumpkin (*Cucurbita Moschata*) Addition As Fat Mimetics Based Carbohydrate In Low Fat Sponge Cake (Doctoral dissertation, Unika Soegijapranata Semarang).
16. Slade, L., M. Kweon, and H. Levine. 2021. Exploration of the functionality of sugars in cake-baking, and effects on cake quality. *Critical Reviews in Food Science and Nutrition*, 61(2), 283-311. <https://doi.org/10.1080/10408398.2020.1729694>
17. Nguyen, T. M. N., and G. B. Tran. 2021. Application of chickpeas aquafaba with pre-treatment as egg replacer in cake production. *Chemical Engineering Transactions*, 89, 7-12. <https://doi.org/10.3303/CET2189002>