

Study the protective role of *Berberis vulgaris* aqueous extract on some Liver enzymes and proteins in male rats treated with Monosodium glutamate

Abstract--- Twenty adult male rats, ages 2-4 months, with an average weight of 150-300 g, have been utilized in the study. The rats have been split up to four groups (G), with five rabbits in each one of the groups. G1 was treated with normal saline extract 15ml orally as the positive control group, G2 has been treated with aqueous extract 14mg/kg as the negative control group, G3 has been treated with aqueous extract of *Berberis vulgaris* (300mg/kg BW), and G4 has been treated with aqueous extract of *Berberis vulgaris* (300mg/kg BW) following 3hrs. This study sought to ascertain protective effect of *Berberis vulgaris* extract on albumin, total protein, aspartate amino transaminase (ALT), globulin, alkaline phosphate (ALP), and alanine amino transaminase (AST). The current study's findings indicated that the mean levels of ALT, AST, ALP, albumin, total protein, and globulin have been much higher ($p \leq 0.050$) in G2 group than in G1 group. We have found no significant difference ($p \geq 0.050$) in levels of ALT, AST, total protein, ALP, albumin, and globulin in the group that had received doses of *Berberis vulgaris* (300mg/kg BW) and Monosodium glutamate (14mg/kg BW) in comparison with G1. We found a significant decrease ($p \leq 0.050$) in levels of (ALP, ALT, and AST) total protein, globulin, and albumin in G4 and G3 in comparison with G2.

Keywords: Monosodium glutamate toxic, *Berberis vulgaris* extract, ALT, AST, ALP and proteins, rats.

Introduction

Therapeutic plants are presently the subject of intense academic interest. Saponins, phytic acids, phytates, proanthocyanidins, cardioprotective flavonoids, tannins, polysaccharides, lipids, and other phyto-components were found in various plant parts. Because of its widespread prevalence, barberry is considered a significant therapy for health difficulties associated with infectious infections. The fruit of the common herb barberry is utilized as food additive in many cultures as a medicinal plant. This shrub, which grows to a height of 1m to 3m, has barbed yellow wood and oval leaves with pendulous yellow flowers, which eventually turn into long, scarlet fruits that are 3 to 5 mm wide and 7 to 10 mm long. Those fruits are edible when they ripen in late summer and early fall. They include high levels of vitamin C and have a tart flavor (4,5). North America, Southern and central Europe, and Southern Asia are among the places in which this plant is found (6,7). This plant's many parts, like its stems, roots, leaves, and fruits, have been utilized in the traditional medicine in Iran as well as other countries (8). This plant has been utilized in the area of traditional medicine for several conditions, like the gastrointestinal problems, kidney stones, gallbladder and liver problems, circulatory system stimulation, and urinary tract disorders (9, 10). *Berberis vulgaris* is used for the treatment of several conditions because of its anti-bacterial, anti-hypertensive, anti-cancer, anti-oxidant, anti-inflammatory, and hepatoprotective properties (11). Triterpenoids, vitamin K, ascorbic acid, at least 10 phenolic compounds, and over 30 alkaloids are found in

Comment [CO1]: The abstract lacks an introduction to what the study is about and also a concluding sentence on the overall impact of the study and/or future directions.

Comment [CO2]: G2 was treated...

Comment [CO3]: G3 was treated...

Comment [CO4]: was

Comment [CO5]: Put this way, this sounds like the outcome of the study. I'd suggest "have been identified" Also, do you mean to say, "have been identified in various parts of different plants"? Because what you have here may be misunderstood as various parts of a particular plant and your plant of interest has not been introduced prior to this sentence.

Comment [CO6]: Prior to this sentence, there should be an introduction of the Barberry plant as one of such plants with therapeutic effects.

Comment [CO7]: Why does your first citation begin from 4 and not 1 ?

Comment [CO8]: For easy of understanding, the first time the Barberry plant is mentioned, you should include the botanical name before subsequent usage

Berberis vulgaris. In Berberis vulgaris, berberine is the main alkaloid that is extracted (12). According to research, it has anti-triglyceridemic, anti-oxidant, anti-bacterial, anti-cancer, anti-parasitic, anti-diabetic, and anti-hepatotoxic properties. Plant roots, stems, and leaves are used as powerful tonic and for therapeutic uses, which include sickness treatment, in the Middle East, Europe, and the US (13). In addition to that, Berberis vulgaris root extract is efficient at treating and preventing calculi development in the renal system (14). Monosodium glutamate (MSG), a non-essential amino acid in human body, has been widely used as amino additive since 1909 as a result of its characteristic umami flavor. Monosodium glutamate is frequently found in foods like tomatoes, cheese, fermented soy products, and yeast extracts. It is mostly absorbed in the small intestine when it enters the human body (15). The following binding of TAS1R1-TAS1R3 promotes intracellular calcium release, which in turn activates the umami flavor of monosodium glutamate, a taste area in the brain. Monosodium glutamate may have harmful effects on people, which include metabolic syndrome, cancer, immunological dysfunction, neurotoxicity, cardiovascular disease, fetal underdevelopment, infertility, and renal toxicity, according to mounting experimental and clinical data. Additionally, studies suggest that nutraceuticals and medicinal herbs could mitigate or reverse the negative effects of monosodium glutamate (16). The purpose of this research is to ascertain how the aqueous extract of Berberis vulgaris affects certain proteins and enzymes that are induced by Monosodium glutamate in male rats.

Comment [CO9]: MSG is not a non-essential amino acid but a sodium salt of Glutamic acid which is a non-essential amino acid

Comment [CO10]: could

Methods and Materials

Materials

Preparation of aqueous extract of *Berberis vulgaris*

From the Baghdad Governorate in Iraq, a Berberis vulgaris plant was obtained. After being washed and chopped into little pieces, the plants have been processed into a fine powder with the use of an electric grinder. A total of 30g of dry Berberis vulgaris powder were taken via a sensitive scale, Sartorius type, and put inside a 1000ml glass flask, which contains 500ml of distilled water. The solution was after that covered and left at room temperature for 24hrs before being filtered by Whatman filter paper No. 101. The filtrate was after that taken and the sediment was left, and the filtrate was then put in clean, sterilized metal dishes and put inside an electric oven at 40 degrees for 24. Put the dry extract (17) in glass bottles and store them in refrigerator until the time that they are needed.

Comment [CO11]: A Berberis vulgaris plant was obtained from the Baghdad Governorate in Iraq

Comment [CO12]: were

Comment [CO13]: fine

Comment [CO14]: contained

Comment [CO15]: then

Comment [CO16]: collected

Comment [CO17]: discarded (I suppose)

Comment [CO18]: for 24 hours

Comment [CO19]: The dry extract (17) was placed in glass bottles and stored in the refrigerator until use.

Comment [CO20]: I don't see the relevance of the exact dates, you can say the study was performed for a time period of one month

Comment [CO21]: were

Experimental animals

Beginning on December 27, 2023, and ending on January 27, 2024, this study was carried out. There were twenty adult male Albino rats in this study, ages 2-4 months, with average weights of 150-300 g. The animals have been raised at a 25°C temperature, with water and adequate ventilation under controlled conditions. Before the experiment began, the animals were given two weeks to acclimate. Twenty rats have been utilized in the present study; they have been split up to 4 groups, with five male rats in every group being treated differently.

1. Group I, the negative control group, received 15 milliliters of normal saline orally as a control.

2. Monosodium glutamate (MSG) at a level of (14mg/kg B.W.) has been administered orally to Group 2's Positive Control group for 30 days.

Comment [CO22]: was

3. Berberis vulgaris extract was given orally to Group 3 rats for 30 days at a 300mg/kg B.W. dose.

4. Group 4 received MSG (14mg/kg B.W.) for 30 days after receiving (300mg/kg BW) of Berberis vulgaris extract orally.

Comment [CO23]: Do you mean this group received both MSG and the extract? If so, rephrase accordingly

Methods

Serum liver enzymes analysis

All groups' blood serum was centrifuged at 3000rpm for 15min. in order to extract the serum, which was after that stored in a refrigerator at a low degree of temperature (-20°C) for the purpose of performing the required physiological tests, including measuring ALT and AST levels using method (18) and ALP using method (19).

Comment [CO24]: was collected and centrifuged

Comment [CO25]: then

Total protein, globulin and albumin estimation.

The blood serum's total protein content was determined with the use of technique (20), albumin levels were determined using method (21) and globulin levels were determined using approach (22).

Statistical Analysis: Every total calculated with a one-way ANOVA at the significance level was regarded as a significant value ($p < 0.05$) (23).

Results and Discussion

By analyzing the liver enzyme levels in the animals' serum during the course of our 30-day experiment, we discovered that MSG significantly raises ($p \leq 0.050$) levels of ALP, ALT, and AST in comparison to G1. We could not detect any significant difference ($p \geq 0.050$) between the group that had received doses of MSG (300mg/kg BW) and Berberis vulgaris (14mg/kg BW). Additionally, as shown in table (1), the Berberis vulgaris (300 mg/kg B.W) G3 significantly lowers ($p \leq 0.050$) levels of ALP, ALT, and AST compared with G1. After a month of oral MSG administration at a dose of 15mg/kg, the liver enzyme levels of male rats increased. This outcome aligned with the research of (24,25). When male rats were given MSG with drinking water at a 4g/L concentration per day for six weeks, Onobrudu and Barine (26) found that levels of liver enzymes increased. (27) showed that the same outcome was obtained when rats were given MSG daily for four weeks at a dosage of 2g/kg. Additionally, during six weeks, rats were given daily oral MSG at a dosage of 0.04–0.08 mg/kg in the study. The increased levels of liver enzymes ALP, AST, and ALT in the blood serum of rats given MSG were indicative of the substance's hepatotoxicity by all of the aforementioned researchers, despite the obvious variation in MSG concentrations and the varying administration durations. The substance's effect on producing damage, liver, and loss of hepatocytes might be the cause of the raised liver enzyme levels in the blood serum of the MSG group; as hepatocyte destruction increases, so does the secretion of these enzymes into the bloodstream (29). Additionally, this substance's generation of oxidative stress and creation of free

Comment [CO26]: Include units of measurements

Comment [CO27]: Put the author before this

radicals will interact with unsaturated fatty acids in the liver cell membrane, causing the mitochondrial membranes to be destroyed and allowing such enzymes to leak (30,31).

Levels of aminotransferases ALT, AST and alkaline phosphatase ALP are an indicator of liver function through the presence or absence of liver cell dysfunction (32) and also result from the breakdown of L-glutamate (MSG) which turns into harmful glutamine inside liver cells, leading to their destruction and the release of their enzymes (33).

The elevation of MSG results in an increase in glutamate dehydrogenase (GDH) due to the release of amine groups as ammonia and the subsequent synthesis of ammonium ion (NH₄⁺), which is toxic and adversely impacts liver function. This effect results in liver damage and destruction, therefore causing the release of liver enzymes (34, 35).

We noticed from Table (1) that the aqueous extract of *Berberis vulgaris* works to reduce liver enzymes and prevents the damage resulting from the MSG. The results of the present study have been found consistent with the study of (24,36, 37,38).

The cause may be a result of the protective function of barberry on the liver cell membrane, resulting in decreased liver enzyme levels. Consequently, this plant functions to preserve the physiological activity of the liver. Chemical study of the plant revealed the presence of active compounds including berberine, berbamine, oxyacanthine and palmatine. These compounds are therapeutically significant and utilized in disease treatment due to their antioxidant, antibacterial, anticancer, and anti-inflammatory characteristics (36).

The aqueous extract also inhibits the production of reactive oxygen radicals due to its strong antioxidant activity. A study indicated that giving rats aqueous extract of barberry at a concentration of (25-50-100) for 11 months improved liver function by reducing enzyme activity. This indicates that the plant has many medicinal properties, which makes the plant have a protective role in protecting liver tissue (39). It works to prevent damage to liver, kidney, heart and brain tissue and their safety (40).

Table (1) Effects of aqueous extract of *Berberis vulgaris* (300mg/kg B.W) on the level of liver enzymes in serum of the male rats that have been treated with MSG for 30 days

	ALP I/U	AST I/U	ALT I/U
Control negative group	40.38 ± 0.75 a	28.40 ± 0.85 a	20.62 ± 0.46 a
Control positive group Monosodium glutamate (14mg/kg BW)	79.18 ± 3.81 b	68.66 ± 1.73 b	37.66 ± 1.33 b
<i>Berberis vulgaris</i> (300mg/kg BW) third group	31.60 ± 0.98 C	20.24 ± 0.69 c	17.10 ± 0.51 c
<i>Berberis vulgaris</i> (300mg/kg B.W) And Monosodium glutamate (14mg/kg B.W.)	38.20 ± 1.32 A	27.40 ± 0.82 a	21.08 ± 0.34 a
LSD	5.18	2.62	1.86

Comment [CO28]: What do the units mean?

P value Significant ≤ 0.05

Results in Table 2 reveal a significant increase ($p \leq 0.050$) in globulin, albumin, and total protein levels in male rats administered MSG at a concentration of 14mg/kg body weight daily for 30 days, in comparison to negative controls. However, no significant differences ($p \geq 0.050$) were observed in globulin, albumin, and total protein levels between groups G3 and G4 and group G1.

Additionally, Table (2) highlights the positive effect of the aqueous extract of *Berberis vulgaris* (300mg/kg body weight) on protein levels. These findings align with the studies of (47, 48, 49, 50). The experiment had also shown a significant increase ($p \leq 0.050$) in serum globulin, albumin, and total protein levels in male rats treated with MSG, consistent with studies (42, 43, 44).

The increase in protein levels is attributed to MSG's toxic effect on liver tissue, leading to its degradation, which results in protein breakdown and release into the bloodstream. This also elevates albumin and globulin levels (30). One study indicated that prolonged daily MSG intake causes liver necrosis and cell breakdown or kidney damage, impairing the proximal convoluted tubules' ability to reabsorb protein due to reduced Na^+/K^+ -ATPase activity (43, 45). Another study reported that orally dosing rabbits with MSG at 0.25–0.5–1 g/kg body weight for 14, 28, or 56 days led to an increased total protein rate (46).

Elevated levels of the amino acids may result in stimulating the synthesis of the albumin in the liver, with some hormones like the thyroxine, growth hormone, and insulin promoting the increase in the production of albumin. The higher levels of total protein and globulin that have been observed in the present study result from the MSG's impact on the metabolic dysfunction, reduced liver efficiency in the storage of vitamins and minerals, in addition to increased generation of the free radicals because of the hepatic damage and inflammation (44).

The oral administration of *Berberis vulgaris*' alcoholic extract results in reducing these 3 levels of proteins, which suggests that the plant has protective roles in the health of the liver through supporting the metabolic processes as well as improving the levels of total protein, albumin, and globulin. This effect results from the plant's capability in decreasing the reactive oxygen species (ROS) that are caused by the oxidative stress, due to its constituents like the potassium, magnesium, flavonoids, vitamins, and unsaturated fatty acids. These components are helpful in safeguarding against diseases at the same time as enhancing liver and kidney functions.

These findings are consistent with studies (47, 48). A study by (49) also found that oral administration of *Berberis vulgaris* aqueous extract for 2–4 weeks, either alone or alongside the toxic substance aflatoxin, preserved protein levels and decreased liver enzymes, urea, creatinine, and blood sugar levels. This supports the plant's role in preventing liver damage. Our results align with these findings. Prior research has shown a significant reduction in total protein and albumin levels following administration of an aqueous extract of barberry leaves at concentration levels of 40–80–120mg/kg daily for 6 weeks in rats treated with the carbon tetra-chloride (50).

Table (2) Effects of aqueous extract of *Berberis vulgaris* (300mg/kg B.W) on levels of Globulin, Albumin and Total protein in serum of male rats with MSG for 30 days

	Globulin mg/dl	Albumin mg/dl	Total protein mg/dl
Control negative group	2.52 ± 0.09 a	4.16 ± 0.82 a	5.52 ± 0.18 a
Control positive group Monosodium glutamate (14 mg/kg BW)	3.76 ± 0.20 b	7.30 ± 0.37 b	8.56 ± 0.40 b
<i>Berberis vulgaris</i> (300mg/kg B.W) third group	2.22 ± 0.07 ac	3.70 ± 0.11 a	4.84 ± 0.19 c
<i>Berberis vulgaris</i> (300mg/kg B.W) And Monosodium glutamate (14 mg/kg BW)	2.56 ± 0.14 a	3.88 ± 0.27 a	5.32 ± 0.16 a
LSD	0.380	0.70	0.63

P value Significant ≤ 0.05

Berberis vulgaris has significant potential to enhance protective functions by increasing liver enzyme and protein levels. These effects are attributed to the direct action of the plant extract on the liver.

REFERENCES

1. Akbulut, M., ÇALIŞIR, S., MARAKOĞLU, T., & Coklar, H. (2009). Some physicochemical and nutritional properties of barberry (*Berberis vulgaris* L.) fruits. *Journal of food process engineering*, 32(4), 497-511.
2. Rahimi-Madiseh, M., Lorigoini, Z., Zamani-Gharaghoshi, H., & Rafieian-Kopaei, M. (2017). *Berberis vulgaris*: specifications and traditional uses. *Iranian Journal of Basic Medical Sciences*, 20(5), 569.
3. Bhardwaj, D., & Kaushik, N. (2012). Phytochemical and pharmacological studies in genus *Berberis*. *Phytochemistry Reviews*, 11(4), 523-542.
4. Aghbashlo, M., Kianmehr, M. H., & Hassan-Beygi, S. R. (2008). Specific heat and thermal conductivity of berberis fruit (*Berberis vulgaris*). *American Journal of Agricultural and Biological Sciences*, 3(1), 330-336.
5. Roshanravan, B., Yousefizadeh, S., Apaydin Yildirim, B., Farkhondeh, T., Amirabadizadeh, A., Ashrafizadeh, M., ... & Samarghandian, S. (2023). The effects of *Berberis vulgaris* L. and *Berberis aristata* L. in metabolic syndrome patients: a systematic and meta-analysis study. *Archives of Physiology and Biochemistry*, 129(2), 393-404.
6. Mokhber-Dezfuli, N., Saaidnia, S., Gohari, A. R., & Kurepaz-Mahmoodabadi, M. (2014). Phytochemistry and pharmacology of berberis species. *Pharmacognosy Reviews*, 8(15), 8-15.
7. Shinwari, Z., Khan, A., Nakaik, T., & Kyōkai, N. (2003). Medicinal and other useful plants of District Swat, Pakistan. Peshawar: Al-Aziz Communications.
8. Imanshahidi, M., & Hosseinzadeh, H. (2008). Pharmacological and therapeutic effects of *Berberis vulgaris* and its active constituent, berberine. *Phytotherapy Research*, 22(8), 999-1012.
9. Gupta, M., & Singh, A. (2022). Studies of the antimicrobial activity and phytochemical properties of *Berberis lycium*. Book Rivers.

10. Sharma, D., Jeet, K., & Kumar, S. (2024). Botanical study of *Berberis vulgaris* L. of family Berberidaceae: A review. *Research Journal of Pharmacognosy and Phytochemistry*, 16(1), 52-56.
11. Zarei, A., Changizi-Ashtiyani, S., Taheri, S., & Ramezani, M. (2015). A quick overview on some aspects of endocrinological and therapeutic effects of *Berberis vulgaris* L. *Avicenna Journal of Phytomedicine*, 5, 485-497.
12. Naaz, I., & Ali, S. A. (2018). Identification and characterization of bioactive compound berberine in the *Berberis vulgaris* root extract using HR-LC-MS analysis. *Journal of Analytical & Pharmaceutical Research*, 7, 146-150. <https://doi.org/10.15406/japlr.2018.07.00268>
13. Abu-Odeh, A. M., & Talib, W. H. (2021). Middle-East medicinal plants in the treatment of diabetes: A review. *Molecules*, 26, 742. <https://doi.org/10.3390/molecules26030742>
14. Bashir, S., Gilani, A. H., Siddiqui, A. A., Pervez, S., Khan, S. R., Sarfaraz, N. J., & Shah, A. (2010). *Berberis vulgaris* root bark extract prevents hyperoxaluria-induced urolithiasis in rats. *Phytotherapy Research*, 24, 1250-1255. <https://doi.org/10.1002/ptr.3196>
15. Yang, L., Gao, Y., Gong, J., Peng, L., El-Seedi, H. R., Farag, M. A., ... & Xiao, J. (2023). A multifaceted review of monosodium glutamate effects on human health and its natural remedies. *Food Materials Research*, 3(1).
16. Thuy, L. N., Salanta, L., Tofana, M., Socaci, S. A., Fărcaș, A. C., & Pop, C. R. (2020). A mini review about monosodium glutamate. *Bulletin UASVM Food Science and Technology*, 77(1), 1-12.
17. Chakravarty, H. L. (1976). *Plant wealth of Iraq*. Botany Directorate, Ministry of Agriculture and Agrarian Reform, Baghdad, Iraq.
18. Bergmeyer, H. U., & Rej, M. H. and R. (1985). International Federation Of Clinical Chemistry (IFCC) Scientific Committee, Analytical Section. *Journal of Clinical Chemistry and Clinical Biochemistry*, 24, 481-495.
19. Belfield, A., & Goldberg, G. M. (1971). Revised assay for serum phenyl phosphatase activity using 4-amino-antipyrine. *Enzyme*, 12, 561-573.
20. Yatizid, H. L. (1987). No title. *Journal of Clinical Chemistry*, 23, 908.
21. Friedman, R. B., & Young, D. S. (2001). *Effects of disease on clinical laboratory tests* (4th ed.). AACC Press.
22. Scimone, J., & Rothstein, R. (1978). *Laboratory Manual of Clinical Chemistry*. AVI Publishing Company, Inc., U.S.A.
23. Moder, K. (2010). Alternatives to F-test in one-way ANOVA in case of heterogeneity of variances (a simulation study). *Psychological Test and Assessment Modeling*, 52(4), 343-353.
24. Ahmed, R. R., Abdul-Hamid, M., Galal, S. R., & Hamdalla, H. M. (2019). Monosodium glutamate-induced liver microscopic and biochemical changes in male rats and the possible amendment of quercetin. *Egyptian Journal of Zoology*, 71, 44-55.
25. Mohamed, P., Radwan, R. R., Mohamed, S. A., & Mohamed, S. (2021). Toxicity of monosodium glutamate on liver and body weight with the protective effect of tannic acid in adult male rats. *Mansoura Journal of Forensic Medicine and Clinical Toxicology*, 29(2), 23-32.
26. Onobrudu, D. A., & Nwiloh, B. I. (2020). Monosodium glutamate alter hepatic functions, redox potential, and lipid metabolism: Omega-3 fatty acids ameliorative intervention. *GSC Biological and Pharmaceutical Sciences*, 13(1), 101-110.

27. Calis, I. U., Cosan, D. T., Saydam, F., et al. (2016). The effects of monosodium glutamate and tannic acid on adult rats. *Iranian Red Crescent Medical Journal*, 18(10), 1-7.
28. Eweka, A. O., & Om-Iniabo, F. A. E. (2011). Histological study of the effect of monosodium glutamate on the ovaries of adult Wistar rats. *Annals of Medical and Health Science Research*, 1(1), 37-44.
29. Karaca, M., et al. (2018). Liver glutamate dehydrogenase controls whole-body energy partitioning through amino acid-derived gluconeogenesis and ammonia homeostasis. *Diabetes*, 67, 1949-1961.
30. Eweka, A. O., & Om-Iniabo, F. A. E. (2008). Histological studies of the effects of monosodium glutamate on the liver of adult Wistar rats. *The Internet Journal of Gastroenterology*, 6.
31. Hazzaa, S. M., El-Roghy, E. S., Abd Eldaim, M. A., & Elgarawany, G. E. (2020). Monosodium glutamate induces cardiac toxicity via oxidative stress, fibrosis, and P53 proapoptotic protein expression in rats. *Environmental Science and Pollution Research*, 27, 20014-20024.
32. Al Aboud, D., Baty, R. S., Alsharif, K. F., Hassan, K. E., Zhery, A. S., Habotta, O. A., Elmahallawy, E. K., Amin, H. K., Moneim, A. E. A., & Kassab, R. B. (2021). Protective efficacy of thymoquinone or ebselen separately against arsenic-induced hepatotoxicity in rats. *Environmental Science and Pollution Research*, 28, 6195-6206.
33. Elbassuoni, E. A., Ragy, M. M., & Ahmed, S. M. (2018). Evidence of the protective effect of L-arginine and Vitamin D against monosodium glutamate induced liver and kidney dysfunction in rats. *Biomedicine and Pharmacotherapy*, 108, 7.
34. Hassan, H. A., El-Kholy, W. M., El-Sawi, M. R., Galal, N. A., & Ramadan, M. F. (2020). Myrtle (*Myrtus communis*) leaf extract suppresses hepatotoxicity induced by monosodium glutamate and acrylamide through obstructing apoptosis, DNA fragmentation, and cell cycle arrest. *Environmental Science and Pollution Research*, 27, 23188-23198.
35. Nilsson, A., et al. (2020). Quantitative analysis of amino acid metabolism in liver cancer links glutamate excretion to nucleotide synthesis. *Proceedings of the National Academy of Sciences of the United States of America*, 117, 10294-10303.
36. Torkamaneh, S., Sharifi, G. H., & Rafeian, M. (2016). The comparison between the effects of *Berberis vulgaris* extract and aerobic exercise on non-alcoholic fatty liver in male rats. *Der Pharma Chemica*, 8(1), 244-247.
37. Rad, S. Z. K., Rameshrad, M., & Hosseinzadeh, H. (2017). Toxicology effects of *Berberis vulgaris* (barberry) and its active constituent, berberine: A review. *Iranian Journal of Basic Medical Sciences*, 20, 516.
38. Ilyas, S., Tabasum, R., Iftikhar, A., Nazir, M., Hussain, A., Hussain, A., Ali, M. S., et al. (2021). Effect of *Berberis vulgaris* L. root extract on ifosfamide-induced in vivo toxicity and in vitro cytotoxicity. *Scientific Reports*, 11(1), 1708.
39. Motalleb, G., Hanachi, P., Fauziah, O., & Asmah, R. (2008). Effect of *Berberis vulgaris* fruit extract on alpha-fetoprotein gene expression and chemical carcinogen metabolizing enzymes activities in hepatocarcinogenesis rats. *International Journal of Cancer Management*, 1(1).
40. Imenshahidi, M., & Hosseinzadeh, H. (2019). Berberine and barberry (*Berberis vulgaris*): A clinical review. *Phytotherapy Research*, 33(3), 504-523.
41. Egbuonu, A. C. C., Obidoa, O., Ezeokonkwo, C. A., Ezeanyika, L. U. S., & Ejikeme, P. M. (2009). Hepatotoxic effects of low dose oral administration of monosodium glutamate in male albino rats. *African Journal of Biotechnology*, 8(13), 3031-3035.

42. Eweka, A. O., Igbigbi, P. S., & Ucheya, R. E. (2011). Histochemical studies of the effects of monosodium glutamate on the liver of adult Wistar rats. *Annals of Medical and Health Sciences Research*, 1(1), 21-30.
43. Bhattacharya, T., & Ghosh, S. K. (2019). Effect of neonatal exposure of monosodium glutamate in kidney of albino mice: A histological study. *Nepal Medical College Journal*, 21(2), 134-141.
44. Abd-Elkareem, M., Soliman, M., Abd El-Rahman, M., & Abou Khalil, N. (2022). Effect of *Nigella sativa* L. seed on the kidney of monosodium glutamate challenged rats. *Frontiers in Pharmacology*, 13.
45. Torii, K., et al. (2013). Physiological roles of dietary glutamate signaling via gut-brain axis due to efficient digestion and absorption. *Journal of Gastroenterology*, 48, 442-451.
46. Okoye, C. N., Ochiogu, I. S., & Onah, C. E. (2016). The effects of monosodium L-glutamate administration on the reproduction and serum biochemistry of adult male rabbits. *Veterinárni Medicína*, 61(3).
47. Khan, I., Najeebullah, S., Ali, M., & Shinwari, Z. K. (2016). Phytopharmacological and ethnomedicinal uses of the Genus *Berberis* (Berberidaceae): A review. *Tropical Journal of Pharmaceutical Research*, 15, 2047-2057. [CrossRef].
48. Rasool, S., Al Meslmani, B., & Alajlani, M. (2023). Determination of hypoglycemic, hypolipidemic and nephroprotective effects of *Berberis calliobotrys* in alloxan-induced diabetic rats. *Molecules*, 28(8), 3533.
49. Ismail, N. F., Ghareeb, D. A., El Hafez, E. L., El-Saadani, M. A., El-Sayed, M. M., & El Sewedy, T. S. (2022). Protective effect of *Cinnamom Zeylanicum*, *Berberis vulgaris* and *Ulva lactuca* extracts on hepatocellular toxicity induced by *Aspergillus flavus* intake in rats. *Bulletin of Egyptian Society for Physiological Sciences*, 38(2), 111-122.
50. Tahmasebi, M., Sadeghi, H., Nazem, H., Panahi Kokhdan, E., & Omidifar, N. (2018). Hepatoprotective effects of *Berberis vulgaris* leaf extract on carbon tetrachloride-induced hepatotoxicity in rats. *Journal of Education and Health Promotion*, 7(1), 147.