

Agronomic Performance of Wheat Varieties in Relation to Sowing Dates in Shivalik Range of Uttarakhand

Abstract

Wheat (*Triticum aestivum* L.) plays a vital role in global agriculture, particularly in India, the world's second-largest producer. Despite its adaptability, wheat cultivation in regions like Uttarakhand faces challenges due to population growth, limited arable land and climate change threats. This study evaluated four wheat varieties (RR-21, SY-255, PBW-550, PBW-292) across three sowing dates (10th, 20th, and 30th December) using a Randomized Block Design during the 2023–2024 Rabi season at Dev. Bhoomi Uttarakhand University's agronomy research farm in Naugaon. The study measured key agronomic parameters including plant height, Leaf Area Index (LAI), Net Assimilation Rate (NAR), Crop Growth Rate (CGR), Relative Growth Rate (RGR), and yield components. Statistical analysis using ANOVA highlighted significant differences in growth and yield attributes among sowing dates and varieties. Sowing on December 20th exhibited superior performance, notably with variety RR-21 showing higher plant height, dry weight, CGR, LAI, RGR, and NAR. Variations in yield attributes such as grains per spike, effective tillers, spike length, 1000-grain weight, grain yield, stover yield, biological yield, and harvest index were also observed. Late sowing negatively affected germination rates, leading to increased weed competition and reduced growth parameters.

Keywords: Late sown, wheat, climate change and yield

Comment [PKAFO1]: italic

Comment [PKAFO2]: cultivars

Comment [PKAFO3]: leaf area index (LAI), net assimilation rate (NAR), crop growth rate

Comment [PKAFO4]: ??

Comment [PKAFO5]: cultivar

Comment [PKAFO6]: ,

Introduction

Wheat (*Triticum aestivum* L.) is the cereal crop that is farmed on the largest scale in the world.

India is the world's second-largest producer of wheat. The introduction of high yielding cultivars is primarily responsible for this extraordinary rise in productivity. Due to its extreme adaptability, which enables it to be cultivated in non-traditional rice-growing regions of northern India as well as late-sown, challenging locations that are resistant to technological innovation, wheat makes up the majority of the crop. Wheat continues to be better in terms of area, productivity, and adaptability to a broad variety of agro-climatic conditions. However, we shouldn't be satisfied with the country's development because there are still many obstacles to overcome, such as population increase combined with dwindling arable land, diminishing water supplies, and climate change (Reynolds *et al.* 2008). Wheat is planted in Uttarakhand as a rabi crops. There is now a significant yield gap between the state's potential and realised wheat output. About 2,38, 000 hectares of wheat are farmed in Uttarakhand, yielding 632000 metric tonnes of productivity per hectare at 2,657 kg. Sowing time and wheat variety are the two most important production parameters that determine productivity. Wheat cultivation requires specific temperature conditions at various stages of its growth and development, which can vary slightly among different varieties. Ideally, wheat thrives in temperatures between 20°C to 25°C, with an upper limit of 35°C. Temperatures exceeding 30°C during the maturity phase can induce forced maturity, leading to a reduction in yield. The optimal temperature range for wheat anthesis and grain filling is between 12°C to 22°C. . In Uttarakhand, wheat is typically sown in November and concludes in late December, depending on the weather, topography, and harvesting of the preceding crop. Early in the growing season, when wheat is planted in late spring or early summer, temperatures are cool, and later in the growing season, they are hot. Optimal moisture levels are also necessary for its proper growth and development. Exposure to temperatures above this range can markedly diminish grain yield, as demonstrated by (Tewolde *et al.* 2006) and (Fisher 2007). Post-anthesis heat stress adversely affects wheat by causing several physiological changes, ultimately resulting in smaller grain weight due to a shortened grain filling period and reduced starch synthesis duration, or a combination of both effects (Hasan and Ahmed, 2005). Thus, identifying wheat varieties that thrive under heat stress and optimizing sowing times to address climate change will significantly enhance wheat production in the state of Uttarakhand.

Comment [PKAF07]: most

Comment [PKAF08]: Tewolde *et al.* 2006 and Fisher 2007.

Comment [PKAF09]: cultivars

Materials and methods

The experiment was carried out in the Rabi season of 2023–2024 at the Dev. Bhoomi Uttarakhand University's agronomy research farm in Naugaon, Dehradun, Uttarakhand, India. The site had exceptional soil qualities and was situated in a subtropical climate zone. The altitude above mean sea level was 638.96 meters. A Randomised Block Design (RBD) was applied in three replications, with four wheat varieties (RR-21, SY-255, PBW-550, and PBW-292) grown under three different planting dates (10, 20, and 30 December) included. Twelve treatments replicated thrice generated 36 plots and the plot size $2\text{m} \times 2\text{m} = 4\text{m}^2$. During the reproductive stage, the mean high and low temperatures were, respectively, 30.7 and 15.12 degrees Celsius. Collected data were statistically analyzed by the ANOVA using the MS-Excel version. The line sowing method was used to seed the wheat, spacing the rows 15 cm apart. The crop received the full dosages of P (60 kg ha⁻¹) and K (40 kg ha⁻¹) as well as half of the nitrogen (60 kg ha⁻¹) as a basal treatment. The residual nitrogen was used as a top dressing for the boot and tillering phases. The sources of nutrients for N, P, and K were urea, DAP, and nutrients of potash, in that order. In 2023, the crop was picked on April 25, April 30, and May 5. The growth, yield component, and yield statistics were recorded in accordance with standard procedure.

Comment [PKAFO10]: (spring)

Comment [PKAFO11]: 30.7 and 15.12 °C

Comment [PKAFO12]: year

Results and discussion

Analysis of variance revealed that there was a significant influence of sowing dates and varieties on the growth and yield of wheat crops.

Comment [PKAFO13]: effect

Effect of sowing dates and varieties on pre-harvest observation of wheat

Plant height at 90 DAS was recorded highest in 20th December (69.23) followed by 30th December (??) and minimum in 10th December (66.52). Among the varieties RR-21 (69.95) showed highest plant height while the lowest plant height was recorded in PBW-292 (64.25). The Dry weight of the wheat variety was recorded as maximum on 20th December (86.14 cm) and minimum on 10th December. Amongst all, variety RR-21 has shown the higher dry weight at 90 DAS and lower dry weight was observed by variety PBW 292. CGR was registered highest by RR21 (115.89) and lowest by PBW292 (105.68). Crop sown on 20th December has recorded

Comment [PKAFO14]: periods?

Comment [PKAFO15]: ??

Comment [PKAFO16]: ??

Comment [PKAFO17]: ??

Comment [PKAFO18]: ??

Comment [PKAFO19]: ??

Comment [PKAFO20]: ??

highest CGR (103.09), whereas lowest was recorded for crop sown on 10th December (79.62). Similarly, performance of variety RR-21 was significantly highest for LAI (3.17), RGR (12.83) and NAR (34.54). Among the sowing dates, crop sown on 20th December was found significantly highest for the traits i.e. LAI (3.06), RGR(12.80), NAR (24.25).

Comment [PKAFO21]: ??

Comment [PKAFO22]: ??

Comment [PKAFO23]: ??

Comment [PKAFO24]: ??

Table 1. Pre -Harvest observation of wheat (*Triticum aestivum* L.)

		Pre - Harvest observation 90 DAS					
Treatments	Plant	Dry	C.G.R	L.A.I	R.G.R	N.A.R	
	Height	weight					
Varieties							
V ₁	RR-21	69.95	88.53	115.89	3.17	12.83	34.54
V ₂	SY-255	68.14	86.44	113.65	3.13	12.80	7.76
V ₃	PBW-550	66.72	84.01	108.13	2.94	12.62	3.98
V ₄	PBW-292	64.25	78.66	105.68	2.66	12.42	2.94
	F test	S	S	S	S	S	S
	SEd (±)	0.91	0.96	2.54	0.07	0.08	6.12
	CD (P= 0.05)	1.88	2.00	4.95	0.15	0.16	12.68
Sowing Date							
S ₁	10 Dec.	66.04	82.44	79.62	2.86	12.50	7.51
S ₂	20 Dec.	69.23	86.14	103.09	3.06	12.80	24.25
S ₃	30 Dec.	66.52	84.65	101.27	3.00	12.73	5.16
	F test	S	S	S	S	S	S
	SEd (±)	0.79	0.83	2.15	0.06	0.07	5.30

Comment [PKAFO25]: m??

Comment [PKAFO26]: gram??

Comment [PKAFO27]: cultivars

CD (P= 0.05) 1.63 1.73 4.35 0.13 0.14 10.99

Effect of sowing dates and varieties on yield attributes

All yields attributes were significantly affected by the date of sowing along with the different varieties used for research study. Delayed sowing decreased grains per spike, effective tillers, spike length, 1000-grain weight, grain yield, Stover yield, biological yield and harvest index.

Sowing on December 20th significantly influenced entire attributing characters and significantly superior to the 10th and 30th December sowing. A comparison of the sowings revealed that crop sown on December 20th had on average higher grains per spike (25.42) than crop sown on 30th December (22.61). Grains per spike were registered significantly highest by RR21 whereas PBW 292 has observed the lowest no. grains per spike. Results of the analysis indicated that the spike length was lengthier for the sowing on December 20th than it was for the sowing on December 10th and December 30th. Likewise, crop sown on 20th of December had a greater number of effective tillers compared to the sowing that took place on 10th and 30th December. Characters like 1000 GW, Grain yield, biological yield and harvest index showed the similar pattern, maximum values were achieved by crop on sowing date 20th December compared to the sowing dates 10th December and 30th December.

Low germination in late-sown wheat is caused due to temperature variations. With delayed sowing, the temperature decreases and fails to meet the necessary conditions for seedling germination. Lately sown crops also encountered a lower number of tillers due to reduced germination rate, caused by lower temperatures. Differences among varieties may be due to their genetic diversity (Tahir *et al.*, 2009). The decrease in plant height, LAI, NAR, biological yield and stover yield in late sowing was due to shorter growing periods. Additionally, low germination of crop, causes weed emergence was more than usual, which resulted into crop weed competition which resulted into lesser plant height, dry weight, CGR and RGR. In contrast to this Early sown crop may have enjoyed better growing environment conditions especially solar radiation and rise in temperature, resulted into increased plant height.

These outcomes are consistent with the findings of (Shehzad *et al.* 2002). The reduction in growing time and shriveling of grain due to high temperatures that occurred throughout the milk

Comment [PKAFO28]: The results

Comment [PKAFO29]: delete

Comment [PKAFO30]: a

Comment [PKAFO31]: in

Comment [PKAFO32]: early

Comment [PKAFO33]: Shehzad *et al.* (2002)

and grain filling stages were the major causes of the test weight decrease on December 30th as a result of the delayed seeding. Additionally, (Ram et al. 2012) showed increased growth degree days, photo-thermal units, and yield characteristics in wheat that was planted immediately, which resulted in superior grain yields.

Comment [PKAFO34]: Ram et al. (2012)

Comparing variety RR-21 to the other three, it has been shown to have a greater test weight (44.68 g), biological yield (95.46 t/ha), number of effective tillers (530.99), grain per spike (25.42), and spike length (13.76). The favorable temperature that RR-21 wheat crop variety needs to accumulate more photosynthetic activity, which raises production, may be the cause of the crop variety's greater yield features. Researchers (Akhtar et al. 2002), (Kumar et al. 2005), and (Patel et al. 1999) have noted a similar pattern across the many types. RR-21 is the cultivar with the highest grain yield i.e., 42.47 t/ha. With a weight of 52.99 t/ha the cultivar RR-21 has produced the largest Stover and has the greatest harvest index (44.49%), according to the data. When seeds were planted on time, the harvest index was much greater than when seeds were sown later. Furthermore, there was no noticeable difference in the varieties' effects on the harvest index. Furthermore, (Ram et al. 2012). reported notable genotype heterogeneity in yield and harvest index.

Comment [PKAFO35]: Many authors?

Comment [PKAFO36]: Ram et al. (2012)

Table 2. Post -Harvest observation of wheat (*Triticum aestivum* L.)

		Post - Harvest observation							
Treatments	1000 GW	Grain	Stover	Biological	Harvest	Effective	Grain	Spike	
		Yield	Yield	Yield	Index	tillers	/spike	length	
V ₁	RR-21	44.68	42.47	52.99	95.46	44.49	530.99	25.42	13.76
V ₂	SY-255	42.42	41.89	52.53	94.28	44.40	523.81	24.07	13.17
V ₃	PBW-550	42.07	39.29	52.39	90.01	43.76	505.33	21.77	13.09
V ₄	PBW-292	40.22	35.82	52.02	88.35	40.51	493.82	19.50	12.62
F test		S	S	S	NS	S	S	S	S
SEd (±)		0.98	0.54	0.50	1.12	0.73	10.31	0.39	0.22

CD(P=0.05) 2.03 1.12 1.03 2.31 1.51 21.37 0.82 0.46

Sowing

Date

S ₁	10 Dec.	41.28	39.11	51.79	91.17	91.17	42.19	22.27	13.00
S ₂	20 Dec.	43.52	40.35	53.10	92.70	92.70	44.32	23.19	13.29
S ₃	30 Dec.	42.25	40.14	52.56	91.55	91.55	43.17	22.61	13.18
F test		S	S	NS	S	S	S	S	S
SEd (±)		0.85	0.47	0.43	0.97	0.97	0.63	0.34	0.19
CD(P=0.05)		1.76	0.97	0.89	2.00	2.00	1.31	0.71	0.39

References

Akhtar M, Cheema MS, Ali L and Jamil M (2002). Sowing date cum varietal trial on wheat. *Asian Journal of Plant Science*, 1(5):550-551.

Comment [PKAFO37]: ???

Anonymous (2013). Progress Report 2012-2013. Directorate of Wheat Research, Karnal, India pp.1

Fisher, R. A. 2007. Understanding the physiological basis of yield potential in wheat. J. Agric.

Hasan, M. A. and Ahmed J U. (2005). Kernel growth physiology of wheat under late planting heat stress, *J. Natl.. Sci. Foundn. Sri Lanka*, 33(3):193-204.

Comment [PKAFO38]: ???

high temperature stress. J. Agron. Crop Sci. 192: 111–120.

Kaur Anureet, Pannu RK and Buttar GS (2010). Impact of nitrogen application on the performance of wheat (*Triticum aestivum*) and nitrogen use efficiency under different dates of sowing. Indian Journal of Agronomy 5(1):40-45.

Comment [PKAFO39]: italic

Kumar Satish, Kadian VS, Singh RC and Malik RK (2005). Effect of planting date on performance of wheat (*Triticum aestivum*) genotypes. Indian Journal of Agricultural Sciences 75(2):03-105.

Comment [PKAFO40]: ??

Mishra B (2006). Wheat research towards national food security. Intensive Agriculture 2:11-13

Muhammad Tahir, Asghar Ali, Ather Nadeem Muhammad, Hussain, Akhtar and Khalid Farhan (2009). Effect of different sowing dates on growth and yield of wheat (*Triticum aestivum*) varieties in district Jhang. Pakistan Journal of Life Science 7(1):66-6

Mukherjee D (2012). Effect of sowing dates on growth and yield of wheat (*Triticum aestivum*) cultivars under mid hill situation of West Bengal. Indian Journal of Agronomy 57(2):152-156.

Comment [PKAFO41]: ???

Pandey B, Pandey RK, Dwivedi DK and Singh RS (2010). Phenology, heat unit requirement and yield of wheat (*Triticum aestivum*) varieties under different crop growing environment. Indian Journal of Agricultural Sciences 80(2):136-140.

Comment [PKAFO42]: ??

Patel SR, Thakur DS and Lal Nageshwar (1999). Yield and nutrient uptake of wheat (*Triticum aestivum*) varieties under different sowing dates. Indian Journal of Agronomy 44(4):276-278.

Comment [PKAFO43]: ??

Ram H, Singh G, Mavi GS and Sohu VS (2012). Accumulated heat unit requirement and yield of irrigated wheat (*Triticum aestivum* L.) varieties under different crop growing environment in central Punjab. Journal of Agrometeorology 14(2):147-153.

Comment [PKAFO44]: ??

Reynolds, M. P., Dixon, J., Ammar, K., Kosina, P. & Braun, H.-J. (2008b). Stakeholders' priorities for internationally-coordinated wheat research. In :Reynolds M.P., J. Pietragalla, and H.-J. Braun, (eds) International Symposium on Wheat Yield Potential: Challenges to International Wheat Breeding. Mexico, D.F.: CIMMYT.

Sci. 145: 99-113.

Shahzad K, Bakht J, Shah WA, Shafi M and Jabeen N (2002). Yield and yield components of various wheat cultivars as affected by different sowing dates. Asian Journal of Plant Sciences 1(5):522-525.

Tahir, M., Ali, A., Nadeem, M. A., Hussain, A., & Khalid, F. (2009). Effect of different sowing dates on growth and yield of wheat (*Triticum aestivum* L.) varieties in district Jhang, Pakistan. Pak. J. Life Soc. Sci, 7(1), 66-69.

Comment [PKAF045]: ??

Tewolde, H., C. J. Fernandez and C. A. Erickson. 2006. Wheat cultivars adapted to post-heading

UNDER PEER REVIEW