

"Influence of Organic Nutrient Enrichment on Growth Pattern, Flowering and Biochemical Profiling in Bottle Gourd (*Lagenaria siceraria*) (Mol.) Standl.) Var. Kashi Ganga"

ABSTRACT

The present investigation was carried out with title at Organic Research Farm (ORF), Department of Horticulture, Institute of Agricultural Sciences, Bundelkhand University, Karguanji, Jhansi, Uttar Pradesh during the *Kharif*-2022 with a view to identify the effects of different combinations of organic nutrition and its role in growth, yield and quality of Bottle gourd variety Kashi Ganga. The experiment was laid in Randomized block design with 8 treatments and 3 replications with different combination in RDM and application of organic nutrition. Under this experiment, overall, 8 treatment was taken T₀ Control (water spray), T₁ (100% Farmyard Manure), T₂ (100% Poultry Manure), T₃ (100% Vermicompost), T₄ (50% Farmyard Manure + 50% Neem cake), T₅ (50% Farmyard Manure + 50% Vermicompost), T₆ (50% Neem cake + 50% Poultry Manure) and T₇ (33% Farmyard Manure + 33% Vermicompost + 33% Poultry Manure). Based on above experimental finding it may be concluded that the treatment T₇ (33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost) was found to be best in the terms of growth yield and quality of Bottle gourd. While the highest net return was found in the T₇ and the highest B:C ratio was found in the same with 6.10.

Keywords: Bottle gourd, Farmyard Manure, Neem Cake, Poultry Manure, Vermicompost

INTRODUCTION

Bottle gourd, scientifically known as *Lagenaria siceraria* (Mol.) Standl., Calabash, also known as bottle gourd, is a vine grown for its fruit. Bottle gourd are horticulture crop belongs to the family *Cucurbitaceae* bearing chromosome number $2n=22$ [31]. fruits have a variety of shapes: they can be huge and rounded, small and bottle-shaped, or slim and serpentine, and they can grow to be over a metre long [20].

The bottle gourd may have been carried from Asiatic Africa, Europe, and the Americas during human migration, [19] or by seeds floating across the oceans inside the gourd. It has been proven to have been globally domesticated and existed in the New World) during the Pre-Columbian era. Bottle gourd is a diploid of parentage [19]. The bottle gourd is a commonly cultivated plant in tropical and subtropical areas of the world and was eventually domesticated in southern Africa. Stands of *L. siceraria*, which may be source plants and not merely domesticated stands, were reported in Zimbabwe [16]. Bottle gourds are grown by direct sowing of seeds or transplanting 15- to 20-day-old seedlings. The plant prefers well-drained, moist, organic rich soil [22].

The plant produces night blooming white flowers. The male flowers have long peduncles, and the females have short ones with an ovary in the shape of the fruit [29].

Hand pollination can be used to solve the problem. Pollen size is ~60 microns [27]. Raw Bottle gourd is 96.1% water, 0.2% protein, 2.5% carbohydrates and contains negligible fat. In a 100-gram reference amount, raw bottle gourd is a source vitamin C (6 mg) with moderate contents of thiamine (0.23 mg), Nicotinic acid (0.20 mg) and magnesium (5 mg), Calcium (20 mg), Phosphorous (10 mg), Potassium (87 mg) [23]. Bottle gourd has valuable anti-cholesterol property. In India area under Bottle gourd production in India accounts to 1.58 million hectares with production of 31.42 metric tonnes in year 2021-22 [18].

Comment [KM1]: Grammar Mistake

Comment [KM2]: Each Letter Capitalize (RBD)

Comment [KM3]: With repeating 2 times in a single sentence

Comment [KM4]: Include expansion

Comment [KM5]: Remove the treatment details and add results

Bihar ranks first in area and production of Bottle gourd in year 2021-22 followed by Uttar Pradesh and Madhya Pradesh. In Uttar Pradesh area under production is 2.38 thousand hectares while production is estimated to be 5.09 million tonnes for year 2021-22. (Source: NHB, Ministry of Agriculture & Farmers Welfare, Government of India, 2021-22.) [21].

Organics are plant and animal wastes that are used as a source of nutrients for plants. They release nutrients when broken down. Natural or synthetic chemicals that contain nutrients for plants are called fertilizers [18]. Though, Manures contains low nutrient per unit amount not only improve soil properties, but also have a longer residual effect than fertilizers with a high nutrient content [29].

Applying FYM improves soil fertility. On an average well decomposed farmyard manure contains 0.5 per cent Nitrogen (N), 0.2 per cent Phosphate (P_2O_5) and .0.5 per cent Potassium (K_2O) [28].

Vermicompost is rich in NKP (nitrogen 2-3%, potassium 1.85-2.25% and phosphorus 1.55- 2.25%), micronutrients, beneficial soil microbes and contain 'plant growth hormones & enzymes [25,26,27].

Being a totally botanical product Neem cake contains 100% natural NPK content and other essential micro nutrients as N (Nitrogen 2.0% to 5.0%), P (Phosphorus 0.5% to 1.0%), K (Potassium 1.0% to 2.0%), Ca (Calcium 0.5% to 3.0%), Mg (Magnesium 0.3% to 1.0%), S (Sulphur 0.2% to 3.0%), Zn (Zinc 15 ppm to 60 ppm), Cu (Copper 4 ppm to 20 ppm), Fe (Iron 500 ppm to 1200 ppm), Mn (Manganese 20 ppm to 60 ppm). It is rich in both sulphur compounds and bitter limonoids [20,18,16].

Poultry manure contains higher nitrogen and phosphorus compared to other bulky organic manures. The average nutrient content is 3.03 percent N; 2.63 per cent P_2O_5 and 1.4 per cent K_2O . Organics such as FYM, Poultry manure and Vermicompost not only promote increased yields and improved crop quality, but also maximize the genetic potential of plants and the presence of nutrients enhances root development, fruit set, affects the vitality and health of plants [11,22,24].

Nutrients are fundamental for balanced nutrition and a tremendous tool to help farmers in increasing crop yield and quality [13,14,17]. The current experiment was conducted in order to help in understanding the growth pattern, flowering and biochemical profiling in bottle gourd by organic nutrient enrichment. Therefore, present study of analysis in Bottle gourd crop was carried out to identify the best treatment combination with high yield, early maturing suited to Bundelkhand zones of U.P.

MATERIAL AND METHODS

The present investigation was done to understand the effect of organic nutrition at different doses combination on fruit growth, yield and quality of Bottle gourd variety Kashi ganga. The details of the materials used, and the procedures adopted in the investigation, which was carried out at Organic Research Farm (HRF), Karguanji, Jhansi Department of Horticulture, Institute of Agricultural Sciences, Bundelkhand University, Jhansi during the *Kharif* season of 2022 are described under the following heads.

In general, soil properties of experimental site showed a typical sandy soil of Bundelkhand region of Uttar Pradesh. Soil was sandy loam in texture, slightly alkaline in reaction and having low electrical conductivity, very low in organic carbon, low in available nitrogen, low in available phosphorus and moderately high in available potassium given in table 2 and 3. Observations were recorded at different stages of growth periods. The data were statistically analysed by the method suggested by [30].

Table 1 Treatment Details

S. No.	Treatment Symbols	Treatment combination
1	T ₀	Control (Water spray)
2	T ₁	100% Farmyard Manure
3	T ₂	100% Vermicompost
4	T ₃	100% Poultry Manure
5	T ₄	50% Farmyard Manure + 50% Neem cake
6	T ₅	50% Farmyard Manure + 50% Vermicompost
7	T ₆	50% Neem cake + 50% Poultry Manure
8	T ₇	33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost

Table 2 Physical and chemical properties of soil at Organic Research Farm (IAS, BU, Karguanji, Jhansi).

Comment [KM6]: in

S. No.	Particulars	Mechanical Properties
1	Fine Sand (%)	42.0
2	Silt (%)	38.2
3	Clay (%)	20.1
4	Textural class	Sandy loam

Table 3: Chemical composition of the soil

S. No.	Ingredients	Quantity
1	Soil pH	7.1
2	Organic carbon (%)	0.52
3	Available Nitrogen (N)	142.33 kg/ha
4	Available Phosphorus (P)	4.56 kg/ha
5	Available Potash (K)	206.11 kg/ha

Source: Soil analysis was done by KVK (Krishi Vigyan Kendra, Jhansi, U.P.).

RESULTS AND DISCUSSION

A) Growth Parameters

The data pertaining to Vine length is presented in Table 4. for various days of transplanting. The treatment with the combined use of 33% Farmyard Manure, 33% Poultry Manure, and 33% Vermicompost (T₇) resulted in the longest vine length across all stages of growth, with a significant increase from 2.09 m at 30 DAS to 3.33 m at 60 DAS. This suggests that a balanced combination of organic amendments can maximize vine elongation, likely due to a well-rounded nutrient profile. Treatment T₁ (100% Farmyard Manure) showed moderate vine growth with 1.68 m at 45 DAS and 2.19 m at 60 DAS, indicating that farmyard manure can be beneficial but not as effective as the combination of inputs followed by T₂ (100% Vermicompost) also contributed to

substantial vine length, especially at 45 DAS (1.83 m) and 60 DAS (2.10 m). Vermicompost may enhance soil structure and nutrient availability, contributing to improved vine growth [5,8,9,10]. T₃ (100% Poultry Manure) showed similar trends, with good vine growth observed at 45 DAS (1.85 m) and 60 DAS (2.22 m), suggesting that poultry manure can also have a positive effect on vine elongation [3,6,9,11].

The combined treatments, particularly T₄ (50% Farmyard Manure + 50% Neem Cake) and T₅ (50% Farmyard Manure + 50% Vermicompost), resulted in better vine growth than individual treatments. Treatment T₅ (50% Farmyard Manure + 50% Vermicompost) had the highest vine length at 60 DAS (2.93 m), surpassing the individual manure treatments, indicating the synergistic effect of combining different organic amendments [1,3,6,24,26]. However, the treatment control (T₀) showed the shortest vine lengths at all stages (0.96 m at 30 DAS, 1.34 m at 45 DAS, and 1.70 m at 60 DAS), supporting the hypothesis that organic amendments promote better vine growth compared to no treatment [2,4,5,25].

2. Number of branches per plant

Treatment T₂ (100% Vermicompost) showed a significant increase in the number of branches (5.17), suggesting that vermicompost may be particularly effective in enhancing branching due to its nutrient richness and ability to improve soil structure. T₃ (100% Poultry Manure) also promoted a higher number of branches (4.67), indicating that poultry manure is beneficial for branching, possibly due to its high nitrogen content [7,9,11,12].

The combination treatments, such as T₅ (50% Farmyard Manure + 50% Vermicompost) (4.58 branches) and T₆ (50% Neem cake + 50% Poultry Manure) (4.42 branches), showed improved branching compared to the control and individual treatments, but they were not as effective as the combined treatment of T₇. Control Treatment (T₀): The control (T₀) exhibited the lowest number of branches (3.85), confirming that organic treatments significantly enhance the number of branches per plant compared to no external amendments [8,10,15].

B) Earliness parameters

1. Days to first male flowering, female flowering, and days to first fruit picking

Among the different applications of organic nutrition, the minimum days to first male flowering was seen in T₇ with 34.67 days, followed by T₁ with 35.78 days whereas maximum days to first flowering 40.08 days was recorded in T₀. Among the application of organic nutrients minimum days to first female flowering was seen in T₇ with 44.17 days, followed by T₁ with 44.63 days whereas maximum days to first female flowering 50.36 days was recorded in T₀. Among the applications of organic nutrition, the minimum days to first fruit picking was seen in T₇ with 75.92 days, followed by T₁ with 76.75 days whereas maximum days to first fruit picking 82.12 days was recorded in T₀. The data of three parameters are given in table 5. The application of organic treatments significantly influenced the timing of flowering and fruit picking in bottle gourd. In the present experiment treatment **T₇ (33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost)** was the most effective treatment, resulting in the earliest male and female flowering as well as the quickest fruit picking. This suggests that a balanced combination of organic inputs is beneficial for accelerating the growth and development of bottle gourd followed by other treatments like **T₁ (100% Farmyard Manure)** and **T₂ (100% Poultry Manure)** also showed good results but were less effective than the combined treatments. The control group showed the latest flowering and fruit picking, reinforcing the advantages of organic amendments in enhancing crop performance as referred by [11,14,17,19].

C) Yield parameters

Fruit yield per plant (kg/plant) and Fruit yield per hectare (q/ha)

The maximum average yield per plant (11.47 kg/plant) were recorded in treatment T₇ followed by T₆ i.e., 10.74 kg/plant and the lowest average yield per plant (7.10 kg/plant) were observed in T₀. The maximum average yield per hectare (611.96 q/ha) were recorded in treatment T₇ followed by T₆ i.e., 572.92 q/ha and the lowest average yield per hectare (378.77 q/ha) were

Comment [KM7]: Check Spelling

observed in T₀. Nutrition play an important role in improving productivity and quality of fruit. Nutrition's doses increased the vigour of plants, assimilating area, size of fruit, thereby resulting into higher weight of fruit. These results are in close conformity with the findings of [9,17,20].

D) Total soluble solid (⁰Brix) and Protein content

From the data presented in table 6. The maximum T.S.S. (6.28 ⁰Brix) was observed in treatment T₇ followed by T₄ with 5.10 ⁰Brix. The minimum T.S.S. (2.81 ⁰Brix) was noticed in treatment T₀. The Ascorbic acid content among different treatments varied significantly. The maximum Ascorbic acid content (4.20 mg/100g) was observed in treatment T₇ followed by T₄ with 4.06 mg/100g. The minimum Ascorbic acid content (2.93 mg/100g) was noticed in treatment T₀. The maximum protein content was reported under the treatment T₇ (33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost) followed by T₅ (50% Poultry Manure + 50% Vermicompost) with 4.20. It may be noted that nutrition play an important role in improving productivity and quality of fruit. Nutrition's doses increased the vigour of plants, assimilating area, size of fruit, thereby resulting into higher weight of fruit. These results are in close conformity with the findings of [10,15,18,19].

Conclusion

From the above experimental finding it may be concluded that the treatment T₇ (33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost) was found to be best in the terms of growth, yield and quality of Bottle gourd.

References

1. Abbas, Z., M. Mubashir, U. Riaz, Z. Javid, M. Ashraf, S. Rehman, M.J. Qamar, A. Zulqadar and S.M. Mehdi. (2019). Combined impacts of compost, poultry manure and NPK fertilizers on yield of bottle gourd plant *Lagenaria siceraria* (Mol.) Standl *Pakistan Journal of Agricultural Research*. 32(3): 456-465.
2. Abha, R., Meena, R. L., Singh, R. and Mandal, R. K. (2019). Effect of various organic manures on growth and yield of bottle gourd *Lagenaria siceraria* (Mol.) Standl *Journal of Pharmacognosy and Phytochemistry*. 8(6): 1203-1205.
3. Adekunle, O. O. (2013). The Effect of Combined Application of Poultry Manure and Sawdust on the Growth and Yield of Bottle gourd. *Journal of Agricultural Science*. 5(10): 169-175.
4. Aderemi, F. T., Aderemi, A. M., Elesho, R. O., Aluko, A. K. and Oyeleke, O. S (2020). Comparative Effects of Different Manure on Growth and Yield of Bottle gourd *Lagenaria siceraria* (Mol.) Standl. *International Journal of Research and Scientific Innovation*. 7(8): 181-183.
5. Adesida, O. A., Smart, M. O., Isola, J. O., Kehinde, O. M. (2020). Methods of poultry manure application and its effect on growth and yield of bottle gourd *Lagenaria siceraria* (Mol.) Standl *Journal of Research in Forestry, Wildlife & Environment*. 12(1): 121- 128.
6. Adhikari A. and Piya, A. (2020). Effect of Different Sources of Nutrient on Growth and

Yield of Bottle gourd (*Abelmoschus esculentus* L. Monech). *International Journal of Environmental & Agriculture Research*. 6(1): 45-50.

7. Agricultural and Processed Food Products Export Development Authority (2020-21). Horticultural Statistics data.
8. Agriculture Statistics at a glance (2021). Directorate of Economics & Statistics. *Government of India Ministry of Agriculture & Farmers Welfare Department of Agriculture & Farmers Welfare*. 130pp.
9. Alam, S. M., Muhammad, A. U., Syed, I. H., Nausherwan, N. N., Syeda, S. A. and Imdad, M. (2019). Effect of Farmyard Manure and Planting Densities on Growth, Yield and Quality of Bottle gourd under Natural Farming. *International Journal of Research in Agriculture and Forestry*. 6(4): 21-25.
10. Amran, H. A., Prasad, V. M. and Saravanan, S. (2014). Effect of FYM on Growth, Yield and Fruits Quality of Bottle gourd *Lagenaria siceraria* (Mol.) Standl. *Journal of Agriculture and Veterinary Science*. 7(3): 7-12.
11. Anonymous, (2019). *Horticultural Statistics at a Glance*, Horticulture statistics division department of agriculture, cooperation & farmers welfare ministry of agriculture & farmers welfare government of India.
12. Bamboriya, J. S., Naga, S. R., Sharma, S. R., Choudhary, M. R. and Bamboriya, S. D. (2018). Productivity, Quality and Profitability of Bottle gourd (*Abelmoschus esculentus*) as Influenced by Organic Manures and Biofertilizers. *International Journal of Bio-resource and Stress Management*. 9(4):506-509.
13. Chaudhary, S. K., Sushil, K. Y., Mahto, D. K., Neha, S. and Sanjay, K. S. (2019). Effect of Organic and Inorganic Sources of Nutrients on Growth and Yield of Bottle Gourd (*Lagenaria siceraria* (Mol.) Standl.) in Bihar. *Current Journal of Applied Science and Technology*. 37(6): 1-7.
14. D. V. S. Chauhan (1972). *Vegetable Production in India*. Ram prasad and Sons, Agra India. 255 pp.
15. De Candolle A. *Origin of cultivated plants*. Hafner Publication Co., New York, (Reprint of 2nd edition. 1959).
16. Decker-Walters, Deena S; Wilkins-Ellert, Mary; Chung, Sang-Min; Staub, Jack E (2004). "Discovery and Genetic Assessment of Wild Bottle Gourd [*Lagenaria Siceraria* (Mol.) Standl; Cucurbitaceae] from Zimbabwe". *Economic Botany*. 58(4): 501–508.
17. Devanda, P., Lakhawat, S. S., Pilania, S., Sharma, S. K., Azad, M., Dudi, D. P. S., Sharvan, K. Y. and Pratihtha, D. (2021). Effect of Organic Manures and Liquid Formulations on Growth, Yield and Quality of Bottle gourd [*Lagenaria Siceraria* (Mol.)

- Standl cv. Arka Anamika. *International Journal of Current Microbiology and Applied Sciences*. 10(06): 426-433.
18. Directorate of Economics and Statistics, (2020-21) Ministry of Agriculture & Farmers Welfare (DAC & FW), Government of India, 2020-21).
19. Erickson, D. L; Smith, B. D; Clarke, A. C; Sandweiss, D. H; Tuross, N (2005). "An Asian origin for a 10,000-year-old domesticated plant in the Americas". *Proceedings of the National Academy of Sciences*. 102(51): 18315–18320.
20. Fagwalawa, L. D. and Yahaya, S. M. (2016). Effect Organic Manure on The Growth and Yield of Bottle gourd. *Imperial Journal of Interdisciplinary Research*. 2(3): 130-133.
21. FAOSTAT (2020). www.fao.org/faostat/en/data/CC. Food Supply- Crops Primary Equivalent. Visited on 15/12/2021.
22. Farnoosh Brock (2013). *The Healthy Juicer's Bible: Lose Weight, Detoxify, Fight Disease, and Live Long*. Skyhorse 6th edition. 214 pp.
23. Fisher, R. A. and Yates, F. (1963). *Statistical Tables for Biological, Agricultural and Medical Research*. Oliver and Boyd, London: 143 p.
24. Ghayal, R. G., Vaidya, K. P. and Tapkeer, P. B. (2017). Effect of different organic manures and inorganic fertilizers on chemical properties of cucumber (*Cucumis sativus* L.) in lateritic soils of Konkan. *International Journal of Chemical Studies*. 5(6): 1626-1630.
25. Ibeh, C. U. Nsoanya, L. N. Nweke, I. A. (2019). Effect of Poultry Manure on Growth and Yield of Local Varieties of Bottle gourd [*Lagenaria Siceraria* (Mol.) Standl. *SSRN Electronic Journal*. 3(1): 22-29.
26. Ibrahim, U., and Hamma, I. L. (2012). Influence of Farmyard Manure and Weeding Regimes on Growth and Yield of Bottle gourd *Lagenaria siceraria* (Mol.) Standl in Zaria. *World Journal of Agricultural Sciences*. 8(5): 453-458.
27. Imnatamsu, Neelam, K., Sandeep, R. and Ronald, D. (2020). Effect of Organic and Inorganic Fertilizers on Growth and Yield of Bottle Gourd (*Lagenaria siceraria*) Under Teak Based Agroforestry System. *Indian Journal of Pure and Applied Biosciences*. 8(3): 388-395.
28. Jackson, M. L., 1973, *Soil and Chemical Analysis*, Prentice Hall of India Private Limited, New Delhi.
29. Kartina, A. M., Utama, P. and Dimiyati, I. (2018). Response growth and yield of bottle gourd [*Lagenaria Siceraria* (Mol.) Standl] to giving dosage levels of chicken manure and variation of plants spacing. *IOP Conference Series: Earth and Environmental Science*. 383(1): 1-9.

30. Yates, F. (1964). Sir Ronald Fisher and the design of experiments. *Biometrics*, 20(2), 307-321.

Table 4 Effect of organic nutrition on Vine length (m) of the Bottle gourd.

Treatment Notation	Treatment details	30 DAS (in m)	45 DAS (in m)	60 DAS (in m)
T ₁	100% Farmyard Manure	1.11	1.68	2.19
T ₂	100% Vermicompost	1.55	1.83	2.10
T ₃	100% Poultry Manure	1.48	1.85	2.22
T ₄	50% Farmyard Manure + 50% Neem cake	1.65	1.94	2.23
T ₅	50% Farmyard Manure + 50% Vermicompost	1.26	2.10	2.93
T ₆	50% Neem cake + 50% Poultry Manure	1.51	1.88	2.25
T ₇	33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost	2.09	2.71	3.33
T ₀	Control	0.96	1.34	1.70
	Mean	1.45	1.92	2.37
	'F' Test	S	S	S
	C.V.	12.19	7.55	6.55
	S.E.(m)	0.10	0.08	0.09

C.D. at 5%	0.31	0.26	0.27
-------------------	-------------	-------------	-------------

Table 5 Effect of plant growth regulators on the number of branches per plant of bottle gourd.

Treatment Notation	Treatment details	In numbers
T₁	100% Farmyard Manure	4.33
T₂	100% Vermicompost	5.17
T₃	100% Poultry Manure	4.67
T₄	50% Farmyard Manure + 50% Neem cake	4.17
T₅	50% Farmyard Manure + 50% Vermicompost	4.58
T₆	50% Neem cake + 50% Poultry Manure	4.42
T₇	33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost	5.30
T₀	Control	3.85
Mean		4.56
'F' Test		S

Comment [KM8]: plant growth regulators ?

C.V.	8.99
S.E.(m)	0.24
C.D. at 5%	0.73

Table 6 Effect of organic nutrients on days to first male flowering, first female flowering and first fruit picking of Bottle gourd.

Treatment Notation	Treatment details	Days to first male flowering	Days to first female flowering	Days to first fruit picking
T₁	100% Farmyard Manure	35.78	44.63	76.75
T₂	100% Poultry Manure	37.25	45.00	76.75
T₃	100% Vermicompost	36.50	45.92	77.67
T₄	50% Farmyard Manure + 50% Poultry Manure	36.75	45.83	77.59
T₅	50% Poultry Manure + 50% Vermicompost	38.75	46.67	78.42
T₆	50% Farmyard Manure + 50% Vermicompost	38.17	47.66	79.41
T₇	33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost	34.67	44.17	75.92
T₀	Control	40.08	50.36	82.12
	Mean	41.51	37.24	46.28
	'F' Test	S	S	S
	C.V.	2.76	2.02	1.25
	S.E.(m)	0.59	0.54	0.56

C.D. at 5%	1.82	1.66	1.72
-------------------	-------------	-------------	-------------

Table 7 Effect of organic nutrition on Average fruit yield per plant and fruit yield per hectare of Bottle gourd.

Treatment Notation	Treatment details	Fruit yield per plant (kg/plant)	Fruit yield per hectare (t/ha)
T₁	100% Farmacyard Manure	8.36	445.90
T₂	100% Poultry Manure	7.41	395.29
T₃	100% Vermicompost	8.47	451.48
T₄	50% Farmacyard Manure + 50% Poultry Manure	8.63	460.07
T₅	50% Poultry Manure + 50% Vermicompost	9.63	513.60
T₆	50% Farmacyard Manure + 50% Vermicompost	10.74	572.92
T₇	33% Farmacyard Manure + 33% Poultry Manure + 33% Vermicompost	11.47	611.96

T₀	Control	7.10	378.77
	Mean	1.63	8.98
	'F' Test	S	S
	C.V.	7.32	7.34
	S.E.(m)	0.37	19.96
	C.D. at 5%	1.14	61.13

Comment [KM9]: Please check the Yield. The average bottle gourd yield is 35.40m

Table 8 Effect of organic nutrition on Total soluble solid and Protein content of Bottle gourd.

Treatment Notation	Treatment details	T.S.S.	Protein Content
T₁	100% Farmyard Manure	3.80	3.70
T₂	100% Poultry Manure	4.70	3.67

T₃	100% Vermicompost	3.72	3.80
T₄	50% Farmyard Manure + 50% Poultry Manure	5.10	4.06
T₅	50% Poultry Manure + 50% Vermicompost	4.34	4.13
T₆	50% Farmyard Manure + 50% Vermicompost	3.78	3.80
T₇	33% Farmyard Manure + 33% Poultry Manure + 33% Vermicompost	6.28	4.20
T₀	Control	2.81	2.93
	Mean	4.40	4.32
	'F' Test	S	S
	C.V.	1.32	6.36
	S.E.(m)	0.03	0.14
	C.D. at 5%	0.10	0.43