

1
2
3
4
5
6
7
8
9

Incidence of *Alternaria* fruit rot as a latent postharvest disease in guava (*Psidium guajava* L.) in Kerala.

10
11

ABSTRACT

A purposive random sampling survey was conducted to study the incidence of latent pathogens associated with postharvest losses in guava sourced from markets and homesteads of Ernakulam, Thrissur and Palakkad districts of Kerala during 2024. Fruits at mature green stage were collected then surface sterilized and incubated at room temperature for symptom development. After three to five days of incubation, few of the fruits collected from Thrissur district started development of dry brown lesions. The disease developed under high humidity and temperature between 30 and 35 °C. Four isolates of *Alternaria* sp. were isolated from the fruits. The highest disease incidence recorded was 75 % from Vettikkal while the highest severity of 20 % was recorded in fruits collected from Shakthan Market. The isolated pathogen appeared white to grey coloured, submerged to aerial mycelia which showed uniform and zonal growth at regular intervals. The isolates exhibited an average growth rate of 0.6-0.9 cm per day. The hyphae were brown-coloured, septate with straight or curved, geniculate, pale brown conidiophores bearing a branched chain of conidia. The conidia were brown, obclavate in shape with 3-4 transverse and 1-2 longitudinal septa having average size of 22.08 to 29.53 × 7.85 to 12.54 µm. Molecular characterization using ITS primers followed by sequence analysis in the NCBI database, identified the pathogen as *Alternaria alternata*.

12
13
14
15

Keywords: guava, postharvest diseases, fruit rot, Alternaria alternata

1. INTRODUCTION

16
17
18
19
20
21
22
23
24
25
26
27
28

Guava (*Psidium guajava* L.), a member of the Myrtaceae family, is native to South America (specifically Peru) and is widely cultivated in tropical and subtropical regions due to its hardiness and adaptability. It contains high amounts of vitamins A, B1, B2, dietary minerals, lutein, zeaxanthin, lycopene, folic acid, potassium, copper, and manganese (Rahman *et al.*, 2003). Guava is also a rich source of vitamin C and ranks second to Aonla in terms of vitamin C content. As one of the most important fruit crops in terms of area and production, after mango, banana, and citrus, guava plays a vital role in human nutrition and health. However, despite its nutritional benefits, guava's perishable nature makes it highly susceptible to significant postharvest losses. One of the major contributors to these losses is fungal infection, which can occur at various stages of the fruit's lifecycle, including during the growing season, harvesting, handling, transport, post-harvest storage, and marketing.

29
30
31

Alternaria alternata, is a destructive pathogen which is responsible for various diseases during the postharvest shelf-life of many horticultural crops. For instance, *Alternaria* causes stem-end rot in mangoes (Amin *et al.*, 2011), fruit spot on apples (Harteveld *et al.*,

2013), black rot in kiwi fruit (Kwon *et al.*, 2011), and *Alternaria* rot on figs (Doster and Michailides, 2007). Studies by Prusky *et al.* (2002) have shown that *Alternaria alternata* can remain quiescent on the cuticles of unripe mango and avocado fruits but can later colonize the mesocarp of these fruits once peeled. This suggests that specific changes occurring during fruit ripening may trigger the transition to active infection. Similarly, *Alternaria* fruit rot has been reported in guava, with Srivastav and Lal (2009) from fruit markets of Allahabad. Nongmaithem (2014) and Fatima (2019) also reported *Alternaria* fruit rot in guava. Fungi like *Alternaria* are capable of sporulating and resisting postharvest treatments, enabling them to infect fruits even during transportation to international markets. Given the significant impact of *Alternaria* infections on postharvest quality of guava fruit, this study aims to document the incidence and severity of *Alternaria* fruit rot in guava in Ernakulam, Thrissur and Palakkad districts in the areas delineated under agroecological units AEU 10 (North Central Laterites) and AEU 12 (Southern Foothills) of Kerala, India. Additionally, the study will examine the disease development in guava under laboratory conditions and the cultural-morphological characteristics and pathogenicity of the pathogen.

2. MATERIAL AND METHODS

2.1 SURVEY AND COLLECTION OF FRUIT SAMPLES

The study was conducted during 2024 at the Department of Plant Pathology, College of Agriculture, Vellanikkara, Kerala. Purposive random sampling surveys were carried out in homesteads and markets of Ernakulam, Thrissur and Palakkad districts delineated under agroecological unit AEU 10 (North Central Laterites) and AEU 12 (Southern Foothills) of Kerala. Apparently, healthy fruits were collected from different places, surface sterilized with a 1 % sodium hypochlorite solution for 5 min and washed three times with sterile distilled water then incubated at room temperature for disease development. The disease symptoms that developed were extensively examined and documented to understand the expression and progression of latent postharvest diseases under natural conditions. Disease incidence and severity were assessed using the formulae given below and expressed as percent disease incidence and percent disease severity (Lakshmi *et al.*, 2011; Wheeler, 1969).

$$\text{Percent disease incidence (PDI)} = \frac{\text{Number of fruits infected}}{\text{Total number of fruits}} \times 100$$

$$\text{Percent disease severity} = \frac{\text{Sum of disease ratings}}{\text{Total number of samples assessed} \times \text{maximum disease grade}} \times 100$$

2.2 ISOLATION OF PATHOGEN AND PATHOGENICITY TESTS

The pathogens associated with symptomatic fruit specimens were isolated using the tissue segmentation method (Rangaswamy, 1958). The fruits were surface sterilized using 1 % sodium hypochlorite solution for 3 min and washed thrice with sterile distilled water. After disinfection, the diseased fruit sample bit was cut into small pieces of 5 mm and placed over PDA media. After incubation, the hyphal tips of the fungal colonies were transferred aseptically to fresh PDA media in Petri plates. The fungal cultures were maintained in PDA slants and stored under refrigeration (4 °C) for future studies. The pathogenicity of fungal isolates was tested to prove the Koch postulates. The mycelial bit inoculation method explained by Rocha *et al.* (1998) was followed for the inoculation of pathogens to confirm pathogenicity. The healthy, semi-ripe guava fruits were surface sterilized by dipping in sodium hypochlorite (1 %) for 15 min followed by three washings with sterile distilled water. Mycelial bits taken from the margins of actively growing colonies (seven days old culture) of the pathogen were placed on

82 healthy fruits. The fruits were pinpricked with a sterile dissection needle before inoculation to
83 facilitate better pathogen establishment. Fruits without pinpricks were also inoculated, and
84 both types were kept under room conditions for disease development. Control was maintained
85 using a PDA disc that was free of pathogens. Moist cotton swab was placed over the mycelial
86 bits or PDA disc. The fruits were then covered in polypropylene cover and maintained at room
87 temperature for symptom development. When the symptoms appeared, the pathogen was re-
88 isolated from the symptomatic fruits into PDA media. The re-isolated pathogen culture was
89 compared with the original culture to prove Koch's postulates.

90 **2.3 CHARACTERISATION OF PATHOGENS**

91
92
93 The cultural characteristics of the pathogen in terms of growth, viz., texture, colour,
94 growth rate, sporulation, and reverse side colour on Petri plates were observed. The
95 morphological identification of the fungus was done based on texture, pigmentation,
96 characteristics of conidia in comparison with standard descriptors. Molecular characterization
97 was used to confirm the species of the isolated pathogen. The genomic DNA was extracted
98 using the modified protocol of Doyle (1991), sequences were amplified using ITS primers, ITS-
99 1F and ITS-4R (White *et al.*, 1990) followed by sequencing and coupling using in silico cap3
100 contig assembly and processed through the BLASTn at NCBI database.

101 **3. RESULTS AND DISCUSSION**

102 **3.1 DISEASE INCIDENCE AND SYMPTOMATOLOGY**

103
104
105 *Alternaria* fruit rot, a latent postharvest disease of guava, was observed in the
106 incubated fruit samples TVG 1, TVG 2, TVG 3, and TSG 1 collected from Vettikkal and
107 Shakthan Market in Thrissur district during March and April 2024 (Table 1). Fruit rot was
108 observed after three to five days of incubation with a disease incidence of 16.6% in Thrissur
109 district. The sample collection sites were delineated within AEU 10 (North Central Laterites)
110 which is characterized by tropical humid monsoon climate and a dry period of around four and
111 half months. The fruit samples were collected during the dry period when diurnal temperature
112 fluctuations were recorded between 23.1 and 40 °C and relative humidity varied from 23 to 95
113 %. The disease developed under incubation at high humid conditions 90-95% and temperature
114 between 30 and 35 °C as light brown to brown irregular lesions, ranging from 0.5 to 3 cm in
115 diameter (Fig.1.). The symptoms on samples TVG 2 and TVG 3 appeared after three days of
116 incubation, and that on TSG 1 and TVG 1 appeared after four and five days of incubation
117 respectively. At the sampling locations, the Percent Disease Incidence (PDI) and Percent
118 Disease Severity (PDS) of *Alternaria* fruit rot recorded were between 33.3-75% and 5-20%
119 respectively. The highest PDI of 75% was observed in fruits collected from Vettikkal, while the
120 highest PDS of 20% was recorded in fruits collected from Shakthan Market. In a survey
121 conducted by Srivastava and Lal (2009) in fruit markets of Allahabad, the incidence of 17.69%
122 *Alternaria* fruit rot in guava was recorded. Jha and Chourasia (2021) also recorded 16.8% fruit
123 decay caused by *Alternaria* during storage under ambient conditions with 18-22 °C and 75-
124 80% RH at Bhagalpur during 2014-2016. Maiti and Lal (2024) reported 11.42% average
125 incidence of *Alternaria* fruit rot in Prayagraj during October 2023 to February 2024. In this
126 study, despite the higher temperature of 30-35 °C, disease development was observed under
127 high humidity conditions. But the elevated temperatures might have reduced the rate of
128 symptom progression. Brown irregular lesions initially developed on the fruit epidermis
129 extended into pulp as the symptoms progressed. Nongmaithem (2014) and Maiti and Lal
130 (2024) observed that *Alternaria* fruit rot in guava, initially caused brown dry or parched spots
131 on the fruit epidermis that subsequently developed into circular to oval or irregular grey-brown
132 lesions that invade the inner pulp at later stages. Similar kinds of symptom development were
133 noticed in this study.




Fig. 1. Symptoms of *Alternaria* fruit rot on guava

Table 1. Samples showing *Alternaria* fruit rot symptoms and days taken for the appearance of symptom, per cent disease incidence, and per cent disease severity


Sl. No.	Samples	Symptoms	Incubation period (DAI)	Percent disease incidence (PDI)	Percent disease severity (PDS)
1.	TVG 1	Numerous small, light brown, irregular lesions with different diameters ranging from 0.3 to 1.5 cm	5	75	10
2.	TVG 2	Irregular, brown lesions with different diameter ranging from 0.5 to 1.5 cm on the skin of the fruit	3	75	5
3.	TVG 3	Brown, irregular, small lesion of 0.5 cm diameter	3	75	5
4.	TSG 1	Brown discoloration with a diameter of 1.5 cm on one side of the fruit	4	33.3	20

141

3.2 ISOLATION OF PATHOGEN AND PATHOGENICITY TESTS

The pathogen was isolated from symptomatic areas of fruit rot samples on potato dextrose agar (PDA) medium. On PDA, the pathogen produced white to grey-coloured aerial mycelia. Seven-day-old pure cultures of the pathogen isolates were used to test pathogenicity. Symptoms started to appear on pinpricked fruits where the mycelial bit inoculum was covered with cotton swab within three to six days of inoculation, but the fruits without pinpricks remained asymptomatic. On these fruits *Alternaria* fruit rot pathogen isolates TVG 1, TVG 2, TVG 3, and TSG 1 resulted in the development of circular to irregular, brown, sunken lesions of 0.6-2 cm diameter on the surface. Notably, the lesion size progression was significantly slower (Fig. 2.). The symptoms were identical to those seen on naturally infected fruits. Previous studies also reported the pathogenicity of *A. alternata* on guava fruits, when inoculated with spore suspensions of 1×10^4 spores/ml (Nongmaithem, 2014) and 1×10^6 spores/ml (Youssef et al., 2015). Among the different pathogen isolates, TVG 3 and TSG 1 produced symptoms within four days after inoculation while TVG 2 took five days and TVG 1 took six days to express symptoms on inoculated fruits. However, the isolate TVG 1 exhibited rapid progression of symptoms. *Alternaria* is generally considered as a weak pathogen that enters plants through wounds or natural openings and remains quiescent in fruits until ripening

159 (Pearson and Hall, 1975). Therefore, the slower symptom expression may be because of the
160 green ripe fruits used for inoculation rather than ripe fruits.
161



162 *DAI – Days after inoculation

163

164 **Fig. 2. Symptoms of *Alternaria* fruit rot of guava upon artificial inoculation at room**
165 **temperature.**

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185


186

187

188


The pathogen isolates initially developed white to grey coloured, submerged to aerial fluffy mycelia which showed uniform and zonal growth at regular intervals with grey or greyish black center and prominent white margins. The mycelia completely turned into greyish black later. The reverse side of the plate shows a cream colour which later turns brown (Fig. 3.). The mycelia covered 9 cm plate after 10-13 days of incubation with an average growth rate of 0.6-0.9 cm per day (Table 2). The isolates TVG 2 and TVG 3 showed higher growth rates, i.e., 0.9 cm per day. The hyphae were brown-coloured, septate with straight or curved, geniculate, pale brown conidiophores bearing a branched chain of conidia. The conidia were brown, obclavate in shape with 3-4 transverse and 1-2 longitudinal septa having an average size of 22.08 to 29.53 × 7.85 to 12.54 μm (Fig. 4.). The pathogen isolates TVG 1, TVG 2, TVG 3 and TSG 1 associated with guava fruit rot were identified as *Alternaria* sp. Similar morphological characteristics were documented by Simmons and Roberts (1993) for *Alternaria* sp. causing black spots in Japanese pear.

Alternaria is well known for exhibiting morphological plasticity under different conditions (Simmons, 1992). Therefore, species level identification of the pathogen isolate *Alternaria* sp. (TSG 1) was done by comparing the amplified ITS sequences (ITS 1F and ITS 4R) of the isolate with that in the NCBI database. The result showed 99.1 per cent similarity and 98 per cent query cover with *Alternaria alternata* with accession number OR687207.1. This confirms the identity of the isolate as *Alternaria alternata*.



189
190
191

Fig. 3. Colony morphology of pathogen isolates on PDA after 10-15 days (a) Front view of the colony (b) Reverse view of the colony



192
193
194
195
196

Fig. 4. Conidia of different *Alternaria* isolates (1000 X)

Table 2. Cultural and morphological characteristics of isolates

Sl. No.	Isolate	Cultural characters	Reverse side of the Petri plate	Dimension of conidia (μm)	Average growth rate (cm per day)
1.	TVG 1	White, submerged mycelia with grey at the center shows diurnal zonation, eventually turned to a greyish brown	Off-white colour with brown center	22.08 to 29.53 × 7.65 to 12.54	0.81

2.	TVG 2	Whitish grey, fluffy mycelia with regular and uniform growth, later colony turned brown.	Cream colour	20.25 to 26.54 × 6.87 to 12.50	0.9
3.	TVG 3	Fluffy, dull whitish-grey mycelium which showed uniform and zonal growth at regular intervals	Cream colour which later turns to brown	20.64 to 29.38 × 7.87 to 11.87	0.9
4.	TSG 1	Fluffy, upraised whitish grey mycelium with zonal growth at regular interval	Cream with the brown colour center	22.29 to 26.12 × 7.87 to 11.24	0.6

197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238

4. CONCLUSION

The prevalence of *Alternaria alternata*, the fungal pathogen responsible for fruit rot of guava, was observed from Thrissur district, one among the three districts surveyed in this study. Temperature and relative humidity appear to have influenced the incidence of *Alternaria* fruit rot. The infection that may have occurred before or after harvest remained dormant in the fruits until incubation under high humidity near saturation. This study confirms previous research that revealed the prevalence of the disease and associated losses at different locations, highlighting the impact of this pathogen on post-harvest life of guava. Given the economic significance of guava, further studies are needed to investigate post-harvest practices that extend shelf life and minimize future losses due to the disease.

ACKNOWLEDGEMENTS

The authors are thankful to Kerala Agricultural University for providing financial assistance to the first author and for providing facilities for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors. all authors read and approved the final manuscript.

REFERENCES

- Rahman, M.A., Ansari, T.H., Meah, M.B., & Yoshida, T. (2003). Prevalence and pathogenicity of guava anthracnose with special emphasis on varietal reaction. *Pakistan Journal of Biological Science*, 6(3), 234-241.
- Amin, M., Malik, A.U., Khan, A.S., & Javed, N. (2011). Potential of fungicides and plant activator for postharvest disease management in mangoes. *International Journal of Agriculture and Biology*, 13(5), 671-676.
- Harteveld, D.O.C., Akisanmi, O.A., & Drenth, A., (2013). Multiple *Alternaria* species groups are associated with leaf blotch and fruit spot diseases of apple in Australia. *Plant Pathology*, 62(2), 289-297.

239 Kwon, J.H., Cheon, M.G., Kim, J.W. & Kwack, Y.B., (2011). Black rot of kiwifruit caused by
240 *Alternaria alternata* in Korea. *The Plant Pathology Journal*, 27(3), 298-298.
241
242 Doster, M.A. & Michailides, T.J., (2007). Fungal decay of first-crop and main-crop figs. *Plant*
243 *Disease*, 91(12), 1657-1662.
244
245 Prusky, D., Shalom, Y., Kobilier, I., Akerman, M., & Fuchs, Y. (2002). The level of quiescent
246 infection of *Alternaria alternata* in mango fruits at harvest determines the postharvest
247 treatment applied for the control of rots during storage. *Postharvest Biology and Technology*,
248 25(3), 339-347.
249
250 Srivastava, R. & Lal, A.A. (2009). Incidence of post-harvest fungal pathogens in guava and
251 banana in Allahabad. *Journal of Horticulture Science*, 4(1), 85-89.
252
253 Nongmaithem, N. (2014). Incidence of post-harvest fungal pathogens associated with guava
254 fruits. *Annals of Plant Protection Sciences*, 22(2), 448-450.
255
256 Fatima, S. (2019). Introduction to major post-harvest diseases of guava. *Journal of Drug*
257 *Delivery and Therapeutics*, 9(4), 591-593.
258
259 Lakshmi, B.K.M., Reddy, P.N., & Prasad, R.D. (2011). Cross-infection potential of
260 *Colletotrichum gloeosporioides* Penz. isolates causing anthracnose in subtropical fruit crops.
261 *Tropical Agricultural Research*, 22(2), 183-193.
262
263 Wheeler, B.E.J. (1969). *An Introduction to Plant Diseases and Fungi*. John Wiley & Sons, New
264 York, 374 p.
265
266 Rangaswamy, G. (1958). *Diseases of Crop Plants in India*. Prentice hall of India Pvt Ltd., New
267 Delhi, 504 p.
268
269 Rocha, J.R.S., Oliveira N.T., & Menezes M. (1998). Comparison of inoculation methods
270 efficiency for evaluation of *Colletotrichum gloeosporoides* isolates pathogenicity on passion
271 fruit (*Passiflora edulis*). *Brazilian Archives of Biology and Technology*, 41(1), 145-153.
272
273 Doyle, J. (1991). DNA protocols for plants. In Hewitt, G.M., Johnston, A.W.B., Young, J.P.W.
274 (eds), *Molecular Techniques in Taxonomy*. Springer, Berlin, Heidelberg, pp. 283-293.
275
276 White, T.J., Bruns, T., Lee, S. and Taylor, J. 1990. Amplification and direct sequencing of
277 fungal ribosomal RNA genes for phylogenetics. In Innis, M.A., Gelfand, D.H., Sninsky, J.J. and
278 White, T.J. (Eds.), *PCR Protocols: A Guide to Methods and Applications*. Academic Press,
279 San Diego, USA, pp. 315-322.
280
281 Jha, G.N. & Chourasia, H.K. (2021). Effect of wrapping and packaging materials on fruit rot of
282 guava (*Psidium guajava* Linn.). *Annals of Plant and Soil Research*, 23(2), 186-191.
283
284 Maiti, S. & Lal, A.A. (2024). Exploration of fungal pathogens associated with postharvest
285 diseases of guava in Prayagraj, India. *Journal of Advances in Biology and Biotechnology*,
286 27(9), 835-844.
287
288 Youssef, K.Y.A., Mustafa, Z.M.M., Mounir, G.A., & Rehab, M.E.A. (2015). Preliminary studies
289 on fungal species associated with guava fruit drop disease and possible management.
290 *Egyptian Journal of Phytopathology*, 43(1-2), 11- 23.
291

- 292 Pearson, R.C. & Hall, D.H. (1975). Factors affecting the occurrence and severity of black mold
293 of ripe tomato fruit caused by *Alternaria alternata*. *Phytopathology*, 65(12), 1352-1359.
294
- 295 Simmons, E. G. & R. G. Roberts. (1993). "*Alternaria* themes and variations (73)." *Mycotaxon*,
296 48, 109-140.
297
- 298 Simmons, E.G. 1992. *Alternaria* taxonomy: current status, viewpoint, challenge. In Chelkowski,
299 J. and Visconti, A. (Eds.), *Alternaria Biology, Plant Diseases and Metabolites*. Elsevier Science
300 Publishers, Amsterdam, pp. 1-35.

UNDER PEER REVIEW