

Original Research Article

Morphological diversity of 23 kola (*Cola nitida*) genotypes from Côte d'Ivoire and Nigeria

Abstract

Twenty-three kola clones from Côte d'Ivoire and Nigeria were assessed for their morphological characters. The objective of this study was to identify potentially promising genotypes for the improvement of kola productivity. The analyses showed that these two populations had contrasted morphological characters, with the exception of nut length. The Côte d'Ivoire genotypes produced large follicles with more nuts (7.21). Those from Nigeria have large leaves, a thick cortex and large but few nuts (5.08). A high correlation was noted among the foliar characters, as well as among the fruiting characters (nuts and follicles). These results provide opportunities for hybridization among these genotypes, both for plant breeding and for QTLs research for certain characters (nut and follicle size, cortex thickness and leaves dimensions).

Keys words: *Cola nitida*, Côte d'Ivoire, genotypes, morphological characters, Nigeria

1 Introduction

Kola, *Cola nitida* (Malvaceae), is a tree native to West Africa [1]. It is a diploecious plant (2n = 40) that can reach 25 m in adulthood [2]. The kola thrives in the humid tropical forest zones of Africa, where rainfall reaches 1,700 mm/year and temperatures range from 23°C to 28°C.

Kola is cultivated for its nuts, which are used in several sectors: in agri-food for the manufacture of energy drinks and wines [3,4], socially for ceremonies such as weddings, baptisms and funerals [5–7], and in the medical field for the manufacture of medicines and the treatment of several diseases [8,9]. Moreover, colaculture contributes significantly to the economic development of producing countries, and to household livelihoods [10–13].

In Côte d'Ivoire, kola is grown by many growers as a pure crop. In addition to these plantations, part of the production comes from spontaneous trees scattered in cocoa or coffee plots [14,15].

Côte d'Ivoire is the world's leading producer and exporter of kola nuts [16]. However, Ivorian colaculture faces major difficulties: aging orchards [14], lack of mastery of technical itineraries [17] and insufficient quality plant material. To guarantee its productivity, Côte d'Ivoire has set up collections of established trees collected in Côte d'Ivoire and Nigeria [18,19].

Analysis of molecular diversity of these genotypes revealed differentiation among the two origins [19]. Thus, morphological characterization of these genotypes, which is the subject of this article, is a step towards setting up a kola breeding program. It enables us to assess the level of variability of morphological characters between genotypes, with a view to extracting those that present characters of interest for genetic improvement.

2 Materials and methods

2.1 Plant material

Morphological characters were assessed in 23 genotypes (13 from Côte d'Ivoire and 10 from Nigeria). For each genotype of the two origins, 3 to 15 trees were considered (Table 1).

Table 1. Origins and numbers of trees used for morphological characters measurements

Genotypes	Origin	Trees numbers
CIV-305	Côte d'Ivoire	5
CIV-306	Côte d'Ivoire	5
CIV-311	Côte d'Ivoire	5
CIV-313	Côte d'Ivoire	5
CIV-314	Côte d'Ivoire	7
CIV-315	Côte d'Ivoire	15
CIV-316	Côte d'Ivoire	5
CIV-318	Côte d'Ivoire	5
CIV-321	Côte d'Ivoire	8
CIV-322	Côte d'Ivoire	5
CIV-323	Côte d'Ivoire	5
CIV-A2	Côte d'Ivoire	5
CIV-A3	Côte d'Ivoire	5
NIG-329	Côte d'Ivoire	5
NIG-330	Nigeria	3
NIG-331	Nigeria	3
NIG-341	Nigeria	5
NIG-342	Nigeria	4
NIG-352	Nigeria	3
NIG-356	Nigeria	7
NIG-364	Nigeria	3
NIG-366	Nigeria	3
NIG-379	Nigeria	3
Total		80

2.2 Methods

2.2.1 Character measurement

Morphological variability of the genotypes was studied using 15 characters inspired by the work of [20] and [21]. Leaves characters measured were: length (LOF), width (LAF), leaf area (SURF = LOF x LAF) and petiole length (LOP). These measurements were recorded on 10 leaves taken at random from each tree.

Follicle characters measured were: length (LOFR), circumference (CIRC), mass (POFR), cortex thickness (ECOR) and number of nuts (NOGR). These measurements were collected on 20 follicles. Follicle length and cortex thickness were measured with a caliper. The circumference, reflecting the size of the follicles, was measured with a tape measure in its median part. Mass was determined on a precision balance shortly after collection.

Regarding nuts, the characters measured were: length (LOGR), width (LAGR), thickness (EPGR), mass (POGR), size (VOGR = LOGR x LAGR x EPGR) and total nut mass (POGR/FR). These measurements were obtained on 30 nuts per genotype. Dimensions

were measured with a caliper. For mass, the nuts were first stripped of their husks and weighed.

2.2.1 Statistical analysis

Means and standard deviations were calculated for each genotype. The two populations were compared with Student's t-test at the 5% level for each character. Genotypes were then compared by analysis of variance to one classification criterion, using the GLM procedure in SAS 9.4 software, with the Newman-Keuls test at the 5% level as the comparison test. Finally, links among pairs of characters were determined using Pearson's correlation test on STATISTICA 7.1 (2005) software.

3 Results

3.1 General characters of the two populations

Genotypes from Côte d'Ivoire were significantly different from those from Nigeria for all characters except nut length ($t = 0.49$; $p = 0.62$). In Côte d'Ivoire origin, follicles were larger (23.47 cm), with the highest mass (346.43 g), the highest number of nuts per follicle (7.21), the thinnest cortices (6.75 mm) and the shortest follicles (13.17 cm). In contrast, the Nigerian population produced smaller follicles (22.89 cm) weighing 302.45 g, with thicker cortex (11.57 mm) and longer follicles (17.78 cm). The number of nuts per follicle was lower (5.08) than the value obtained in genotype native from Côte d'Ivoire.

Regarding nuts, the genotype population originating in Nigeria produced the largest (37.26 cm³) with the highest mass (24.86 g); while those originating in Côte d'Ivoire produced smaller nuts (25.77 cm³) with a lower mass than the former (22.81 g). The total nut mass per follicle was greater in genotypes from Côte d'Ivoire (157.72 g) than in genotypes from Nigeria (120.86 g) (Table 2).

Table 2. Mean values of morphological characters of kola trees from Côte d'Ivoire and Nigeria

Characters	Côte d'Ivoire	Nigeria	Means	p
LoF	19.29 ± 3.76	20.36 ± 4.71	19.65 ± 4.13	0.003
LaF	6.29 ± 1.49	6.79 ± 2.31	6.46 ± 1.82	0.002
Surf	125.67 ± 82	147.99 ± 65	133.24 ± 65.85	< 0.001
LoP	4.4 ± 1.67	5.01 ± 2.58	4.61 ± 2.04	< 0.001
Circ	23.47 ± 2.52	22.89 ± 3.47	23.28 ± 2.89	0.019
PoFr	346.43 ± 124.08	302.45 ± 137.21	331.52 ± 130.24	< 0.001
LoFr	13.17 ± 2.13	17.78 ± 3.39	14.74 ± 3.41	< 0.001
Ecor	6.75 ± 1.85	11.57 ± 2.53	8.39 ± 3.11	< 0.001
NoGr	7.21 ± 2.61	5.08 ± 2.5	6.49 ± 2.76	< 0.001
LoGr	4.02 ± 0.49	4 ± 0.72	4.02 ± 0.58	0.62
LaGr	2.99 ± 0.46	3.14 ± 0.71	3.04 ± 0.56	0.002
EpGr	2.05 ± 0.57	2.69 ± 0.72	2.27 ± 0.69	< 0.001
VoGr	25.77 ± 11.39	37.26 ± 21.59	29.67 ± 16.52	< 0.001
PoGr	22.81 ± 9.09	24.86 ± 13.39	23.51 ± 10.78	0.028
PoGr/Fr	157.72 ± 79.92	120.86 ± 84.97	145.22 ± 83.44	< 0.001

LoF: leaf length, LaF: leaf width, Surf: leaf area, LoP: petiole length PoFr: average follicle mass, LoFr: follicle length, Circ: follicle circumference, Ecor: cortex thickness, NoGr: number of nuts/follicle, LoGr: nut length, LaGr: nut width, EpGr: nut thickness, VoGr: nut volume, PoGr: average nut mass; PoGr/Fr: total nut mass/follicle

3.2 Morphological characteristics of genotypes

3.2.1 Leaf characteristics

Analysis of variance revealed a significant difference among genotypes for leaf characters ($p < 0.001$). Genotypes NIG-331 and NIG-352 had the largest leaves and longest petioles. In contrast, leaves and petioles were less developed in genotypes CIV-313 and NIG-329. All other genotypes showed intermediate values for the same characters (Table 3).

Table 3. Mean leaf characteristics of 23 kola genotypes from Côte d'Ivoire and Nigeria

Genotypes	LOF (cm)	LAF (cm)	SURF (cm ²)	LOP (cm)
CIV-305	20.98 ± 1.53 cde	8.16 ± 0.82 b	172.06 ± 27.33 cd	5.29 ± 1.35 cde
CIV-306	17.24 ± 2.89 gh	5.57 ± 1.13 f	98.92 ± 36.81 g	4.81 ± 1.39 def
CIV-311	18.23 ± 2.37 g	5.75 ± 0.97 ef	106.84 ± 31.69 g	4.15 ± 1.15 efg
CIV-313	14.39 ± 1.36 i	4.31 ± 0.52 g	62.45 ± 12.28 h	2.37 ± 0.84 j
CIV-314	16.82 ± 2.25 gh	5.74 ± 1.24 ef	98.89 ± 34.38 g	3.99 ± 1.49 fghi
CIV-315	18.75 ± 1.61 fg	7.42 ± 0.92 bc	139.59 ± 23.35 ef	3.6 ± 0.84 fghi
CIV-316	17.39 ± 2.08 gh	6.03 ± 0.91 ef	106.49 ± 28.21 g	3.37 ± 1.02 ghij
CIV-318	22.03 ± 2.71 bcd	7.09 ± 1.11 cd	158.42 ± 41.25 de	5.76 ± 2.11 cd
CIV-321	17.93 ± 2.52 g	5.52 ± 0.95 f	100.46 ± 29.21 g	3.48 ± 0.85 ghij
CIV-322	22.46 ± 2.19 bc	7.77 ± 1.17 bc	175.91 ± 39.76 cd	5.97 ± 1.36 cd
CIV-323	19.15 ± 2.64 efg	5.3 ± 0.96 f	103.35 ± 31.07 g	3.61 ± 1.05 fghi
CIV-A2	18.69 ± 2.69 fg	5.45 ± 0.92 f	103.43 ± 30.32 g	4.42 ± 0.77 efg
CIV-A3	26.68 ± 2.44 a	7.75 ± 0.71 bc	206.94 ± 26.04 b	6.39 ± 1.26 c
NIG-329	15.62 ± 1.6 hi	3.54 ± 0.49 h	55.88 ± 13.12 h	2.81 ± 0.81 ij
NIG-330	20.9 ± 2.33 cde	6.68 ± 0.61 de	140.86 ± 27.94 ef	5.65 ± 1.47 cd
NIG-331	26.59 ± 4.41 a	10.34 ± 2.09 a	282.84 ± 105.43 a	8.77 ± 2.87 a
NIG-341	23.68 ± 3.33 b	7.94 ± 1.35 b	189.91 ± 52.19 bc	5.31 ± 1.67 cde
NIG-342	17.05 ± 3.39 gh	6.12 ± 1.09 ef	107.55 ± 39.4 g	3.88 ± 1.48 fghi
NIG-352	26.31 ± 3.53 a	9.92 ± 1.81 a	266.1 ± 81.99 a	7.64 ± 2.51 b
NIG-356	20.37 ± 1.98 def	5.84 ± 0.74 ef	120.06 ± 25.43 fg	5.93 ± 1.53 cd
NIG-364	18.16 ± 1.98 g	5.78 ± 0.87 ef	106.04 ± 25.09 g	2.83 ± 1.14 ij
NIG-366	17.48 ± 2.86 gh	5.94 ± 1.29 ef	105.83 ± 34.84 g	4.32 ± 1.51 efg
NIG-379	17.46 ± 2.56 gh	5.86 ± 1.28 ef	104.89 ± 37.32 g	2.92 ± 1.14 hij
Means	19.65 ± 4.13	6.46 ± 1.82	133.24 ± 65.85	4.61 ± 2.04
p	< 0.001	< 0.001	< 0.001	< 0.001

Comment [U1]: Data in the table should be written in single or double-digit after decimal

Means with the same letter in a column are statistically equal at the 5% level.

LoF: leaf length, LaF: leaf width, Surf: leaf surface, LOP: petiole length

3.2.2 Follicle characteristics

All genotypes showed significant differences in follicle characteristics ($p < 0.001$). The largest follicles with the highest masses were observed in four genotypes: CIV-323 (25.52 cm, 508.23 g), CIV-322 (26.68 cm, 501.74 g), NIG-341 (26.41 cm, 433.07 g), NIG-352 (26.5 cm, 412.42 g). In contrast, the smallest follicles with the lowest masses were observed in genotypes NIG-342 (19.5 cm, 184.61 g), NIG-366 (20.49 cm, 217.99 g) and NIG-379 (20.5 cm, 213.81 g).

Five genotypes (NIG-330, NIG-341, NIG-356, NIG-364 and NIG-379), all of Nigerian origin, presented the longest follicles (18.34 to 19.49 cm). The Nigerian genotype NIG-352 had the thickest cortex (14.9 mm), on average three times that of genotypes CIV-313 and CIV-323. The number of nuts in the follicles was highest in genotype CIV-316 (10.68). On the other hand, 3.4 to 4.05 nuts were counted in genotypes NIG-342, NIG-330 and NIG-379. All other genotypes showed intermediate values for the same characters (Table 4).

Table 4. Mean values for follicle characters in 23 kola genotypes from Côte d'Ivoire and Nigeria.

Genotypes	CIRC (cm)	POFR (cm)	LOFR (cm)	ECOR (cm)	NOGR
CIV-305	24.1 ± 1.24 bcd	363.56 ± 74.08 bc	13.21 ± 1.19 gh	7.41 ± 1.11 fgh	7.32 ± 1.57 cdefg
CIV-306	23.83 ± 1.89 bcde	328.72 ± 72.02 cd	12.23 ± 1.08 hi	5.17 ± 1.03 jk	6.28 ± 1.74 fgh
CIV-311	21.49 ± 2.92 fg	332.7 ± 49.27 cd	14.53 ± 0.81 fg	8.17 ± 0.79 f	8.08 ± 1.87 bcd
CIV-313	25.1 ± 2.63 abc	436.75 ± 134.87 b	14.31 ± 1.65 fg	4.56 ± 1.15 k	7.64 ± 2.44 bcdef
CIV-314	23.19 ± 1.66 cdef	364.71 ± 79.72 bc	16.1 ± 1.27 def	7.02 ± 1.54 fgh	5.48 ± 2.26 ghi
CIV-315	22.64 ± 1.99 def	289.65 ± 89.87 cde	12.36 ± 1.85 hi	7.06 ± 1.67 fgh	6.24 ± 2.04 fgh
CIV-316	21.86 ± 1.42 efg	300.7 ± 45.03 cde	12.95 ± 0.84 gh	9.32 ± 0.82 e	10.68 ± 2.18 a
CIV-318	21.52 ± 1.88 fg	225.4 ± 71.48 ef	11.67 ± 1.64 hi	6.72 ± 1.83 ghi	5.72 ± 2.62 ghi
CIV-321	22.56 ± 2.81 def	220.7 ± 61.09 ef	10.15 ± 1.17 j	5.78 ± 1.30 ij	4.93 ± 2.45 hij
CIV-322	26.68 ± 2.03 a	501.74 ± 139.87 a	14.74 ± 1.68 efg	8.1 ± 1.68 f	8.93 ± 2.09 b
CIV-323	25.52 ± 1.87 ab	508.23 ± 119.55 a	15.03 ± 1.65 ef	4.54 ± 0.71 k	8.93 ± 1.46 b
CIV-A2	22.66 ± 1.7 def	268.49 ± 72.97 de	10.92 ± 1.29 ij	6.26 ± 0.65 hi	5.6 ± 1.52 ghi
CIV-A3	24.03 ± 1.66 bcd	362.7 ± 74.5 bc	13.11 ± 1.04 gh	7.68 ± 1.09 fg	8.24 ± 2.01 bc
NIG-329	22.45 ± 1.92 def	299.91 ± 71.14 cde	15.92 ± 2.59 def	11.45 ± 1.76 cd	6.5 ± 2.21 defg
NIG-330	24.3 ± 3.48 bcd	323.1 ± 135.93 cd	19.49 ± 3.56 a	11.50 ± 2.42 cd	3.55 ± 1.85 j
NIG-331	22.62 ± 2.67 def	298.67 ± 133.39 cde	17.08 ± 3.39 cd	11.35 ± 1.182 d	4.75 ± 2.02 hij
NIG-341	26.41 ± 3.66 a	433.07 ± 150.42 b	18.54 ± 3.47 abc	10.75 ± 3.32 d	5.45 ± 2.35 ghi
NIG-342	19.5 ± 2.53 h	184.61 ± 98.87 f	16.37 ± 2.86 de	13 ± 1.78 b	3.4 ± 1.81 j
NIG-352	26.5 ± 3.05 a	412.42 ± 137.57 b	17.62 ± 3.05 bcd	14.9 ± 1.39 a	6.35 ± 1.87 efgh
NIG-356	23.46 ± 1.28 cdef	379.02 ± 85.51 bc	18.96 ± 1.96 ab	11.6 ± 1.53 cd	7.95 ± 1.95 bcde
NIG-364	22.66 ± 2.88 def	261.87 ± 111.02 def	18.52 ± 4.55 abc	10.95 ± 2.19 cd	7.3 ± 2.71 cd
NIG-366	20.49 ± 1.39 gh	217.99 ± 35.32 ef	17 ± 2.17 cd	8.05 ± 0.51 f	5.1 ± 1.65 hij
NIG-379	20.5 ± 2.92 gh	213.81 ± 119.01 ef	18.34 ± 4.26 abc	12.15 ± 2.28 bc	4.05 ± 2.39 ij
Means	23.27 ± 2.89	331.52 ± 130.24	14.74 ± 3.41	8.38 ± 3.11	6.48 ± 2.76
p	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001

Comment [U2]: Write one or two digits after decimal in the entire data set

Means with the same letter in a column are statistically equal at the 5% level.

Circ: follicle circumference, PoFr: mean follicle mass, LoFr: follicle length, Ecor: cortex thickness, NoGr: number of nuts per follicle

3.2.3 Nut characteristics

Analysis of variance revealed a significant difference among genotypes for nut characteristics ($p < 0.001$). Nut volume (size) ranged from 13.11 to 71.99 cm³. The largest nuts were produced by NIG-330, followed by NIG-341 (63.15 cm³). However, CIV-316 produced the smallest nuts, followed by CIV-A3, CIV-318, CIV-311, NIG-352 and CIV-A2.

The biggest nuts are heavier: NIG-330 (45.32 g) and NIG-341 (40.88 g). In contrast, the smallest nuts are light: genotypes CIV-316 (10.63 g) and NIG-352 (12.09 g). Total nut mass per follicle was highest in CIV-323 (308.99 g), CIV-313 (232.96 g), CIV-322 (232.15 g), NIG-356 (224.8 g) and NIG-341 (215.93 g). On the other hand, NIG-342 (65.67 g), NIG-379 (70.08 g) and NIG-366 (72.7 g) were the lowest (Table 5).

UNDER PEER REVIEW

Table 5. Mean follicle characteristics of 23 kola genotypes from Côte d'Ivoire and Nigeria

Genotypes	LOGR (cm)	LAGR (cm)	EPGR (cm ²)	VOGR (cm)	POGR	POGR/FR
CIV-305	3.77 ± 0.32 defg	2.91 ± 0.27 fghijkl	2.14 ± 0.42 efg	23.87 ± 7.13 fgh	22.56 ± 6.08 efghi	154.58 ± 38.65 cd
CIV-306	4.34 ± 0.23 c	3.11 ± 0.37 defghi	2.38 ± 0.39 cdef	32.16 ± 6.61 def	26.58 ± 7.49 def	154.21 ± 31.36 cd
CIV-311	3.78 ± 0.22 defg	2.77 ± 0.27 hijkl	1.73 ± 0.31 hij	18.21 ± 4.51 hi	17.66 ± 3.66 hijk	139.1 ± 31.27 cde
CIV-313	4.63 ± 0.31 b	3.20 ± 0.42 defg	2.08 ± 0.5 fgh	30.98 ± 9.82 def	31.99 ± 10.55 cd	232.96 ± 83.88 b
CIV-314	4.03 ± 0.47 cd	2.89 ± 0.49 fghijkl	2.05 ± 0.47 fgh	25.01 ± 11.55 fgh	24.68 ± 6.84 efg	129.16 ± 52.47 cdef
CIV-315	4.27 ± 0.38 c	3.22 ± 0.38 defg	2.26 ± 0.32 def	31.37 ± 7.69 def	17.63 ± 5.73 hijk	109.37 ± 47.71 defg
CIV-316	3.53 ± 0.14 fg	2.65 ± 0.24 ijkl	1.41 ± 0.37 j	13.11 ± 2.95 i	10.63 ± 1.41 l	112.59 ± 23.18 defg
CIV-318	3.52 ± 0.4 fg	2.65 ± 0.37 ijkl	1.82 ± 0.49 ghi	17.68 ± 8.51 hi	18.77 ± 10.33 ghij	93.58 ± 45.22 efg
CIV-321	4.06 ± 0.38 cd	3.02 ± 0.65 efghji	2.69 ± 0.51 bc	32.97 ± 10.48 def	21.17 ± 7.86 fghi	88.4 ± 30.11 efg
CIV-322	4.03 ± 0.41 cd	3.16 ± 0.35 defgh	2.46 ± 0.57 cdef	31.62 ± 9.52 def	26.36 ± 7.69 def	232.15 ± 80 b
CIV-323	4.63 ± 0.59 b	3.59 ± 0.5 bc	2.52 ± 0.35 cde	42.09 ± 10.58 c	34.97 ± 7.77 c	308.99 ± 76.48 a
CIV-A2	3.92 ± 0.23 de	2.88 ± 0.3 ghijkl	1.7 ± 0.39 hij	19.17 ± 4.68 hi	22.74 ± 2.7 efghi	127.08 ± 37.44 cdef
CIV-A3	3.85 ± 0.36 def	2.83 ± 0.35 ghijkl	1.53 ± 0.36 ij	16.79 ± 5.43 hi	20.82 ± 3.29 fghi	168.09 ± 41.18 c
NIG-329	3.84 ± 0.21 def	2.79 ± 0.34 hijkl	2.38 ± 0.45 cdef	25.46 ± 5.79 fgh	17.52 ± 2.94 hijk	114.52 ± 42.24 def
NIG-330	4.98 ± 0.37 a	4.17 ± 0.31 a	3.47 ± 0.65 a	71.99 ± 15.73 a	45.32 ± 5.57 a	158.04 ± 77.25 cd
NIG-331	3.78 ± 0.75 defg	3.25 ± 0.57 def	2.75 ± 0.65 bc	36.53 ± 19.44 cde	23.55 ± 12.51 efgh	112.34 ± 68.24 defg
NIG-341	4.83 ± 0.42 ab	3.79 ± 0.51 b	3.42 ± 0.52 a	63.15 ± 15.37 b	40.88 ± 9.54 b	215.93 ± 91.64 b
NIG-342	3.64 ± 0.72 efg	2.83 ± 0.64 ghijkl	2.61 ± 0.53 bcd	28.88 ± 14.31 efg	19.65 ± 9.57 ghij	65.67 ± 40.2 g
NIG-352	3.43 ± 0.44 g	2.49 ± 0.41 l	2.04 ± 0.7 fgh	18.29 ± 8.57 hi	12.09 ± 5.04 kl	78.47 ± 46.34 fg
NIG-356	4.3 ± 0.69 c	3.32 ± 0.59 cde	2.66 ± 0.67 bcd	39.32 ± 16.85 cd	28.2 ± 10.28 de	224.8 ± 100.82 b
NIG-364	4.14 ± 0.44 cd	3.44 ± 0.5 cd	2.94 ± 0.66 b	42.81 ± 14.3 c	30.64 ± 9.47 cd	96.07 ± 51.56 efg
NIG-366	3.46 ± 0.47 g	2.56 ± 0.47 kl	2.23 ± 0.45 def	21.12 ± 9.84 ghi	13.95 ± 6.4 jkl	72.7 ± 43.47 g
NIG-379	3.6 ± 0.33 efg	2.76 ± 0.51 ijkl	2.47 ± 0.48 cdef	25.01 ± 8.95 fgh	16.81 ± 5.29 ijk	70.08 ± 54.51 g
Means	4.02 ± 0.58	3.04 ± 0.57	2.27 ± 0.69	29.66 ± 16.52	23.51 ± 10.78	145.22 ± 83.44
p	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001

Means with the same letter in a column are statistically equal at the 5% level.

LoGr: nut length, LaGr: nut width, EpGr: nut thickness, VoGr: nut volume index, PoGr: average nut mass; PoGr/Fr: total nut mass per follicle.

3.3 Correlations among characteristics

Several pairs of characters are well correlated ($r > 0.8$), as is the case with leaf characters ($0.85 < r < 0.97$). Nut mass and size were positively correlated ($r = 0.89$), as were follicle circumference and mass ($r = 0.89$). These correlations indicate that the largest nuts are the heaviest. Follicle size and mass evolve in the same direction ($r = 0.89$). Furthermore, follicle mass and total nut mass per follicle were positively correlated ($r = 0.86$).

Other pairs of characters were strongly correlated ($0.5 < r < 0.8$). Nut mass was positively correlated with circumference ($r = 0.52$) and follicle mass ($r = 0.54$). This shows that nut mass evolves in the same direction as follicle size and mass. The same applies to the number and total mass of nuts per follicle ($r = 0.56$), and the average and total mass of nuts per follicle ($r = 0.65$). On the other hand, the number of nuts per follicle and nut thickness are inversely proportional ($r = -0.56$): the more nuts there are in the follicle, the thinner the nuts.

Follicle circumference was positively related to total nut mass ($r = 0.7$). This suggests that larger follicles contain more nuts. Correlation among follicle length and thickness ($r = 0.73$), indicates that longer follicles have a thicker cortex. Leaf length was moderately correlated with follicle circumference ($r = 0.44$). However, correlations among follicle circumference and other leaf characteristics were not significant ($p > 0.05$) (Table 6).

Table 6. Matrix correlations among the 23 kola genotypes characters.

	LOF	LAF	SURF	LOP	CIRC	POFR	LOFR	ECOR	NOGR	LOGR	LAGR	EPGR	VOGR	POGR
LAF	0.88*** p < 0.001													
SURF	0.95*** p < 0.001	0.97*** p < 0.001												
LOP	0.89*** p < 0.001	0.85*** p < 0.001	0.9*** p < 0.001											
CIRC	0.44* p = 0.03	0.32* p = 0.14	0.39* p = 0.06	0.35* p = 0.1										
POFR	0.29* p = 0.18	0.17 p = 0.44	0.23* p = 0.28	0.23* p = 0.29	0.89*** p < 0.001									
LOFR	0.15 p = 0.48	0.15 p = 0.49	0.17 p = 0.44	0.17 p = 0.43	0.09 p = 0.66	0.13 p = 0.55								
ECOR	0.32* p = 0.13	0.36* p = 0.09	0.37* p = 0.07	0.32* p = 0.13	-0.11 p = 0.61	-0.19 p = 0.37	0.73** p < 0.001							
NOGR	0.05 p = 0.82	-0.06 p = 0.77	-0.01 p = 0.93	0.001 p = 0.99	0.37* p = 0.08	0.6** p = 0.002	-0.34* p = 0.11	-0.36* p = 0.09						
LOGR	-0.09 p = 0.68	-0.2 p = 0.35	-0.17 p = 0.43	-0.13 p = 0.54	0.54* p = 0.01	0.5* p = 0.02	0.2 p = 0.36	-0.25* p = 0.23	-0.03 p = 0.88					
LAGR	0.1 p = 0.64	0.01 p = 0.95	0.03 p = 0.89	0.04 p = 0.83	0.45* p = 0.03	0.37* p = 0.07	0.38* p = 0.07	-0.02 p = 0.93	-0.18 p = 0.4	0.91*** p < 0.001				
EPGR	0.08 p = 0.71	0.08 p = 0.69	0.08 p = 0.71	0.07 p = 0.72	0.22* p = 0.31	0.06 p = 0.78	0.61** p = 0.002	0.35 p = 0.1	-0.56** p = 0.005	0.61** p = 0.002	0.78** p < 0.001			
VOGR	0.11 p = 0.61	0.05 p = 0.81	0.06 p = 0.78	0.07 p = 0.72	0.38* p = 0.07	0.25* p = 0.24	0.53** p = 0.01	0.17 p = 0.43	-0.36* p = 0.08	0.84*** p < 0.001	0.95*** p < 0.001	0.91*** p < 0.001		
POGR	0.07 p = 0.75	-0.07 p = 0.75	-0.03 p = 0.89	0.03 p = 0.87	0.52** p = 0.01	0.54** p = 0.03	0.35* p = 0.09	-0.12 p = 0.59	-0.18 p = 0.39	0.91*** p < 0.001	0.93*** p < 0.001	0.71** p < 0.001	0.89*** p < 0.001	
POGR /FR	0.08 p = 0.69	-0.11 p = 0.61	-0.04 p = 0.86	0.37* p = 0.86	0.7** p < 0.001	0.86*** p < 0.001	0.05 p = 0.82	-0.4* p = 0.06	0.56** p = 0.006	0.7** p < 0.001	0.57** p = 0.006	0.17 p = 0.44	0.4* p = 0.06	0.65** p = 0.001

Comment [U3]: No need to mention the p-value in the data set

LoF: leaf length, LaF: leaf width, Surf: leaf area, LoP: petiole length, PoFr: mean follicle mass, LoFr: follicle length, Circ: follicle circumference, Ecor : cortex thickness, NoGr: number of nuts per follicle, PoGr: mean nut mass, LoGr: nut length, LaGr: nut width, EpGr: nut thickness, VoGr: nut volume; PoGr/FR: total nut mass per follicle.

*** very strong correlations at 5% level;

** strong correlations at 5% level;

* average correlations at 5% level;

no * weak correlations at 5% level.

4 Discussion

The aim of this study was to analyze the morphological diversity of *C. nitida* genotypes in CNRA's collection. At the end of the analyses, a significant effect of genotype origin on the characters measured was noted. Only nut length showed no significant effect. The superiority of genotypes from Nigeria over those from Côte d'Ivoire was thus demonstrated for leaf size, nut size and mass, and cortex thickness. But for follicle size and mass, genotypes from Côte d'Ivoire showed the highest values.

Genotypes evaluated had contrasting morphological characters. Analyses revealed significant differences between them for these characters. The variations observed suggest that these characters could serve as a genotype distinguishing index [22].

In terms of cortex thickness, genotypes from Nigeria all have a thick cortex. In contrast, genotypes of Ivorian origin have thin cortexes. These variations could be due to the appearance of the cortex. Indeed, [14] indicates that most genotypes of Ivorian origin in the collection have a smooth cortex, as opposed to Nigerian genotypes. The blisters formed on the follicle cortex help to increase its thickness [23]. In fact, for this reason, genotypes with rough follicles have a thicker cortex.

Regarding production parameters, analyses revealed that the largest follicles contain large nuts with high mass. These results concur with those reported by [24]. In this respect, genotypes CIV-313, CIV-322, CIV-323, NIG-330 and NIG-341 could be genotypes of interest. On the same genotypes, [25] obtained different results when the trees were younger. In addition, as mean nut mass is highly heritable in kola [25], the above-mentioned genotypes could be included in a varietal creation and selection program.

Some characters are positively correlated; this suggests that when one character varies, the character correlated with it may vary in the same direction, as these correlations are not genetic correlations. This is the case, for example, between follicle mass and nut mass, and between nut size and nut mass. These results concur with those of [26] on cocoa. Indeed, this author revealed the existence of a strong positive correlation between the mass of the pods and that of the fresh beans, organs analogous to the follicles and nuts of the kola. Furthermore, in *C. nitida*, correlations between nut mass and follicle mass have been demonstrated by [25]. Moreover, follicle mass and average nut mass are characters associated with selection for yield. So, to improve plot yield, it is possible to select directly for the "follicle mass" or "nut mass" character. What's more, the fresh nut production of a plot can be directly estimated from measurements of follicle mass, rather than that of all the nuts harvested.

A negative correlation was observed between thickness and the number of nuts per follicle. This highlights the existence of a trade-off between the two linked characters. Thus, a reduction in nut thickness would favor an increase in the quantity of nuts produced, through resource allocation.

4 Conclusion

This study described and assessed the morphological diversity of kola genotypes in the existing collection in Côte d'Ivoire. The 23 genotypes analyzed showed contrasting characteristics. Genotypes originating in Nigeria are characterized by a thick cortex, large nuts and leaves. In the Côte d'Ivoire genotype, however, the cortex is thinner, and the leaves and nuts smaller. Several pairs of characters were correlated. Large nuts had the highest masses. Similarly, the biggest follicles were the heaviest. In view of these results,

these genotypes could serve as broodstock for the creation of large-nut varieties by introgression of other characters such as cortex thickness and number of nuts per follicle.

References

1. Aké-Assi L (1984) Les surprenantes vertus de la kola. *Histoire Naturelle* 65: 28–33.
2. Adebola PO, Morakinyo JA (2005) Chromosome numbers of four Nigerian species of *Cola* Schott. & Endlicher (Sterculiaceae). *Silvae Genetica* 54: 42–44.
3. Jayeola CO (2001) Preliminary studies on the use of kolanuts (*Cola nitida*) for soft drink production. *Journal of Food Technology in Africa* 6: 1–5.
4. Burdock GA, Carabin IG, Crincoli CM (2009) Safety assessment of kola nut extract as a food ingredient. *Food Chem Toxicol* 47: 1725–1732.
5. Adedokun MO, Soaga JAO, Olawumi AT, et al. (2013) Socio-economic Contribution, Marketing and Utilization of Edible Kolanut (*Cola acuminata* and *Cola nitida*) to Rural Women Livelihood in Abeokuta, Nigeria. *ijmec*.
6. Ouattara B (2013) Vertus et usages de la noix de cola dans les sociétés Ouest Africaines. *Revue Africaine d'Anthropologie, Nyansa-Pô* 1–17.
7. Dah-Nouvlessounon D, Adoukonou-Sagbadja H, Diarrassouba N, et al. (2015) Indigenous knowledge and socio- economic values of three kola species (*Cola nitida*, *Cola acuminata* and *Garcinia kola*) used in southern Benin. *Eur Sci Jour* 11: 206–227.
8. Ekalu A, Habila JD (2020) Phytochemistry, pharmacology and medicinal uses of *Cola* (Malvaceae) family: a review. *Med Chem Res* 29: 2089–2105.
9. Mbembo BM, Inkoto CL, Amogu J-JO, et al. (2022) Mini-review on the phyto-chemistry, pharmacology and toxicology of *Cola nitida* (Vent.) Schott & Endl. (Malvaceae): A medically interesting bio-resource of multiple purposes in Africa. *NRFHH* 2: 335–342.
10. Ndagi I, Babalola FD, Mokuwonye IU, et al. (2012) Potentials and Challenges of Kolanut Production in Niger State, Nigeria. *ISRN Agronomy* 2012: 1–9.
11. Deigna-Mockey V, Kouadio R, Konan Y, et al. (2016) Diagnosis in production and post-harvest processing of nuts of *Cola nitida* (Malvaceae) in Côte d'Ivoire. *JAERI* 9: 1–11. Formatted: Font: Italic
12. Dibi DV (2018) Les Contraintes Des Coopératives Dans Le Commerce De La Noix De La Cola À Bouaké (Côte d'Ivoire). *ESJ* 14: 68.
13. Savi MK, Noumonvi R, Chadaré FJ, et al. (2019) Synergy between traditional knowledge of use and tree population structure for sustainability of *Cola nitida* (Vent.) Schott. & Endl in Benin (West Africa). *Environ Dev Sustain* 21: 1357–1368. Formatted: Font: Italic
14. Ouattara Y (2021) Caractérisation agro-morphologique et moléculaire de colatiers (*Cola nitida* (Vent.) Schott & Endl.) en collection et de plants en pépinière. Formatted: Font: Italic

15. Ouattara Y, Gba KKM, Séry DJ-M, et al. (2023) Manual pollination in kola (*Cola nitida*): A technique for increasing seeds production. *JACR* 11: 42–50. Formatted: Font: Italic
16. FIRCA (2020) Le FIRCA et la filière Cola, Abidjan (Côte d'Ivoire), FIRCA.
17. Séry DJ-M, Bonsson B, Kouadjo CGZ, et al. (2020) Optimisation de la technique de greffage en fente terminale du colatier (*Cola nitida* [Vent.] Schott and Endlicher). *Journal of Applied Biosciences* 155: 15994–16004. Formatted: Font: Italic
18. Sié RS, N'Goran JAK, Montagnon C, et al. (2009) Characterization and evaluation of two genetic groups and value of intergroup hybrids of *Cola nitida* (Vent.) Schott and Endlicher. *Euphytica* 167: 107. Formatted: Font: Italic
19. Ouattara Y, Koffi KG, Séry DJ-M, et al. (2022) Genetic diversity and structure of kola tree (*Cola nitida*) clones germplasm in Côte d'Ivoire using Single Nucleotide Polymorphism markers. *Biodiversitas* 23: 4677–4685. Formatted: Font: Italic
20. Adebola PO, Morakinyo JA (2006) Evaluation of morpho-agronomic variability of wild and cultivated kola Schott et Endl.) in South Western Nigeria. *Genetic Resources and Crop Evolution* 53: 687–694.
21. Dah-Nouvlessounon D, Adoukonou-Sagbadja H, Nafan D, et al. (2016) Morpho-agronomic variability of three kola trees accessions [*Cola nitida* (Vent.) Schott et Endl., *Cola acuminata* (P. Beauv.) Schott et Endl., and *Garcinia kola* Heckel] from Southern Benin. *Genet Resour Crop Evol* 63: 561–579. Formatted: Font: Italic
22. Gbemavo CJ, Gandji K, Gnangle CP, et al. (2015) Variabilité morphologiques et conservation des morphotypes de *Jatropha curcas* Linn. (Euphoraceae) au Benin. *Journal of Agriculture and Environment for International Development* 109. Formatted: Font: Italic
23. Dublin P (1965) Le colatier en République Centrafricaine : culture et amélioration, 2ème partie, Biologie florale. *Café Cacao Thé* 9: 175–192.
24. Egbe EA, Kuchambi IE, Tchoundjeu Z (2013) Phenotypic variation in fruits and nuts of *Cola acuminata* in three populations of the centre region of Cameroon. *Int Res J Plant Sci* 4: 236–247. Formatted: Font: Italic
25. Sié RS (1999) Etude de la diversité et des paramètres génétiques du colatier *Cola nitida* (Vent.) Schott et Endl. de la collection de Côte d'Ivoire. Formatted: Font: Italic
26. Lachenaud P (2007) Fruit trait variability in wild cocoa trees (*Theobroma cacao* L.) from the Camopi and Tanpok basins in French Guiana. *Acta Botanica Gallica* 154: 117–128. Formatted: Font: Not Italic

Comment [U4]: Reference numbers 5, 14 and 16 should be added with volume and page number