

EFFECT OF TILLAGE AND NUTRIENT MANAGEMENT PRACTICES ON GROWTH, DEVELOPMENT AND NUTRIENT CONTENT OF SOYBEAN (*Glycine max* L.) IN THE NORTH WESTERN HIMALAYAS

ABSTRACT

A field experiment was conducted for 2 years (2022 & 2023) at Research Farm of the Department of Agronomy, COA, CSKHPKV, Palampur on silty clay loam soils to study the performance of soybean to varied tillage and nutrient management practices. In terms of tillage practices, significantly better plant growth in terms of plant height, dry matter accumulation, number of nodules and nodule fresh and dry weight per plant, were observed in conventional tillage and zero tillage with residue retention, while zero tillage without residue resulted in the least growth parameters. The days to 50 % flowering and physiological maturity were not significantly influenced by both tillage and nutrient managements. The content and uptake of N, P and K in soybean were also found to be significantly higher in CT during the first year and in ZTR during the second, although, during both the years, CT and ZTR was at par with each other. ZT practices marked the least content and uptake of nutrients irrespective of the years. The nutrient management practices performed alike during both the years, where significantly higher growth parameters and nutrient content and uptake were obtained in organic farming practice, which was followed by RDF and INM, nano-urea and SPNF. Thus, it can be concluded that though both CT and ZTR performs similarly in plant growth and development, however, in the long term sustainability, ZTR is superior to CT. Organic farming practices is a environment friendly approach and soybean being leguminous in nature greatly responds to it, thereby, adoption of ZTR and organic farming in leguminous crops can be a recommended practices for the long term quality and productivity of agriculture.

Comment [I1]: delete

Comment [I2]: number of days

Key words:Conventional tillage,Dry matter accumulation, Nodule count, Zero tillage with residue, organic farming

Introduction

Soybean (*Glycine max* L.) is one of the most important leguminous crops grown worldwide, serving as a significant source of protein and oil (Thrane *et al.*, 2017). In regions such as the Northwestern Himalayas, where the agro-climatic conditions are unique, optimizing soybean cultivation practices is crucial for achieving high yields and sustaining soil health. Among the various agronomic practices, tillage and nutrient management are critical in determining the growth and development soybean plants.

Tillage practices involves the mechanical manipulation of the soil, plays a pivotal role in influencing soil structure, moisture retention, aeration and nutrient availability (Bushari *et al.*, 2015). Different tillage systems, ranging from conventional tillage to reduced or no-till methods, have varying effects on soil health and plant performance. The choice of tillage system can impact the microbial activity, organic matter content and soil compaction, all of which influence nutrient uptake by crops (Hu *et al.*, 2021).

Nutrient management, on the other hand, involves the efficient application of fertilizers and organic amendments to meet the nutritional demands of crops while minimizing environmental impact. The application of macro and micronutrients is essential for the optimum growth and development of soybean, particularly in regions with soil nutrient deficiencies (Bagale, 2021). Fertilizer use efficiency (FUE) and integrated nutrient management (INM) practices can help maintain soil fertility, enhance nutrient availability, and reduce dependence on chemical fertilizers (Panta and Parajulee, 2021).

Comment [I3]: Maximizing fertilizer use efficiency

In India, intensive cereal based cropping systems are the most dominant systems which are exhaustive in nature. Therefore, legume based cropping systems, such as the offers a promising and sustainable alternative (Yan *et al.*, 2024). Soybean, grown during the *kharif*season, is well suited to rainfed conditions and can essentially improve soil fertility through biological nitrogen fixation.

Comment [I4]: as such

In the context of the Northwestern Himalayas, where soil fertility is often low and erratic rainfall patterns pose additional challenges, understanding the interaction between tillage practices and nutrient management is vital. This region's complex agro-ecological characteristics require tailored agricultural practices that balance soil conservation with high productivity. The present study aims to investigate the effect of different tillage practices and nutrient management strategies on the growth, development and nutrient content of soybean with a focus on optimizing these practices for sustainable soybean cultivation in the Northwestern

Comment [I5]: delete

Himalayas. By exploring the influence of tillage and nutrient management on soybean production, this research seeks to contribute valuable insights into best practices for improving both crop yields and soil health in this region.

Material and method

The field experiment was conducted at Research Farm of the Department of Agronomy, COA, CSKHPKV, Palampur for 2 years during *kharif* 2022 and 2023. The experimental site lied in Palam valley of district Kangra, with an elevation of 1290.8 meter above mean sea level, 32°6' N latitude, 76°3' E longitude during *kharif* 2022 and 2023. A critical perusal of weekly meteorological data showed that comparatively more rainfall was received during *kharif* 2023 than *kharif* 2022, with total rainfall during the respective crop seasons were 1864.6 mm and 2350.2 mm, respectively. The mean relative humidity during *kharif* 2022 and *kharif* 2023 ranged between 53 to 91 % and 53 to 92 %, respectively. A critical resume of the meteorological data showed that during *kharif* 2022, maximum and minimum temperatures were comparatively higher than *kharif* 2023. The mean maximum temperature ranged from 23.7 °C to 33.4 °C during *kharif* 2022 and respective range during *kharif* 2023 was 24.3 °C to 29.8 °C. The mean minimum temperature ranged from 10.4 to 20.4 °C and 9.9 to 20 °C during *kharif* 2022 and *kharif* 2023, respectively. The soil of the experimental site was silty clay loam in texture and acidic in reaction. The soil was medium in organic carbon, medium in available nitrogen and available phosphorus and high in available potassium. The experiment was laid out in split plot design with three replications, with fifteen treatments comprising of three tillage practices in the main plot and five nutrient management practices in subplots. The treatments in main plot included conventional tillage (CT), zero tillage with residue retention (ZTR) and zero tillage without residue retention (ZT). The sub plot treatments comprised of Recommended dose of fertilizers (RDF), 50% RDN + 100% PK (basal) + foliar spray of nano-urea @ 2ml l⁻¹ (25 DAS), Integrated Nutrient Management (INM): 50% RDF + 2.5 t ha⁻¹ vermicompost + *Rhizobium* + PSB, Organic farming practice (OFP): 5.0 t ha⁻¹ vermicompost + *Rhizobium* + PSB + 2 sprays of vermish @ 10% at 25 and 45 DAS and Subhash Palekar Natural Farming practice (SPNF). The experimental field was prepared as per treatments related to tillage practices. The plots under zero tillage were cleaned by removing the stones, stubbles and weeds. The plots for conventional tillage were prepared with the help of a power tiller then levelled. Clean and bold seeds were selected and treated with *Rhizobium* and PSB in INM and OFP and with beejamrit @ 1 litre per 10 kg of seeds in case of SPNF before sowing. The Hara soya variety of soybean was used for sowing. The recommended dose of nitrogen, phosphorous and potassium were 20:60:40 kg ha⁻¹. At 15 DAS, in each plot, one meter row length was marked at two places in the net plot area and the mean number of plants was expressed as number of plants per meter row length to record the emergence count. Five plants in second row from the border row from each plot were sampled. The samples were dried in shade and then oven dried till a constant weight was obtained and was used for growth attribute estimation. The developmental studies in soybean included days taken to 50 per cent flowering and days taken to physiological maturity. The date on which 50 percent plants showed flowering in the net plot was recorded and number of days to attain this was calculated from the date of sowing. The date when 95 per cent of pods on a plant turned brown in colour and leaves turned yellow and started shedding, it was considered as physiological maturity stage and the date was noted for number of days taken for maturity. To estimate nitrogen, phosphorous and potassium content and uptake of soybean standard methods given by Jackson 1973 were followed.

Comment [I6]: delete

Comment [I7]: being

Comment [I8]: the same

Comment [I9]: and

Comment [I10]: number of days

The nutrient uptake by crops was computed with the help of following relationship:

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = \frac{\text{Nutrient content (\%)} \times \text{Straw/Grain yield (kg ha}^{-1}\text{)}}{100}$$

Comment [I11]: To avoid confusion, such formula should be written using the 'insert formula' option in MS Word or other such softwares.

Total uptake of particular nutrient was calculated as follows:

Total uptake = uptake by seed + uptake by straw

The data on growth, development and nutrient contents and uptakes were recorded and subjected to statistical analysis. Statistical significance was tested by applying F-test at 0.05 level of probability.

Results and Discussion

Effect of tillage and nutrient management practices on growth parameters of soybean

A close perusal of data in Table 1 revealed that emergence counts were not significantly impacted by various tillage and nutrient management systems during both years of study. However, administration of various

tillage practices significantly influenced the plant height, dry matter accumulation, nodule count per plant, fresh and dry weight of nodule of soybean during both *kharif* 2022 and 2023. During the first *kharif*, higher growth attributes were obtained with conventional tillage, which was followed by zero tillage with residue retention and the lowest with zero tillage without residues. In the following year, the ZTR resulted in higher growth parameters followed by CT, while the shortest were recorded with ZT. However, during both the years, CT and ZTR were at par with each other. Better plant growth with CT during the initial year could be attributed to improved soil aeration, nutrient availability and root growth (Bushari *et al.*, 2015). CT practices break up the soil, enhancing aeration, improving root penetration and allowing easy access to nutrients and water initially (Rodríguez *et al.*, 2022). However, over time, excessive tilling led to degraded soil structure, reduced organic matter, compaction and lower moisture retention (Toth *et al.*, 2024). This might have failed CT in producing the highest growth in the later year. By the second year, ZTR practice resulted in superior plant growth probably due to better nutrient cycling and enhanced soil structure over time. Residue retention acted as mulch, covering the soil and reducing evaporation, thereby conserving soil moisture. The residue decomposed and contributed to higher soil organic matter, enhancing water holding capacity and providing a more stable environment for root growth (Li *et al.*, 2018). ZT consistently resulted in the least growth in both years due to poor moisture retention and soil compaction. Absence of residue cover possibly exposed the soil to direct sunlight, increasing evaporation and reduced soil moisture (Sarker *et al.*, 2022). ZT practices led to soil compaction over time, which restricted root growth, limiting the plant's ability to access water and nutrients deep in the soil, thereby, contributing to lesser growth attributes.

In terms of nutrient management practices, a similar trend was observed during both *kharif* 2022 and 2023, where, organic farming practices (OFP) resulted in significantly higher growth attributes, which was followed by INM, RDF, with both being at par with each other. Next in line was combined application of inorganic fertilisers and nano-urea and the least was observed in natural farming practices. Organic farming practices relied on organic inputs such as vermicompost and vermiwash which are rich in essential nutrients. These nutrients were released slowly and steadily as the organic matter decomposed, providing a continuous supply of nutrients throughout the crop growth stages (Mohite *et al.*, 2024). This also improved soil structure by increasing the organic matter content and encouraging diverse soil microbial activity, which further breaks down organic matter, releasing nutrients in a form that the plants can absorb easily (Lim *et al.*, 2015). INM combines both organic and inorganic nutrient sources, ensuring a balance between immediate nutrient availability from inorganic fertilizers and the slower, sustained release of nutrients from organic sources (Graham *et al.*, 2017). This balanced approach ensured that plants have access to nutrients throughout the growing season, promoting strong and sustained growth (Kumar *et al.*, 2024). RDF and combined application of inorganic fertilizers with nano-urea provided a rapid supply of essential nutrients like nitrogen, phosphorus and potassium, which are crucial for early plant development. However, these nutrients are often more readily available in the early stages, resulting in rapid initial growth. Nano-urea provides highly efficient nitrogen delivery, improving plant growth in the short term, but due to lack of contribution to soil organic matter it fails in the long term improvement of soil structure and fertility (Swify *et al.*, 2024). Natural farming practices relied on natural inputs like jeevamrit, beejamrit, ghanjeevamrit, etc., while these inputs provided some nutrients, their availability is often lower and less consistent than organic or inorganic systems, leading to limited nutrient availability at critical growth stages, leading to shorter plants throughout the growing season (Biswas and Pakhira, 2023).

Comment [I12]: resulting

In general, the growth of soybean was found to be greater in *kharif* 2023 than *kharif* 2022, which might be attributed to higher rainfall and humidity in the next year that ensured adequate water supply, nutrient uptake and efficient photosynthesis, besides reducing water stress. The interaction between the various tillage and nutrient management practices was found to be non-significant.

Comment [I13]: the year 2023

Effect of tillage and nutrient management practices on developmental phases of soybean

A close perusal of the data enshrined in Table 2 revealed that the applied treatments had no significant effect on the number of days taken to attain 50 % flowering and days to physiological maturity. However, during the first year, among the tillage practices, numbers of days to 50 % flowering and physiological maturity were found to be lesser in CT and longer duration was observed in ZT. During *kharif* 2023, numbers of days to 50 % flowering and days to physiological maturity were found to be lesser with ZTR and longer with ZT.

Comment [I14]: attain 50%

Table 1: Effect of tillage and nutrient managements on growth parameters of soybean

Treatments	Emergence count at 15 DAS		Plant height (cm)		Dry matter accumulation (g m ⁻²)		Number of nodules per plant		Fresh weight of nodules per plant (g)		Dry weight of nodules per plant (g)	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
Tillage practices												
C ₁	23.36	23.35	89.3	89.3	308.6	316.4	56.9	55.8	0.91	0.88	0.42	0.42
C ₂	22.94	23.25	87.1	91.8	292.6	332.0	54.6	57.3	0.87	0.94	0.40	0.44
C ₃	22.23	22.33	83.7	84.2	269.1	293.6	50.6	51.9	0.81	0.82	0.37	0.39
SEm±	0.46	0.48	0.8	0.8	5.1	3.5	0.6	0.6	0.01	0.01	0.004	0.005
CD (P =0.05)	NS	NS	3.3	3.3	20.1	13.7	2.4	2.3	0.04	0.04	0.02	0.03
Nutrient management practices												
M ₁	22.78	23.01	87.2	89.5	294.7	316.7	53.3	54.2	0.85	0.87	0.40	0.41
M ₂	22.54	22.73	83.9	85.8	272.2	295.8	52.4	53.5	0.84	0.86	0.39	0.40
M ₃	23.03	23.42	89.2	90.5	305.9	330.8	55.4	56.8	0.89	0.91	0.41	0.43
M ₄	23.53	23.85	92.8	94.1	328.3	351.8	57.7	58.5	0.92	0.94	0.43	0.44
M ₅	22.32	22.60	80.3	82.3	249.6	275.0	51.5	52.2	0.82	0.84	0.38	0.39
SEm±	0.42	0.34	0.9	0.9	6.6	6.2	0.7	0.6	0.01	0.01	0.005	0.004
CD (P =0.05)	NS	NS	2.7	2.8	19.2	18.1	2.1	1.6	0.03	0.03	0.02	0.02

C₁: Conventional tillage, C₂: Zero tillage with Residue, C₃: Zero tillage without residue, M₁: RDF, M₂: 50% RDN + 100% PK + nano-urea @ 2 ml l⁻¹ (25 DAS), M₃: INM-50% RDF + 2.5 t ha⁻¹ VC + *Rhizobium*+ PSB, M₄: OF-5 t ha⁻¹ VC + *Rhizobium* + PSB + vermi wash @ 10% (25, 40 and 55 DAS), M₅: Subhash Palekar's natural farming

In terms of nutrient management practices, organic farming practices recorded earliest days while natural farming practices recorded the maximum days. The number of days required by a crop to achieve a particular phenophase is a varietal character and depends on weather conditions, mainly temperature varying slightly due to the cultivation practices. The interaction between the various tillage methods and nutrient management practices was found to be non-significant.

Comment [I15]: the least number of

Comment [I16]: number of days to achieve the phenophases under observation.

Table 2: Effect of tillage and nutrient managements on developmental stages of soybean

Treatments	Days to 50 per cent flowering		Days to physiological maturity	
	2022	2023	2022	2023
Tillage practices				
C ₁	57.20	60.33	117.81	119.90
C ₂	59.47	59.33	118.50	119.26
C ₃	60.13	61.53	120.47	120.80
SEm±	1.12	1.49	0.76	0.63
CD (P =0.05)	NS	NS	NS	NS
Nutrient management practices				
M ₁	58.56	60.00	118.97	119.99
M ₂	59.44	61.67	119.44	120.16
M ₃	58.33	59.44	118.49	119.53
M ₄	57.56	58.33	117.98	119.36
M ₅	60.78	62.56	119.75	120.78
SEm±	1.63	1.49	0.98	0.63
CD (P =0.05)	NS	NS	NS	NS

C₁: Conventional tillage, C₂: Zero tillage with Residue, C₃: Zero tillage without residue, M₁: RDF, M₂: 50% RDN + 100% PK + nano-urea @ 2 ml l⁻¹ (25 DAS), M₃: INM-50% RDF + 2.5 t ha⁻¹ VC + *Rhizobium*+ PSB, M₄: OF-5 t ha⁻¹ VC + *Rhizobium* + PSB + vermi wash @ 10% (25, 40 and 55 DAS), M₅: Subhash Palekar's natural farming

Effect of tillage and nutrient management practices on N, P and K content in grain and straw of soybean

The data pertaining to the effect of various tillage and nutrient management practices on the nitrogen, phosphorous and potassium content in grain and straw of soybean have been embodied in Table 3. The result revealed the significant influence of the treatments on the nutrient content of grain and straw during both the years of study (2022 and 2023). A close perusal of data revealed that CT and ZTR resulted in significantly higher NPK content over ZT. In the first year, CT practices exhibited higher NPK content in soybean while in the following year, maximum content was obtained with ZTR treatments. However, during both the years, both CT and ZTR were at par with each other. Conversely, ZT exhibited lowest NPK content irrespective of the year. The variation in nutrient content between CT and ZTR could be due to differences in soil nutrient availability and microbial activity over time. In the first year, CT likely provided better mineralization due to soil disturbance, enhancing nutrient release (Vilakazi et al., 2022). However, in the succeeding year, ZTR treatments may have improved soil health, organic matter retention and microbial activity, leading to better nutrient availability (Chandra et al., 2023).

With respect to nutrient management studies, during both the years, significantly higher nutrient content in grain was noticed with OFP. It was closely followed by INM and RDF, where both were at par with each other and were superior over conjoint application of inorganic fertilizers and nano-urea. The treatment comprising natural farming practices showed the lowest content of NPK in grain. Similar trends were observed in the NPK content of straw. The higher nutrient content in grain and straw with OFP could be attributed to the gradual release of nutrients from organic sources, improved soil structure and enhanced microbial activity, promoting root growth and better nutrient uptake (Paramesh et al., 2023). INM and RDF also provided sufficient nutrient, though slightly lower than organic systems, due to a combination of organic and inorganic inputs which might be due to reduced organic matter and nutrient cycling imbalances (Kumari et al., 2023). In contrast, natural farming practices, relying primarily on minimal external inputs, resulted in the lowest content,

Comment [I17]: delete

as the nutrient availability may not have been sufficient to meet the crop's full nutrition demand (Awasthi *et al.*, 2020).

The interactions between different treatments had no significant effect on the NPK content of grain and straw of soybean.

Effect of tillage and nutrient management practices on N, P and K uptake in grain and straw of soybean

The mean values of uptake of NPK by grain and straw along with total uptake in soybean have been presented in Table 4. A close analysis of data revealed that CT practices resulted in higher nutrient uptake in both grain and straw, which was however statistically equivalent to ZTR. The lowest uptake of NPK was recorded with ZT. However, in the succeeding year, higher NPK uptake in both grain and straw was noticed when zero tillage was practiced along with residue retention, and it was statistically at par with CT. ZT resulted in the lowest uptake of nutrients. In terms of nutrient management systems, during both the years, NPK uptake in grain and straw was recorded to be significantly higher with organic farming practices, which was followed by INM, RDF, integrated application of inorganic fertilisers and nano-urea and natural farming practices. However, INM was at par with RDF. The total uptake of nutrients in soybean followed the same trend as that in grain and straw for both tillage and nutrient management practices.

Comment [18]: NPK uptake

The nutrient uptake is a function of both nutrient content and yield, therefore, higher NPK content in grain and straw along with significantly larger grain and straw yields in the aforementioned treatments resulted in significantly higher uptake of NPK by both grain and straw (Dubey *et al.*, 2022). The total nutrient uptake by a crop is the sum of the nutrient uptake by grain as well as straw, the higher uptake by grain and straw was reflected in superior uptake of total NPK in the aforesaid treatments.

Interaction effect of tillage practices and nutrient management practices was found to be non-significant during both the years.

Conclusion

The result of the experiment revealed that both CT and ZTR practices resulted in better growth and development of soybean. However, considering the long term benefits and sustainability of future agriculture, ZTR is more preferable than CT. Among the nutrient managements, organic farming outperformed other practices, providing steady nutrient release, enhanced soil structure and microbial activity. Therefore, ZTR and OFP emerged as the most sustainable practices for long-term soybean cultivation in the region.

Table 3: Effect of tillage and nutrient managements on N, P and K content in per cent (%) of soybean

Comment [I19]: delete

Treatments	N content in grain		N content in straw		P content in grain		P content in straw		K content in grain		K content in straw	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
Tillage practices												
C ₁	6.22	6.28	1.56	1.60	0.472	0.494	0.182	0.192	1.51	1.53	2.29	2.32
C ₂	6.20	6.36	1.54	1.66	0.464	0.507	0.177	0.198	1.50	1.55	2.26	2.37
C ₃	6.07	6.20	1.45	1.51	0.453	0.476	0.160	0.175	1.46	1.49	2.19	2.24
SEm±	0.02	0.02	0.01	0.01	0.002	0.003	0.001	0.002	0.01	0.004	0.02	0.02
CD (P =0.05)	0.08	0.08	0.04	0.03	0.010	0.014	0.006	0.007	0.03	0.02	0.08	0.07
Nutrient management practices												
M ₁	6.19	6.30	1.52	1.60	0.466	0.496	0.175	0.188	1.50	1.53	2.28	2.34
M ₂	6.09	6.21	1.49	1.56	0.446	0.478	0.164	0.182	1.46	1.50	2.20	2.26
M ₃	6.23	6.33	1.54	1.62	0.478	0.507	0.181	0.194	1.51	1.55	2.30	2.36
M ₄	6.32	6.44	1.57	1.66	0.499	0.525	0.153	0.206	1.55	1.58	2.34	2.41
M ₅	5.98	6.12	1.46	1.52	0.426	0.455	0.189	0.172	1.42	1.47	2.11	2.17
SEm±	0.03	0.03	0.01	0.01	0.006	0.006	0.003	0.003	0.01	0.008	0.02	0.01
CD (P =0.05)	0.08	0.07	0.02	0.03	0.017	0.017	0.009	0.009	0.02	0.02	0.07	0.04

C₁: Conventional tillage, C₂: Zero tillage with Residue, C₃: Zero tillage without residue, M₁: RDF, M₂: 50% RDN + 100% PK + nano-urea @ 2 ml l⁻¹ (25 DAS), M₃: INM-50% RDF + 2.5 t ha⁻¹ VC + *Rhizobium*+ PSB, M₄: OF-5 t ha⁻¹ VC + *Rhizobium* + PSB + vermi wash @ 10% (25, 40 and 55 DAS), M₅: Subhash Palekar's natural farming

Table 4: Effect of tillage and nutrient managements on NPK uptake (kg ha⁻¹) of soybean

Treatments	N uptake						P uptake						K uptake					
	Grain		Straw		Total		Grain		Straw		Total		Grain		Straw		Total	
	2022	2023	2022	2023	2023	2022	2022	2023	2022	2023	2023	2022	2022	2023	2022	2023	2023	2022
Tillage practices																		
C ₁	86.0	88.4	33.2	34.5	119.2	122.9	6.6	7.0	3.9	4.1	10.5	11.1	21.0	21.6	48.9	56.2	69.9	71.7
C ₂	80.5	94.2	31.4	37.6	111.9	131.7	6.0	7.5	3.6	4.5	9.7	12.0	19.5	23.0	46.0	59.5	65.5	76.6
C ₃	71.9	79.7	27.3	30.5	99.2	110.2	5.4	6.1	3.0	3.6	8.4	9.7	17.3	19.2	41.1	50.1	58.4	64.4
SEm±	1.4	1.3	0.0	0.5	2.0	1.6	0.1	0.2	0.1	0.03	0.2	0.1	0.4	0.4	1.2	0.9	1.5	1.2
CD (P =0.05)	5.6	5.1	2.4	2.0	7.9	6.2	0.4	0.4	0.4	0.1	0.7	0.4	1.6	1.6	4.7	3.5	5.9	4.6
Nutrient management practices																		
M ₁	80.6	88.5	31.1	34.5	111.7	123.0	6.1	7.0	3.6	4.1	9.7	11.1	19.6	21.5	46.7	50.5	66.3	72.1
M ₂	72.9	80.8	28.4	31.7	101.2	112.5	5.3	6.2	3.1	3.7	8.5	9.9	17.5	19.5	41.7	45.9	59.2	65.4
M ₃	85.3	93.2	32.5	36.4	117.8	129.5	6.6	7.5	3.8	4.4	10.4	11.8	20.7	22.8	48.6	53.1	69.3	75.9
M ₄	93.5	101.7	35.5	39.4	129.0	141.1	7.4	8.3	4.4	4.9	11.7	13.2	23.0	24.9	52.9	57.5	75.8	82.4
M ₅	65.0	73.1	25.7	28.9	90.7	101.9	4.6	5.4	2.7	3.3	7.3	8.7	15.5	17.5	36.9	41.2	52.4	58.7
SEm±	2.3	2.3	0.8	0.7	2.8	2.8	0.2	0.1	0.1	0.1	0.3	0.3	0.6	0.6	1.2	1.2	1.6	1.6
CD (P =0.05)	6.9	6.8	2.3	2.1	8.2	8.2	0.6	0.6	0.3	0.3	0.8	0.9	1.7	1.7	3.5	3.4	4.8	4.5

C₁: Conventional tillage, C₂: Zero tillage with Residue, C₃: Zero tillage without residue, M₁: RDF, M₂: 50% RDN + 100% PK + nano-urea @ 2 ml l⁻¹ (25 DAS), M₃: INM-50% RDF + 2.5 t ha⁻¹ VC + *Rhizobium*+ PSB, M₄: OF-5 t ha⁻¹ VC + *Rhizobium* + PSB + vermi wash @ 10% (25, 40 and 55 DAS), M₅: Subhash Palekar's natural farming

Reference

- Awasthi, N., Upadhyay, R., Singh, A., Kumar, R., & Sharma, G. (2020). Effect of different organic inputs on growth and yield of Soybean (*Glycine max.* L) under mountainous conditions of Himachal Pradesh. *Environment Conservation Journal*, 21(3): 195–199.
- Bagale, S. (2021). Nutrient Management for Soybean Crops (Macronutrients and Micronutrient Management). *International Journal of Agronomy*, 2: 10.
- Biswas, S. & Pakhira, R. (2023). Natural Farming in India: Concept and Current Scenario. *VigyanVarta*, 4(9): 76-79.
- Bushari, M. A., Kukal, S. S., Kaur, A., Bhatt, R. & Dulazi, A.A. (2015). Conservation tillage impacts on soil, crop and the environment. *International Soil and Water Conservation Research*, 3(2): 119-129.
- Chandra, P., Khippal, A. K., Prajapat, K., Barman, A., Singh, G., Rai, A. K., Ahlawat, O. P., Verma, R. P. S., Kumari, K. & Singh, G. (2023). Influence of tillage and residue management practices on productivity, sustainability, and soil biological properties of rice-barley cropping systems in Indo-Gangetic plain of India. *Frontiers in Microbiology*, 14:1130397.
- Dubey, S., Kumar, M., Shukla, V., Singh, G., Singh, D., Shahi, A., Uday, & Rao, D. (2022). Nutrient content and uptake in grain and straw of late-sown wheat as a result of chemical weed control and precision nitrogen management. *Scientist*, 1(2): 189-196.
- Graham, R. F., Wortman, S. E & Pittelkow, C. M. (2017). Comparison of Organic and Integrated Nutrient Management Strategies for Reducing Soil N₂O Emissions. *Sustainability*, 9(4):510.
- Hu, X., Liu, J., Liang, A., Li, L., Yao, Q., Yu, Z., Li, Y., Jin, J., Liu, X & Wang, G. (2021). Conventional and conservation tillage practices affect soil microbial co-occurrence patterns and are associated with crop yields. *Agriculture, Ecosystems & Environment*, 319: 107534.
- Jackson, M. L. (1973). Soil chemical analysis, Prentice Hall Indian Private Limited, New Delhi, pp 179-180.
- Kumar, A., Barkha, A. & Chandel, N. (2024). Organic Farming vs. Integrated Nutrient Management: A Comparative Review of Agricultural Productivity and Sustainability. *International Journal of Plant & Soil Science*, 36: 460-473.
- Li, Z., Zhao, B., Olk, D. C., Jia, Z., Mao, J., Cai, Y. & Zhang, J. (2018). Contributions of residue-C and -N to plant growth and soil organic matter pools under planted and unplanted conditions. *Soil Biology and Biochemistry*, 120: 91-104.
- Lim, S. L., Wu, T. Y., Lim, P. N., & Shak, K. P. (2015). The use of vermicompost in organic farming: overview, effects on soil and economics. *Journal of the science of food and agriculture*, 95(6): 1143–1156.
- Mohite, D. D., Chavan, S. S. & Jadhav, V. S. (2024). Vermicomposting: a holistic approach for sustainable crop production, nutrient-rich bio fertilizer, and environmental restoration. *Discover Sustainability*, 5: 60.
- Panta, S. & Parajulee, D. (2021). Integrated Nutrient Management (INM) in Soil and Sustainable Agriculture. *International Journal of Applied Science & Biotechnology*, 9(3): 160-165.
- Paramesh, V., Kumar, M. R., Rajanna, G. A., Gowda, S., Nath, A.J., Madival, Y., Jinger, D., Bhat, S. & Toraskar, S. (2023). Integrated nutrient management for improving crop yields, soil properties, and reducing greenhouse gas emissions. *Frontiers in Sustainable Food Systems*, 7:1173258.
- Rodríguez, C., Durán-Zuazo, V. H., Rodríguez, S. M., García-Tejero, I. F., Gálvez, R. B. & Cuadros, T. S. (2022). Conservation Agriculture as a Sustainable System for Soil Health: A Review. *Soil Systems*, 6(4): 87.
- Sangam, P. & Dipika, P. (2021). Integrated Nutrient Management (INM) in Soil and Sustainable Agriculture. *International Journal of Applied Sciences and Biotechnology*, 9: 160-165.

Sarker, M. R., Galdos, M. V., Challinor, A. J., Huda, M. S., Chaki, A. K. & Hossain, A. (2022). Conservation tillage and residue management improve soil health and crop productivity—Evidence from a rice-maize cropping system in Bangladesh. *Frontiers in Environment Science*, 10: 969819.

Swify, S., Mazeika, R., Baltrusaitis, J., Drapanauskaitė, D. and Barčauskaitė, K. (2024). Review: Modified Urea Fertilizers and Their Effects on Improving Nitrogen Use Efficiency (NUE). *Sustainability*, 16(1):188.

Thrane, M., Paulsen, P.V., Orcutt, M.W. and Krieger, T.M. (2017). *Soy Protein: Impacts, Production, and Applications* (Eds. Nadathur, S.R., Wanasundara, J.P.D and Scanlin, L.). In: Sustainable Protein Sources, Academic Press, p 23-45.

Toth, M., Stumpp C., Klik, A., Strauss, P., Mehdi-Schulz, B., Liebhard, G. & Strohmeier, S. (2024). Long-term effects of tillage systems on soil health of a silt loam in Lower Austria. *Soil and Tillage Research*, 241: 106120.

Vilakazi, B. S., Zengeni, R. & Mafongoya, P. (2022). The effects of different tillage techniques and N fertilizer rates on nitrogen and phosphorus in dry land agriculture. *Agronomy*, 12(10):2389.

Yan, Z., Chu, J., Nie, J., Qu, X., Sánchez-Rodríguez, A. R., Yang, Y., Pavinato, P. S., Zeng, Z. & Zang, H. (2024). Legume-based crop diversification with optimal nitrogen fertilization benefits subsequent wheat yield and soil quality. *Agriculture, Ecosystems & Environment*, 374: 109171.

UNDER PEER REVIEW