

Investigating the Anti-diabetic and Lipid-regulating Properties of *Eclipta alba* in Alloxan-induced Diabetes

Abstract

Since the dawn of human civilization, humanity has used herbal medicine for medicinal purposes. This research sought to evaluate the antidiabetic effectiveness and lipid profile of *Eclipta alba*. We evaluated the antidiabetic efficacy using the alloxan-induced diabetic model. The group receiving a 1200 mg/kg dosage had statistically significant outcomes for total cholesterol and LDL, with findings of $196.15 \pm 8.91^*$ and $123.77 \pm 6.50^*$, respectively ($p < 0.05$). However, no groups exhibited statistically significant results for HDL and triglyceride levels, despite a reduction in these parameters in the blood after the administration of the extract. The group 6, with a dosage of 1200 mg/kg, exhibited statistically significant results ($p < 0.05$) for SGPT and SGOT, with values of $88.71 \pm 6.23^*$ and $92.82 \pm 7.50^*$, respectively. In the kidney function test, groups 5 and 6 exhibited statistically significant results ($p < 0.05$) at doses of 800 and 1200 mg/kg.

Keywords: *Eclipta alba*, HDL, LDL, Diabetes, Herbal medicine, Triglyceride.

Comment [a1]: Avoid literary language.

Comment [a2]: give actual p value

Comment [a3]: conclusion?

Introduction

Diabetes mellitus (DM) is a metabolic condition characterized by a persistent increase in blood glucose levels (BGL) due to reduced insulin action, secretion, or both in target tissues. It is known that all types of diabetes mellitus are linked to long-lasting microvascular (retinopathy, neuropathy, and nephropathy) and macrovascular (coronary and peripheral artery disorders and stroke) complications. These issues occasionally lead to the detection of organ injury and mortality at a late stage or without sufficient medical supervision [1]. The liver, the body's largest glandular organ, is responsible for the regulation of the majority of physiological activities. The liver receives an individual's entire blood volume numerous times throughout the day. It is essential for the metabolic functions of humans [2]. Rising levels of reactive oxygen species (ROS), which include OH, H₂O₂, and O₂, can happen when someone drinks too much, is addicted to drugs, is exposed to some dangerous chemicals, or has a virus or parasite infection [3]. Hepatocellular injury may result from this. The Centers for Disease Control and Prevention conducted research on 1492 clinicians who provide ambulatory treatment in non-government facilities. The survey revealed that these physicians encounter hyperlipidemia as the second most common chronic disease, with hypertension being the only condition they encounter more frequently [4]. The results of the study indicate that the primary factor contributing to hyperlipidemia is the excessive consumption of high-fat meals [5]. The liver plays a crucial role in the metabolism of commonly prescribed anti-hyperlipidemic medications such as atorvastatin, pravastatin, fluvastatin, simvastatin, lovastatin, and rosuvastatin. As a result, the bioavailability of these medications is low [6]. Statins can transiently inhibit the enzyme 3-hydroxy-3-methylglutaryl-co-A reductase (HMG-CoAR). This enzyme lowers cholesterol levels. This enzyme facilitates the reduction of cholesterol synthesis within the cells. The reason for this is that statins have the capacity to enter hepatocytes and inhibit HMG-CoAR, which is responsible for their pharmacological effects [7]. Statin-associated muscle symptoms (SAMS), also known as muscular problems, are the primary adverse effects that restrict the use of statins. The onset of diabetes mellitus (DM) and complications influencing the central nervous system are two additional potentially detrimental consequences [8]. These synthetic medicines are not only costly, but they also have significant adverse effects, which may result in financial hardships for patients who are required to continue taking them throughout the duration of the therapy [9]. Consequently, it is imperative to create antihyperlipidemic medications that are highly effective

and have minimal adverse effects. The discovery and synthesis of novel therapies are contingent upon the presence of plants [10]. They function as a plentiful and beneficial source of naturally occurring compounds that are suitable for therapeutic purposes. Experts in the field suggest that specific chemical constituents extracted from medicinal plants possess therapeutic properties. Consequently, researchers are perpetually in pursuit of innovative herbal remedies and other plant-based medications that can effectively address a variety of ailments [4]. For centuries, numerous countries worldwide have employed traditional medicines as remedies derived from botanicals, dietary supplements, and alternative medical methods. Traditional medicine has experienced a substantial surge in popularity in recent years, with a significant number of individuals throughout the nation relying on it as their primary form of care [11]. Plants used for medical purposes contain a wide variety of chemical constituents, enabling them to produce a wide range of therapeutic and pharmacological effects. Tanning agents, glycosides, alkaloids, saponins, polysaccharides, essential oils, terpenoids, resins, and plant lipids are among the numerous constituents of these substances [12–14]. Genetically engineered plants ultimately achieve the desired therapeutic outcome by precisely regulating chemical levels. Increasing the production of secondary metabolites, which includes making alkaloids, is one of the many possible uses of reverse genetics [15]. Global advancements in scientific research have facilitated the investigation of the therapeutic properties of plant species [16]. In comparison to synthetic pharmaceuticals, plants are becoming increasingly popular due to their inherent safety, potent pharmacological properties, and cost-effectiveness.

Eclipta alba (L.) Hassk., commonly referred to as false daisy or ink plant, is an annual herb belonging to the Asteraceae family. *E. alba* is a medium-sized, branched plant characterized by small white flowers. Warm, humid climates across Asia, South America, and Africa frequently harbor this plant, with reports of its presence in French rice fields in 1991. Often considered a common weed by farmers, traditional medicine uses the plant in various native countries such as India, China, Thailand, Brazil, Korea, and Ivory Coast [18, 17]. China, India, and Thailand traditionally use *E. alba* to treat skin conditions such as atopic dermatitis, which is associated with inflammatory processes, and vitiligo [19]. People extensively use the plant to promote hair growth and treat hepatitis and jaundice. In Eastern Ivory Coast, pregnant women utilize the plant to promote fetal development and facilitate childbirth [18]. It exhibits antimicrobial, anti-hyperlipidemic, and anti-diabetic properties (21). The plant extract of *E. alba* contains a diverse

array of secondary metabolites, such as saponins, sterols, flavonoids, terpenoids, phenolic acids, thiophenes, polyacetylenes, and coumestans [22].

This study investigates the anti-diabetic effects and lipid profile of an ethanolic extract from *Eclipta alba*.

Materials and methods

Drugs, Chemicals, and Instruments

We obtained the ethanol and alloxan from Sigma Aldrich in Germany. Healthcare Pharmaceutical Limited provided us with a complimentary sample of metformin, a commonly used medication for diabetes. The blood serum analysis kits for various biomarkers were acquired from Plasmatic Laboratory Products Ltd. in the United Kingdom. Alere Inc. glucometer is used in this study. We acquired it from Shahbag in Dhaka, Bangladesh. We assessed the biochemical parameters using the Humalyzer 3000, a semiautomated clinical chemistry analyzer.

Plant Collection and Extract Preparation

Three distinct regions in Bangladesh were utilized to collect *Eclipta alba* plants: North Bengal, a hill-track area, and a lowland area. The next step involved authentication and taxonomic identification. The National Herbarium of Bangladesh maintained the plant specimen in accordance with applicable regulations. The leaves were dried in a shaded area for a duration of seven to ten days, followed by fine grinding. The powdered leaves were agitated for 96 hours in a 70% ethanol solution. Following the soaking process, the extract underwent filtration, and the resultant liquid was collected. The concentration was achieved using a rotary evaporator. The dried extract was collected and stored in the refrigerator for future use.

Experimental Animal Handling

100 male Wistar rats weighing 125-150 grams were obtained from the Pharmacy Department of Jahangirnagar University in Dhaka, Bangladesh. The rats were maintained in a controlled environment at the Institute of Nutrition and Food Science, University of Dhaka, with a 12-hour dark/light cycle and a constant temperature of 25°C. We regularly provided the subjects with a standard pellet meal and clean water. The rats were housed in the facility to acclimatise prior to the commencement of the study. The rat trials adhered to the guidelines set forth by the Institutional Animal Ethics Committee (IEAC). The ethical approval was taken from the Dhaka University, Department of Zoology with the issue no 147/pharm.science.ewu. The researchers

Comment [a4]: Still it is not clear why this plant was chosen for anti diabetic activity? what is the rationale?

Comment [a5]: product ID

Comment [a6]: kit number

Comment [a7]: MakeModel?

Comment [a8]: what is meant by distinct? what was speciality?

Comment [a9]: accession number?

Comment [a10]: Could not understand

care for and managed the animals in accordance with the guidelines set by the Swiss Academy of Medical Sciences (SAMS) and the Swiss Academy of Sciences (SCNAT).

Comment [a11]: add references.

Experimental Guidelines

The tests were conducted in adherence to the ethical principles stated in the 2013 Helsinki Declaration. The study strictly adhered to the "3R" standards, which are fundamental principles in Swiss and global legislation regarding the use of animals in research. The prefix "R" represents the concept of "replacement," encompassing both absolute replacements (such as substituting animal models with computer-generated models) and relative replacements (such as substituting live animals with cell or tissue cultures or vertebrates with invertebrates). For the purpose of conducting thorough research, an animal model was utilised. Rats were chosen as test subjects due to their distinct pancreas and beta cells, which makes them suitable for antidiabetic research. This is in contrast to invertebrates, as mammals are vertebrates. The second "R" represents "reduction," which pertains to techniques that minimise the number of animals needed to gather sufficient data for research purposes or maximise the information obtained from each animal. We selected ten rats for this study based on the sample size estimate determined by the power analysis method. We employed this approach to ensure adherence to the recommended guidelines. Refinement, the third "R," involves reducing the pain and distress experienced by experimental animals. In order to enhance the comfort of the rats during surgery and minimise any discomfort caused by pinching, the tail tips of the rats were gently rubbed with isopropyl alcohol both before and after each blood glucose level measurement. The rats were provided with sufficient nourishment throughout the trial, and they were euthanized painlessly at the conclusion, in accordance with the 2013 amendment to the Guidelines for the Euthanasia of Animals.

Comment [a12]: reference?

Experimental Design

We divided the rats into groups based on their body weight and subsequently tested them for antihyperglycemic action (Table 1). The rodents were categorized into groups according to their body weight, with 10 rats in each group. Table 1 illustrates the alloxan control group, consisting of rats that received only alloxan therapy. N/A indicates the absence of therapeutic treatment in this group.

Table 1: Anti-hyperglycemic Activity Analysis

Group number	Group Status	Treatment specimen	Dose of treatment specimen (mg/kg)	Group Abbreviation
1	Negative Control	Physiological Saline	10 mL/kg	N
2	Alloxan control	Alloxan	150 mg/kg	A
3	Alloxan + Metformin	Alloxan + Metformin	150 mg/kg + 100mg	A + M100
4	Alloxan + <i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract low dose	150 mg/kg + 400 mg/kg	A + EA ₄₀₀
5	Alloxan + <i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract medium dose	150 mg/kg + 800 mg/kg	A + EA ₈₀₀
6	Alloxan + <i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract high dose	150 mg/kg + 1200 mg/kg	A+ EA ₁₂₀₀
7	Metformin	Metformin	100 mg/kg	M
8	<i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract low dose	400 mg/kg	EA ₄₀₀
9	<i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract medium dose	800 mg/kg	EA ₈₀₀
10	<i>Eclipta alba</i>	Alloxan + <i>Eclipta alba</i> extract high dose	1200 mg/kg	EA ₁₂₀₀

Comment [a13]: Give appropriate nales. Eg. NC for Negative control, DC for disease control, PC Positive control etc. Donot use single alphabets

Biological Sample Collection

Blood glucose levels were measured by obtaining samples through puncturing the tip of the rat's tail. Blood was collected from the slaughtered animal immediately after a heart puncture and transferred to a microcentrifuge tube. The supernatant fluid was obtained by centrifuging the collected samples for 5 minutes at 5,000 rpm. The fluid was transferred to a different

microcentrifuge tube for biochemical testing. The kidneys and liver were promptly extracted from the animal's body post-sacrifice and thoroughly rinsed with an ice-cold saline solution for subsequent analysis of kidney and liver function. Rats were categorized into distinct groups based on body weight, followed by tests to assess their antihyperglycemic action (Table 1). Rodents were categorized according to body weight, with each group consisting of 10 rats. The control group in Table 1 consisted exclusively of rats treated with alloxan. This group does not receive therapeutic treatment when not applicable is indicated.

Estimation of Biochemical Parameters

By using a glucometer, the blood glucose level was ascertained. The Humaluzer 3000 was one of many tests administered, along with those for the lipid profile (HDL, LDL, Cholesterol, triglyceride), kidneys (Urea, Creatinine), and liver (SGPT and SGOT). We also tested liver and kidney samples for gluconeogenic and glycolytic enzyme activity

Statistical Analysis

All of our findings (raw data) in terms of numerical parameters were recorded and analyzed on a spreadsheet using the MS Excel application. The gathered data were subjected to descriptive statistics, with the findings reported as mean SD. To evaluate statistical significance, we used the SPSS 16 software's "One-way Anova test" to interpret inter-group heterogeneity in terms of several biological factors. The occurrences are considered statistically significant since the 'p' value was less than 0.05($p < 0.05$).

Results and discussion

Herbal medicine involves the application of medicinal plants for the prevention and treatment of illnesses, encompassing both traditional remedies prevalent in various cultures and the utilization of standardized and titrated herbal extracts. This study examined the antidiabetic effects and lipid profile of the herb *Eclipta alba* in a murine model. Diabetes represents a significant health challenge in the twenty-first century. Diabetes is a significant contributor to mortality, with its macro- and microvascular complications leading to increased disability and substantial healthcare expenditures. The dosages of 800 mg/kg and 1200 mg/kg in groups 5 and 6, respectively, yielded statistically significant findings ($p < 0.05$) regarding antidiabetic efficacy (Figure 1). Multiple studies on plant extracts produced comparable results [22, 23]. The group receiving a dose of 1200 mg/kg demonstrated statistically significant outcomes for total cholesterol and LDL, with values of $196.15 \pm 8.91^*$ and $123.77 \pm 6.50^*$, respectively ($p < 0.05$).

However, no groups exhibited statistically significant outcomes regarding HDL and triglyceride levels, despite a reduction in these parameters in the blood following the administration of the extract. Two studies on plant extracts produced comparable results [24, 25]. The group 6, administered a dose of 1200 mg/kg, exhibited statistically significant results for SGPT and SGOT, with values of $88.71 \pm 6.23^*$ and $92.82 \pm 7.50^*$, respectively ($p < 0.05$) (Table 2). Two studies on plant extracts produced comparable results [26, 27]. Groups 5 and 6 demonstrated statistically significant results ($p < 0.05$) in the kidney function test at doses of 800 mg/kg and 1200 mg/kg. Two studies on plant extract produced comparable results [28, -30].

Comment [a14]: NO mention of histopathology




Figure 1: Antidiabetic activity of different dose of *Eclipta alba*

Table 2: Lipid profile after administration of different dose of *Eclipta alba*

Groups	Total Cholesterol	HDL	LDL	Triglyceride	SGPT	SGOT	Urea	Creatinine
N	123.24± 5.25	93.50±6.18	25.77 ± 2.90	50.53 ± 5.81	42.34± 3.08	41.28±4.10	35.77±3.28	0.84±0.092
A	209.50 ± 9.28	46.24±4.20	134.77± 12.13	110.77±6.83	98.01± 9.75	103.73±12.21	107.21±6.21	2.72±0.75
A+M ₁₀₀	137.70 ± 8.21	70.50±4.28	55.42± 4.87	58.162±4.97	58.72± 8.24	54.28±11.10	54.39±7.12	1.41±0.83
A+EA ₄₀₀	204.23 ± 7.50	48.10±5.73	130.33± 9.82	107.53±9.28	96.28± 6.23	100.25±9.87	103.55±7.50	2.4±0.73*
A+EA ₈₀₀	201.19 ± 8.24	49.70±4.73	127.81± 8.37	104.73±7.08	94.73± 7.28	97.82±8.73	100.29±6.20	1.9±0.084*
A+EA ₁₂₀₀	196.15 ± 8.91*	52.37±3.08	123.77± 6.50*	98.50±6.25	88.71± 6.23*	92.82±7.50*	94.90±5.29*	1.48±0.084*
EA ₄₀₀	120.15 ± 6.25	95.08±5.73	27.10± 3.10	52.28±6.24	44.28± 3.29	42.29±4.10	34.08±2.05	0.77±0.052
EA ₈₀₀	124.19 ± 7.10	91.73±4.88	24.50± 2.30	49.28±5.10	44.70± 4.10	39.73±3.15	37.50±3.90	0.88±0.073
EA ₁₂₀₀	128.17 ± 6.10	94.28±5.77	22.08± 1.90	50.50±4.20	43.93± 3.08	41.40±4.02	38.20±4.01	0.81±0.094
M ₁₀₀	123.5 ± 5.10	93.93±4.70	22.77± 2.80	54.81±5.10	40.73± 3.19	44.50±3.25	37.20±3.20	0.76±0.091

Note: The results were expressed in Mean±SEM (standard mean error) *p< 0.05, **p< 0.01, and ***p< 0.001 were considered as statistically significant. The statistical analysis followed by one-way analysis of variance (Dunnett's test) compared to the control.

Comment [a15]: give actual p value

Conclusion

The ethanol extract of *Eclipta alba* demonstrates significant protective effects against diabetes, hypercholesterolemia, liver damage, and impaired kidney function. The extract exhibited a notable effect on the specified outcomes, indicating its possible therapeutic significance. Further

research is required to isolate and identify the specific active compounds that contribute to its anti-diabetic and lipid-lowering effects. This may yield a better understanding of the mechanisms of action and facilitate the advancement of more effective treatments derived from this plant.

Comment [a16]: discussion need to be precisely written.

References:

1. Yikna BB, Yehualashet AS. Medicinal plant extracts evaluated in vitro and in vivo for antidiabetic activities in Ethiopia: Bases for future clinical trials and related investigations. *Evidence-Based Complementary and Alternative Medicine*. 2021;2021(1):9108499..
2. Himi HZ, Rahman MM, Hasan SA, Cruze LR, Ishraat ST, Chowdhury MM. An Evaluation of Hepato-protective Activity of Ethanolic Extract of *Solanum nigrum* with Varying Doses on CCL4 Induced Hepatic Injured Rat. *Asian Journal of Advanced Research and Reports*. 2024 Mar 16;18(4):75-80.
3. FM SS, Juliana AB, Bornila M, Puja B, Nur-Neasha D, Rafat T. An assessment of hepato-protective activity of *Psidium guajava* fruit extract against hepatic injured rodent model. *Asian Journal of Medical Principles and Clinical Practice*. 2023 Oct 7;6(2):240-5.
4. Baroi JA, Hossian MR, Chowdhury MM, Dolon NN, Maliha F, Rupak MA, Lima NN, Ullah MR, Tahsin R. An assessment of anti-hyperlipidemic potentialities of ethanolic extract of *hemidesmus indicus* in high fat induced rat model. *Asian Journal of Food Research and Nutrition*. 2023 Jul 9;2(4):323-30.
5. Zhang Y, Li X, Yang Q, Zhang C, Song X, Wang W, Jia L, Zhang J. Antioxidation, anti-hyperlipidaemia and hepatoprotection of polysaccharides from *Auricularia auricular* residue. *Chemico-Biological Interactions*. 2021 Jan 5;333:109323.
6. Srinivasa Rao K, Prasad T, Mohanta GP, Manna PK. An Overview of Statins as Hypolipidemic Drugs. *International Journal of Pharmaceutical Sciences and Drug Research*. 2011; 3(3):178-183.

7. Schachter M. Chemical, pharmacokinetic and pharmacodynamic properties of statins: an update. *Fundam Clin Pharmacol*. 2005;19:117-125.
8. Thompson PD, Panza G, Zaleski A, Taylor B. Statin-associated side effects. *Journal of the American College of Cardiology*. 2016;67(20):2395-410.
9. Rupak MA, Chowdhury MM, Shurovi FS, Ferdous J, Tahsin MR, Sarif S, Hasan MM, Chowdhury JA, Kabir S, Chowdhury AA, Aktar F. An Evaluation of Analgesic and Anti-Inflammatory Activity of Ethanolic Extract of *Cynodon Dactylon* on Stressed Rodent Model. *Biomedical Journal of Scientific & Technical Research*. 2022;42(3):33550-7.
10. Islam M, Rupak AH, Nasrin N, Chowdhury MM, Sen P, Foyosal AU, Uddin MJ, Ferdous J, Tahsin MR, Aktar F, Kabir S. An evaluation of potential hepatoprotective properties of *Hylocereus undatus* fruit in experimental rat model. *Biomedical Journal of Scientific & Technical Research*. 2022;43(2):34405-16.
11. Chowdhury MM, Sikder MI, Islam MR, Barua N, Yeasmin S, Eva TA, et al. A review of ethnomedicinal uses, phytochemistry, nutritional values, and pharmacological activities of *Hylocereus polyrhizus*. *J Herbmed Pharmacol*. 2024;13(3):353-365. doi: 10.34172/jhp.2024.49411.
12. Lima NN, Dolon NN, Maliha F, Ullah MR, Humayra F, Chowdhury MM, Rupak MA, Baroi JA, Shohan FS, Tashin R. An Evaluation of Analgesic and Anti-Inflammatory Activity of *Ficus racemosa* in Rat Model.
13. Chowdhury M, Chakma B, Islam A, Sikder I, Sultan RA. Phytochemical investigation and in vitro and in vivo pharmacological activities of methanol extract of whole plant *Argyreiacapitiformis* (Poir.) Ooststr. *Clinical Phytoscience*. 2024 Dec;10(1):1-8.
14. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry*. 2013; 1(6):168-182
15. Lima NN, Dolon NN, Maliha F, Ullah MR, Humayra F, Chowdhury MM, Rupak MA, Baroi JA, Shohan FS, Tashin R. An Evaluation of Analgesic and Anti-Inflammatory Activity of *Ficus racemosa* in Rat Model.

16. Pracheta SS, Sharma V, Paliwal R, Sharma S, Yadav S, Singh L, et al. Chemoprotective activity of hydro-ethanolic extract of *Euphorbia nerrifolia* Linn. Leaves against DENA-induced liver carcinogenesis in mice. *Biol Med.* 2011; 3(2):36–44.
17. Feng L, Zhai YY, Xu J, Yao WF, Cao YD, Cheng FF, Bao BH, Zhang L. A review on traditional uses, phytochemistry and pharmacology of *Ecliptaprostrata* (L.) L. *Journal of ethnopharmacology.* 2019 Dec 5;245:112109.
18. Jahan R, Al-Nahain A, Majumder S, Rahmatullah M. Ethnopharmacological significance of *Eclipta alba* (L.) hassk.(Asteraceae). *International scholarly research notices.* 2014;2014(1):385969.
19. Galli SJ, Tsai M, Piliponsky AM. The development of allergic inflammation. *Nature.* 2008 Jul 24;454(7203):445-54.
20. Datta K, Singh AT, Mukherjee A, Bhat B, Ramesh B, Burman AC. *Eclipta alba* extract with potential for hair growth promoting activity. *Journal of ethnopharmacology.* 2009 Jul 30;124(3):450-6.
21. Mithun NM, Shashidhara S, Vivek Kumar R. *Eclipta alba* (L.) A review on its phytochemical and pharmacological profile. *Pharmacologyonline.* 2011;1(1):345-57.
22. Chung IM, Rajakumar G, Lee JH, Kim SH, Thiruvengadam M. Ethnopharmacological uses, phytochemistry, biological activities, and biotechnological applications of *Ecliptaprostrata*. *Applied microbiology and biotechnology.* 2017 Jul;101:5247-57.
22. Mandal SK, Rahmat S, Sakib K, Mehjabin B, Rahman T, Rasna IJ. An Assessment of Anti-diabetic Effect of *Gymnemasylvestre* in Alloxan-induced Rat Model. *International Research Journal of Gastroenterology and Hepatology.* 2024 Feb 6;7(1):29-36.
23. Ahmed AB, Rao AS, Rao MV. In vitro callus and in vivo leaf extract of *Gymnemasylvestre* stimulate β -cells regeneration and anti-diabetic activity in Wistar rats. *Phytomedicine.* 2010 Nov 1;17(13):1033-9.
24. Bhowmik P, Shohan FM, Baroi JA, Pranto TI, Ullah MR, Rupak MA, Zaman TS, Rasna IJ, Tashin R. Evaluation of the Effects of Ethanolic Extract of *Ficus*

benghalensis on the Lipid Profile and Kidney Function in Rat Model. International Research Journal of Gastroenterology and Hepatology. 2024 Feb 3;7(1):22-8.

25. Kumar P, Rani S, Arunjyothi B, Chakrapani P, Rojarani A. Evaluation of antidiabetic activity of *Gymnemasylvestre* and *Andrographis paniculata* in streptozotocin induced diabetic rats. *Int J PharmacognPhytochem Res.* 2017;9(1):22-5.
26. Kang MH, Lee MS, Choi MK, Min KS, Shibamoto T. Hypoglycemic activity of *Gymnemasylvestre* extracts on oxidative stress and antioxidant status in diabetic rats. *Journal of agricultural and food chemistry.* 2012 Mar 14;60(10):2517-24.
27. El Shafey AA, El-Ezabi MM, Seliem MM, Ouda HH, Ibrahim DS. Effect of *Gymnemasylvestre* R. Br. leaves extract on certain physiological parameters of diabetic rats. *Journal of King Saud University-Science.* 2013 Apr 1;25(2):135-41.
28. Osman M, Fayed SA, Ghada IM, Romeilah RM. Protective effects of chitosan, ascorbic acid and *gymnemasylvestre* against hypercholesterolemia in male rats. *Australian Journal of Basic and Applied Sciences.* 2010;4(1):89-98.
29. Mishra B, Pancholi SS. Investigation of a new antidiabetic combination based on *Gymnemasylvestre* and *Momordica charantia* along with Pioglitazone in major diabetic complications. *Molecular & Clinical Pharmacology.* 2013;4(1):11-25.
30. Kishore L, Singh R. Preventive effect of *gymnemasylvestre* homeopathic preparation on streptozotocin-nicotinamide induced diabetic nephropathy in rats. *Oriental Pharmacy and Experimental Medicine.* 2017 Sep;17(3):223-32.

UNL