

Prevalence and Antibiotic Resistance of Salmonella in Broiler Meat from Rupandehi, Nepal

ABSTRACT

Salmonella is recognized as a significant foodborne pathogen responsible for causing severe infection. It is one of the main causes of huge economic losses due to mortality and decreased production in poultry sector throughout world. This study was conducted to determine the prevalence and antibiogram of *Salmonella* isolated from retail broiler meat. A cross-sectional study was done from August to September 2023 with a total of 152 samples (106 muscle and 46 liver) from different retail meat shops of Siddharthanagar municipality, Rupandehi and transported to Veterinary Medicine Lab, Paklihawa Campus for further analysis according to standard culture-based methods. Antibiogram of isolated *Salmonella* was evaluated against five different groups of antibiotics by disc diffusion method following CLSI guidelines. Data was analysed in SPSS using Chi-Square test at confidence level of 95%. The overall prevalence of Salmonella was recorded to be 18.42% whereby prevalence from liver sample was 19.57% and muscle sample was 17.92%. Statistical analysis showed no significant difference in prevalence among the sample types. The antibiogram study revealed that none of the antibiotics showed 100% effectiveness. The most resistance was seen with Ciprofloxacin (89.28%) followed by Ceftriaxone (85.71%), Tetracycline (82.14%), Chloramphenicol (57.14%), and Amikacin (53.57%). 26 out of 28 isolates (92.86%) were found to be multidrug-resistant (≥ 3 antibiotic groups). The study revealed a higher prevalence of Salmonella in the retail market, highlighting it as a potential public health threat due to the risk of infection from poultry meat consumption. Regular surveillance on antibiotic resistance and justifiable use of antibiotics in the commercial poultry industry is highly recommended.

Keywords: Antibiogram, Antibiotic Resistance, Prevalence, *Salmonella*, Broiler

1. INTRODUCTION

The poultry industry in Nepal has been rapidly expanding in the last decade to meet the increased demand for poultry meat, with an annual growth rate between 17% and 18% and a 261% increase in meat production from 2008 to 2018 (FAO,2020 and Fowler, 2021). In Nepal, poultry

farming is important for both economic growth and food security. However, this growth has been met with many challenges, such as bacterial, viral, and protozoal diseases, and increased antibiotic resistance by bacterial pathogens, which eventually increases the cost of production of poultry products by the farmers (Gompo, 2019). High bacterial load in poultry meat and their antimicrobial resistance have led to potential health issues (Neupane and Kaphle, 2019). Antimicrobial resistance in bacteria, including *Salmonella* spp. is considered a major food safety concern and the spread of resistant strains along the food chain is to be handled for both veterinary medicine and public health (Telli *et al.*, 2022).

Salmonella is a rod-shaped, flagellated, gram-negative facultative anaerobe which infects multiple animal hosts including human by contaminating a wide variety of foods (Wang *et al.*, 2020). There is an increased risk of bacterial infection like *Salmonella* in poultry with increasing number of unorganized poultry farms, which increases the risk of zoonosis and risk of antibiotic resistance due to overuse of antibiotics as growth enhancers (Mahmud *et al.*, 2011). Salmonellosis is a pathogenic bacterial zoonotic disease caused by pathogenic *Salmonella* strains has causing substantial public health impacts. Non-typhoidal *Salmonella* (NTS) is responsible for food-borne diarrheal disease while invasive NTS causes major blood stream infection (Sanni *et al.*, 2023). Contaminated meats, mainly from avian origins are the prospective source of human salmonellosis and become the most therefore the most important source of meat-borne public health hazard (Buncic *et al.*, 2014). *Salmonella enterica* subsp. *enterica* (I) is responsible for major zoonotic diseases.

Tetracyclines, Aminoglycosides and Fluoroquinolones are the most used antibiotic classes among poultry farmers (Subedi *et al.*, 2023). According to Acharya *et al.* (2023) livestock, particularly poultry sectors are supposed to have the highest burden of antimicrobial resistance. Among six subspecies of *Salmonella enterica*, Typhimurium and Enteritidis demonstrated high virulence and resistance to multiple antibiotics (Alegria-Moran *et al.*, 2017). Antibiotic resistance is a significant public health challenge. In the poultry industry, antibiotics are commonly employed both for treating and preventing (prophylactic use) *Salmonella* infections. However, *Salmonella* species have been demonstrated resistance to quinolones, nalidixic acid, and their derivatives, including fluoroquinolones. (Yasmin *et al.*, 2020)

There are a significant number of commercial poultry farms in the Rupandehi district, but very scanty research has been performed, and no adequate baseline information is available till now. The data obtained from present study will help to alert the veterinarians, the public health sector, to strengthen one health approach. Furthermore, According to estimates provided globally, the number of deaths directly linked to AMR has risen to more than 1.2 million in 2019 and this is forecast to increase to approximately 10 million deaths per year by 2050, if insufficient action is taken to control AMR. Thus, this study is intended to provide insights into the prevalence and antibiotic resistance of *Salmonella* in commercial poultry meat of Rupandehi, Nepal.

3. METHODOLOGY

3.1 Study site




Fig1. Map of Nepal Showing the study site.

The current study was conducted in Siddharthanagar municipality of Rupandehi, Nepal. Samples were collected from retail broiler meat shops. The collected samples were transported to the Medicine lab, Institute of Agricultura and Animal Science, Paklihawa Campus, Rupandehi.

3.2 Study population and sample size

Purposive sampling was conducted in retail broiler meat shops of Siddharthanagar municipality. The sample size for the prevalence of *Salmonella* was calculated using Epi-tools epidemiological

calculators by Ausvet. The calculated sample size was with expected precision of 5% and 95% confidence level. Total number of samples collected was 152.

3.5 Sample collection and processing

Collection of samples

For culture, a sample of the liver and muscle of the broiler was taken aseptically in a zip lock bag from different retail meat shops.




Fig 2. Sample collection in Zip-lock bag

Enrichment and selective culturing techniques



Fig.3. Pre-enrichment in Buffer Peptone Water.



Fig.4. *Salmonella* in XLD agar

All samples were treated aseptically. One gram of minced liver and muscle were separately inoculated in a test tube with 10 ml buffer peptone water (BPW) and incubated at 37°C for 20-24 hrs. A loopful of the enriched BPW broth was inoculated into Rappaport Vasiliadis broth and incubated at 37°C for overnight. Selective media XLD was used as the selective media for the isolation of the *Salmonella*. A loopful of culture from Rappaport Vasiliadis (R10 broth) was streaked on the XLD agar and incubated at 37°C for 24 hrs.

Biochemical tests



Fig. 5 MR-VP test
(+)



Fig. 6 Oxidase test (-)




Fig. 7 Catalase test



Fig.8 Gram's stain Gram negative medium sized rod shaped bacterium

To confirm suspected *Salmonella* species, a series of biochemical tests were conducted. The primary tests included Gram staining, catalase, and oxidase tests. Gram staining revealed Gram-negative bacteria, catalase testing showed effervescence with 5% hydrogen peroxide, and oxidase testing indicated a positive result with a purple color change on the oxidase disc. Secondary tests for colonies passing the primary tests included the Simmons Citrate test, which indicated citrate utilization with a blue color change, and the Triple Sugar Iron (TSI) test, assessing fermentation and hydrogen sulfide (H₂S) production through characteristic color changes. The Sulphur Indole Motility (SIM) test evaluated motility, indole production, and H₂S generation, indicated by black precipitate and turbidity. The Methyl Red (MR) test assessed acid production with a red color, indicating a positive result, while the Voges-Proskauer (VP) test indicated butanediol fermentation with no colour change, signifying a negative outcome.

Antibiotic Sensitivity test (AST)






Fig.9 Measurement of zone of inhibition using Vernier calliper scale

Fig.10. Carpet culture in Mueller Hinton Agar using sterile cotton swab

After a biochemical test, 5-6 pure, well-isolated colonies were transferred into a test tube containing 3ml of buffer peptone water from the confirmed culture of Salmonella. Initially, BPW density was measured using a McFarland Densitometer before inoculation and then again after inoculation until the turbidity equivalent to a 0.5 McFarland standard was obtained. If the turbidity is high, extra buffer peptone is added, whereas if the turbidity is low, more colonies are added.



Fig.11 Zone of inhibition *Salmonella* against selected antibiotic discs

A sterile non-toxic cotton swab was dipped into the adjustable suspension and spread over the entire agar surface of the Mueller Hinton Agar (MHA) plate four times, turning the plate at a 60⁰ angle between each streaking. After that antibiotic discs were placed in the inoculated MHA with the help of an antibiotic dispenser. The antibiotics tested, along with their concentrations, were Ceftriaxone (30 mcg), Chloramphenicol (30 mcg), Tetracycline (30 mcg), Ciprofloxacin (5 mcg), and Amikacin (30 mcg). The plates were incubated at 37⁰c for 24 hours after which the zone of inhibition was measured with vernier caliper scale. Using an interpretation chart, according to the zone size of each antimicrobial reporting the organism was interpreted as 'Resistant', 'Intermediate', and 'Sensitive'.

3.6 Statistical analysis

The entry of data and graphical representation was done using Microsoft Office Excel 2019. The statistical association was tested by Chi-square analysis using SPSS version 25 with a significant level defined at $p < 0.05$.

RESULT

The prevalence of *Salmonella* was 19.57% in 46 liver samples and 17.92% in 106 muscle samples. Thus, 28 samples (18.42 %) out of the total 152 samples taken were found positive for *Salmonella* sps.




Fig.12. Overall Prevalence of *Salmonella* in muscle and Liver

Table 1. Statistical association of *Salmonella* with sample type

Sample type	Number of sample (X)	Number of positive (Y)	Prevalence (Y/X)	Odds ratio	<i>p</i> - value	Association
Liver	46	9	19.57%	0.89 (0.37-2.16)	0.82	Statistically non-significant (<i>p</i> >0.05)
Muscle	106	19	17.92%			

The result showed that there is no significant different in prevalence of *Salmonella* with type of sample examined.




Figure 13. Antibiotic resistance pattern with different antibiotics used

The most resistance was seen with Ciprofloxacin (89.28%) followed by Ceftriaxone (85.71%), Tetracycline (82.14%), Chloramphenicol (57.14%), and Amikacin (53.57%) according to our research.

DISCUSSION:

Salmonella is a major zoonotic food borne pathogen causing a variety of clinical manifestations ranging from mild gastroenteritis to bacteriemia and involve many organs with localized extraintestinal infections (Gharieb *et al.*, 2015) Our findings reveal alarming levels of antibiotic resistance among *Salmonella* isolates, with Ciprofloxacin showing the highest resistance (89.28%), followed by Ceftriaxone (85.71%) and Tetracycline (82.14%). These results are consistent with global trends of increasing resistance to these commonly used antibiotics. Chloramphenicol and Amikacin, while showing relatively lower resistance. The prevalence is pretty comparable to the study of Baral *et al.* (2023) and Fowler (2021). Higher prevalence of *salmonella* was observed by Bhandari *et al.* (2013), Lamichanne (2018), Bantawa *et al.* (2018), and Mahato (2019), with prevalence of 46.2%, 25%, 60%, and 40%, respectively. The finding obtained was higher than the research conducted by Dhakal and Manandhar (2005) and Maharjan *et al.* (2006), whose findings were 12% and 14.5%, respectively.

Out of 5 antibiotics tested, none of the antibiotics showed 100% effectiveness. Indiscriminate use of antibiotics intended for rapid cure may trigger the development of resistant strains of bacteria, thus reducing the efficacy of antibiotics. Amikacin showed a maximum sensitivity of 42.86% while none of any isolates showed sensitivity with Ciprofloxacin. A similar result was found in the study of Poudel (2021). High percentage of isolates were resistant towards Tetracycline which is similar to the study of Lamichanne (2018), Ellerbroek *et al.* (2010), and Dhakal *et al.* (2016). 57.14% of *Salmonella* isolates were resistant to Chloramphenicol, which is comparatively lower than 87.50% (R. Selvaraj *et al.*, 2010) but higher than 7.69% (Dhakal *et al.*, 2016).

Previous study by Crump *et al.* (2023) showed 80% of non-typhoidal *Salmonella* isolates to be multidrug resistant. But lower percentage of MDR (45.16%) was found in the study of Adhikari *et al.* (2023). Indiscriminate use of antibiotics for preventive, metaphylactic, therapeutic and as growth promoters might be the main causative factor for increased MDR (Subedi *et al.*, 2023).

CONCLUSION

The prevalence of *Salmonella* in broiler meat from retail meat shops of Siddharthanagar was found to be 18.42%. No significant difference in prevalence among the sample type was recorded. In this study, Amikacin was found to be the most sensitive and Ciprofloxacin was the least sensitive against *Salmonella* among tested antibiotics. Among total *Salmonella* isolates, very high percentage (92.86%) multidrug resistance was seen, thus prudent use of antimicrobials among poultry farmers is necessary. Precautions are required by farmers, retailers as well as consumers to reduce zoonotic risk. These findings reinforce the need for regional surveillance to track resistance trends and inform appropriate policy interventions.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

6. REFERENCES

1. Acharya, K. P., Phuyal, S., & Yon, D. K. (2023). Antibiotic Resistance Among Poultry Farms in South Asia: A Scoping Review. *Nepal Journal of Biotechnology*, *11*(1), 1–15. <https://doi.org/10.54796/njb.v11i1.276>
2. Adhikari, S., Awasthi, P., Bogati, S., & Singh, S. (2023). A Review on An Economic Analysis of Poultry Meat Production in Nepal *Malaysian Animal Husbandry Journal A REVIEW ON AN ECONOMIC ANALYSIS OF POULTRY MEAT PRODUCTION IN. September*, 101–104. <https://doi.org/10.26480/mahj.02.2023.101.103>
3. Adhikari, S., Sharma Regmi, R., Sapkota, S., Khadka, S., Patel, N., Gurung, S., Thapa, D., Bhattarai, P., Sapkota, P., Devkota, R., Ghimire, A., & Rijal, K. R. (2023). Multidrug resistance, biofilm formation and detection of blaCTX-M and blaVIM genes in *E. coli* and *Salmonella* isolates from chutney served at the street-food stalls of Bharatpur, Nepal. *Heliyon*, *9*(5), e15739. <https://doi.org/10.1016/j.heliyon.2023.e15739>

4. Alegria-Moran, R., Rivera, D., Toledo, V., Moreno-Switt, A. I., & Hamilton-West, C. (2017). First detection and characterization of Salmonella spp. In poultry and swine raised in backyard production systems in central Chile. *Epidemiology and Infection*, *145*(15), 3180–3190. <https://doi.org/10.1017/S0950268817002175>
5. Bantawa, K., Rai, K., Subba Limbu, D., & Khanal, H. (2018). Food-borne bacterial pathogens in marketed raw meat of Dharan, eastern Nepal. *BMC Research Notes*, *11*(1), 1–5. <https://doi.org/10.1186/s13104-018-3722-x>
6. Baral, R., Gurung, K., Bhandari, P., Thakuri, P., Gurung, T. W., Sharma, S. P., & Thagunna, B. (2023). *Microbiological Assessment of Broiler Chicken Meat from Different Slaughterhouses of Pokhara Valley* *Malaysian Animal Husbandry Journal* *MICROBIOLOGICAL ASSESSMENT OF BROILER CHICKEN MEAT FROM DIFFERENT*. January. <https://doi.org/10.26480/mahj.02.2023.82.86>
7. Bhandari, N., Nepali, D., & Paudyal, S. (2013). Assessment of bacterial load in broiler chicken meat from the retail meat shops in Chitwan, Nepal. *International Journal of Infection and Microbiology*, *2*(3), 99–104. <https://doi.org/10.3126/ijim.v2i3.8671>
8. Buncic, S., Nychas, G. J., Lee, M. R. F., Koutsoumanis, K., Hébraud, M., Desvaux, M., Chorianopoulos, N., Bolton, D., Blagojevic, B., & Antic, D. (2014). Microbial pathogen control in the beef chain: Recent research advances. *Meat Science*, *97*(3), 288–297. <https://doi.org/10.1016/j.meatsci.2013.04.040>
9. Chakrabarti, A. (2017). *A Textbook of Preventive Veterinary Medicine* (6th ed.). New Delhi: Kalyani Publishers.
10. Coburn, B., Grassl, G. A., & Finlay, B. B. (2007). Salmonella, the host and disease: A brief review. *Immunology and Cell Biology*, *85*(2), 112–118. <https://doi.org/10.1038/sj.icb.7100007>
11. Crump, J. A., Nyirenda, T. S., Kalonji, L. M., Phoba, M. F., Tack, B., Platts-Mills, J. A., Gordon, M. A., & Kariuki, S. M. (2023). Nontyphoidal Salmonella Invasive Disease: Challenges and Solutions. *Open Forum Infectious Diseases*, *10*(Suppl 1), S32–S37. <https://doi.org/10.1093/ofid/ofad020>

12. Dar, M. A., Ahmad, S. M., Bhat, S. A., Ahmed, R., Urwat, U., Mumtaz, P. T., Dar, T. A., Shah, R. A., & Ganai, N. A. (2017). *Salmonella typhimurium* in poultry: a review. *World's Poultry Science Journal*, 73(2), 345–354
13. Dhakal, I.P., and Manandhar, P. (2005). Isolation of *Salmonella* in the pooled samples of litter, food, and water in Chitwan poultries. *Proc. Natl. Expo, 2005*, 43-46.
14. Dhakal, L.B., Ishwari Prasad, D., Saroj Kumar, Y., Md., A., & Md. Zohorul, I. (2016). Prevalence and antibiotic resistance profile of *Salmonella* from livestock and poultry raw meat, Nepal. *International Journal of Molecular Veterinary Research*, January.
15. Dudhane, R. A., Bankar, N. J., Shelke, Y. P., & Badge, A. K. (2023). *The Rise of Non-typhoidal Salmonella Infections in India : Causes , Symptoms , and Prevention*. 15(10). <https://doi.org/10.7759/cureus.46699>
16. Ellerbroek, L., Narapati, D., Phu Tai, N., Poosaran, N., Pinthong, R., Sirimalaisuwan, A., Tshering, P., Fries, R., Zessin, K. H., Baumann, M., & Schroeter, A. (2010). Antibiotic resistance in salmonella isolates from imported chicken carcasses in bhutan and from pig carcasses in Vietnam. *Journal of Food Protection*, 73(2), 376–379. <https://doi.org/10.4315/0362-028X-73.2.376>
17. FAO Sustainable Small-Scale livestock Production. [(accessed on 21 April 2020)]; Available online: <http://www.fao.org/3/ca7660en/CA7660EN.pdf>
18. Fowler, P. D. (2021). *prevalence among poultry farms and slaughterhouses in Chitwan , Nepal*. 14, 437–446.
19. Geetha, M., and Palanivel, K. (2018). A brief review on salmonellosis in poultry. *International Journal of Current Microbiology and Applied Sciences*, 7(05), 1269-1274.
20. Gharieb, R. M., Tartor, Y. H., & Khedr, M. H. E. (2015). Non-Typhoidal Salmonella in poultry meat and diarrhoeic patients: Prevalence, antibiogram, virulotyping, molecular detection and sequencing of class I integrons in multidrug resistant strains. *Gut Pathogens*, 7(1), 1–11.
21. Gompo, T. R., Pokhrel, U., Shah, B. R., & Bhatta, D. D. (2019). Epidemiology of

Important Poultry Diseases in Nepal. *Nepalese Veterinary Journal*, 36, 8–14.
<https://doi.org/10.3126/nvj.v36i0.27746>

22. Hetta, H. F., Ramadan, Y. N., Al-Harbi, A. I., A. Ahmed, E., Battah, B., Abd Allah, N. H., Zanetti, S., & Donadu, M. G. (2023). Nanotechnology as a Promising Approach to Combat Multidrug Resistant Bacteria: A Comprehensive Review and Future Perspectives. *Biomedicines*, 11(2). <https://doi.org/10.3390/biomedicines11020413>
23. Hossain, M. J., Attia, Y., Ballah, F. M., Islam, M. S., Sobur, M. A., Islam, M. A., Ievy, S., Rahman, A., Nishiyama, A., Islam, M. S., Hassan, J., & Rahman, M. T. (2021). Zoonotic Significance and Antimicrobial Resistance in Salmonella in Poultry in Bangladesh for the Period of 2011–2021. *Zoonotic Diseases*, 1(1), 3–24. <https://doi.org/10.3390/zoonoticdis1010002>
24. Kuffmann, F. (1975). Classification of bacteria. Munksgarrd, Copenhagen.
25. Lamichhane R. (2018). *Assessment of bacterial load, isolation and antibiogram of Salmonella from fresh broiler meat in registered meat shop of Bhaktapur municipality, Bhaktapur*. Internship final report submitted to Institute of Agriculture and Animal Science (IAAS), Tribhuvan University, Kathmandu, Nepal.
26. Lynn, M. K. A. G., Opp, C. H. B., Ewitt, W. D., Abney, P. D., Mokhtar, M., & J., F. (1998). Typhimurium Dt104 Infections in the United States. *The New England Journal of Medicine*, 338(19), 1333–1338.
27. Maharjan, M., Joshi, V., Joshi, D. D., & Manandhar, P. (2006). Prevalence of Salmonella species in various raw meat samples of a local market in Kathmandu. *Annals of the New York Academy of Sciences*, 1081, 249–256. <https://doi.org/10.1196/annals.1373.031>
28. Mahmud, Md Showkat, Md Latiful Bari, and M. Anwar Hossain. "Prevalence of Salmonella serovars and antimicrobial resistance profiles in poultry of Savar area, Bangladesh." *Foodborne pathogens and disease* 8.10 (2011): 1111-1118.
29. Mahato, S. (2019). Relationship of Sanitation Parameters with Microbial Diversity and Load in Raw Meat from the Outlets of the Metropolitan City Biratnagar, Nepal. *International Journal of Microbiology*, 2019.

30. MoALD. (2023). Statistical Information on Nepalese Agriculture. *MoALD*, 269. <https://medium.com/@arifwicaksanaa/pengertian-use-case-a7e576e1b6bf>
31. Montville, T. and Matthews, K. (2008). *Food microbiology: an introduction*, 2 ed., ASM Press, Washington, USA.
32. Neupane, R., & Kaphle, K. (2019). Bacteriological quality of poultry meat in Nepal Animal Welfare View project Parasitology View project Bacteriological quality of poultry meat in Nepal. *International Journal of Veterinary Sciences and Animal Husbandry*, 4(5), 10–15. <https://www.researchgate.net/publication/336115104>
33. Poudel R. (2021). *Prevalence and antibiogram of isolated Salmonella from apparently healthy poultry of Morang district*. Internship final report submitted to Institute of Agriculture and Animal Science (IAAS), Tribhuvan University, Kathmandu, Nepal.
34. R. Selvaraj, R. Das, S. Ganguly, M. Ganguli, S. Dhanalakshmi, & Kumar Mukhopadhyay, S. (2010). Characterization and antibiogram of *Salmonella* spp. from poultry specimens. *Journal of Microbiology and Antimicrobials*, 2(9), 123–126. <https://www.researchgate.net/publication/267549402>
35. Sanni, A. O., Onyango, J., Rota, A. F., Mikecz, O., Usman, A., PicaCiamarra, U., & Fasina, F. O. (2023). Underestimated economic and social burdens of non-Typhoidal *Salmonella* infections: The One Health perspective from Nigeria. *One Health*, 16, 100546. <https://doi.org/10.1016/j.onehlt.2023.100546>
36. Subedi, D., Jyoti, S., Thapa, B., Paudel, S., Shrestha, P., Sapkota, D., Bhatt, B. R., Adhikari, H., Poudel, U., Gautam, A., Nepal, R., & Al-Mustapha, A. I. (2023). Knowledge, Attitude, and Practice of Antibiotic Use and Resistance among Poultry Farmers in Nepal. *Antibiotics*, 12(9), 1369. <https://doi.org/10.3390/antibiotics12091369>
37. Telli AE, Biçer Y, Telli N, Güngör C, Turkal G and Onmaz NE, 2022. Pathogenic *Escherichia coli* and *Salmonella* spp. in chicken rinse carcasses: Isolation and genotyping by ERIC-PCR. *Pak Vet J*, 42(4): 493- 498. <http://dx.doi.org/10.29261/pakvetj/2022.049>
38. Vegad, J. L., and Katiyar, A. K. (2015). *A Textbook of Veterinary Special Pathology* (1st ed.). New Delhi: CBS Publishers & Distributors Pvt. Ltd.

39. Wang, M., Qazi, I. H., Wang, L., Zhou, G., & Han, H. (2020). Salmonella virulence and immune escape. *Microorganisms*, 8(3), 1–25.
40. WOA. (2018). *Salmonellosis* (Vol.2).
41. Yasmin S, Nawaz M, Anjum AA, Ashraf K, Ullah N, Mustafa A, Ali MA and Mehmood A, 2020. Antibiotic susceptibility pattern of salmonellae isolated from poultry from different districts of Punjab, Pakistan. *Pak Vet J*, 40(1): 98-102. <http://dx.doi.org/10.29261/pakvetj/2019.080>

UNDER PEER REVIEW