

# Effects of chemical thinning agents on the growth, fruit set, yield, and quality of the nectarine cultivar 'Silver King'

## Abstract

The investigation was conducted at the Research Farm of the Division of Fruit Science, SKUAST-Kashmir, Shalimar campus, during 2016 and 2017 to evaluate the effects of chemical thinning agents on vegetative growth, fruit set, yield, and fruit quality of the nectarine cultivar 'Silver King.' The chemicals tested, were Naphthalene Acetic Acid (NAA), Urea, and Ethephon and were applied after petal fall. The findings revealed that NAA (30 ppm) significantly enhanced vegetative parameters such as plant height, shoot elongation, and leaf area. Both NAA and ethephon effectively reduced fruit set and overall yield while improving fruit quality traits, including fruit size, weight, and coloration. Additionally, ethephon (150 ppm) accelerated fruit maturation by 7–8 days. These results indicate that chemical thinning, particularly with NAA and ethephon, holds potential for optimizing fruit quality and marketability in the 'Silver King' nectarine cultivar.

**Keywords** - Nectarine 'Silver King', thinning agents, chemical thinning, Naphthalene Acetic Acid (NAA), Urea, Ethephon

## Introduction

Nectarines (*Prunus persica* L. Batsch var. *nucipersica*), a relatively recent introduction to the Kashmir Valley, have gained substantial popularity due to their attractive appearance, smooth epidermis, and high market value. However, similar to peach trees, nectarine trees tend to produce an excessive number of flowers, which can lead to overcrowded branches and reduced fruit size if not properly regulated. Traditionally, thinning has been carried out manually by removing surplus blossoms or fruitlets by hand. Although effective, manual thinning is highly labor-intensive, time-consuming, and economically impractical on a commercial scale (Jackson & Looney, 1999). As a result, there has been a transition toward chemical thinning as a more efficient and cost-effective alternative. Chemical thinning agents, including plant growth regulators such as naphthalene acetic acid (NAA) and ethephon, along with other chemicals like urea, thiourea, and ammonium thiosulfate, have become indispensable in contemporary horticultural practices. These agents effectively manage fruit

load while significantly reducing labor requirements and operational costs. These chemical agents are applied at specific stages of fruit development to regulate fruit load and optimize key quality attributes such as size, color, and sugar content. Research has shown that the application of chemical thinners during critical developmental stages, including the closed pink stage, full bloom, and early fruitlet stages, can effectively reduce fruit set, leading to increased individual fruit weight and overall fruit quality (Zilkha *et al.*, 1988; Meitei *et al.*, 2013). For example, urea applied at these stages helps to mitigate excessive fruit production and promotes the development of larger, higher-quality fruits.

Plant growth regulators such as NAA and ethephon, in particular, have demonstrated significant benefits in nectarine production. When applied post-bloom, these agents not only reduce fruit set but also enhance important attributes such as fruit size, coloration, and sugar concentration. Ethephon, for instance, can accelerate fruit maturation, allowing growers to harvest and market their produce earlier, thereby gaining a competitive edge (Rajiv *et al.*, 2017; Rimpika *et al.*, 2017). NAA, specifically, has been shown to stimulate vegetative growth, which in turn supports more efficient fruit development by ensuring a more balanced nutrient distribution to the developing fruits. The use of chemical thinning agents also offers the advantage of reducing physiological stress on the trees, mitigating risks such as limb breakage from excessive fruit load. By controlling fruit numbers early in the development process, trees can allocate more resources to fewer fruits, resulting in higher-quality harvests. This practice not only improves the marketability of the produce by enhancing fruit size and visual appeal but also contributes to the long-term health and productivity of the trees.

## **Material and methods**

The study was carried out in the Research Farm of the Division of Fruit Science at SKUAST-Kashmir, Shalimar campus during the year 2016 and 2017. The experimental orchard is situated at an elevation of 1611 m above mean sea level and lies at 34° 09' N latitude and 74° 52' E longitude. Kashmir has a temperate climate with severe winters from December to March, often dropping below freezing and covered in snow. The valley's altitude ranges from 1500 to 2500 meters above sea level. It experiences a mean maximum temperature of 24.5°C and a minimum of 1.2°C, with a relative humidity of 43.9 per cent. Precipitation averages 650 mm, primarily falling between March and May. The study used four-year-old Nectarine plants of the cultivars Snow Queen, Silver King, and Red Gold, grafted onto peach seedling rootstocks. Planted at a distance of 3 x 3 meter and trained in an open centre system. The plants were

chosen for their uniform vigour and maintained under consistent cultural practices throughout the research.

Experimental plants were pruned each year during December by thinning out all the weak, thin, very vigorous and diseased shoots. Among the remaining healthy shoots, only one year old shoots, well distributed throughout the tree canopy, was retained and headed back to 10-12 nodes. These plants were subjected to growth regulators / chemical treatments and hand thinning as per details given in Table 1.

To evaluate the effects of chemical thinning on *Silver King* nectarines, solutions of Naphthalene Acetic Acid (NAA), urea, and ethephon were prepared by dissolving the specified quantities of each chemical in 100 mL of water, with a few drops of Teepol added to reduce surface tension. Each solution was prepared in 5-liter volumes and applied using a knapsack sprayer to achieve uniform coverage of the developing fruits, ensuring no runoff. Spraying commenced with lower concentrations to prevent excessive dilution, and the sprayer was thoroughly rinsed before applying higher concentrations. The treatments were conducted on clear, calm mornings to minimize environmental variability. Upon fruit maturity, two kilograms were randomly harvested from different sections of the tree for physico-chemical analysis. Any malformed or damaged fruits were excluded from the samples to maintain consistency and accuracy in the assessment.

**Table 1: Treatment details**

S.No.	Treatment	Concentration	Time of application
T <sub>1</sub>	Control	No thinning	The sprays were performed one week after petal fall
T <sub>2</sub>	Naphthalene acetic acid	10 ppm	
T <sub>3</sub>	Naphthalene acetic acid	20 ppm	
T <sub>4</sub>	Naphthalene acetic acid	30 ppm	
T <sub>5</sub>	Ethephon	50 ppm	
T <sub>6</sub>	Ethephon	100 ppm	
T <sub>7</sub>	Ethephon	150 ppm	
T <sub>8</sub>	Urea	0.2 %	
T <sub>9</sub>	Urea	0.4 %	
T <sub>10</sub>	Urea	0.6 %	

Observations recorded during the study was Plant Girth (cm), Trunk Cross Section Area (cm<sup>2</sup>), Plant Height (cm), Leaf Area (cm<sup>2</sup>), Annual Shoot Extension Growth (cm), Date of

Initial Bloom (about 10 % flowering), Date of Full Bloom (above 80 % flowering), Percent Fruit Set (%), Date of Fruit Maturity, Fruit Length (cm), Fruit Breadth (cm), Fruit Weight (g), Fruit Firmness (kg/cm<sup>2</sup>), Fruit Colour (hunter colour lab), Soluble Solids Concentration, SSC (%), Titratable Acidity (%), SSC/Acidity Ratio, Total Sugars (%), Yield (kg/tree), Yield Efficiency (kg/TCSA) Fruit Nutrient Status and Statistical Analysis.

## **Results and Discussion**

The present study investigated the effects of chemical treatments on vegetative growth, fruit set, yield, and fruit quality of nectarine. The application of naphthalene acetic acid (NAA) at 30 ppm significantly enhanced vegetative growth parameters, including annual shoot elongation (52.21 and 52.72 cm), plant height (273.78 and 283.61 cm), trunk girth (23.76 and 26.51 cm), trunk cross-sectional area (44.95 and 55.97 cm<sup>2</sup>), and leaf area (36.54 and 37.36 cm<sup>2</sup>), compared to other treatments during the years 2016 and 2017. Urea, applied at various concentrations, also improved annual shoot elongation (50.67 and 50.82 cm) and leaf area (35.76 and 35.62 cm<sup>2</sup>).

The enhanced vegetative growth observed with NAA treatments may be attributed to improved allocation of photosynthates and nutrients, which facilitated cellular division and increased cell wall elasticity. Moreover, fruit thinning resulted in a reduced crop load, allowing for more vigorous shoot development. These findings align with the results of previous studies by Zilkha *et al.* (1989), Devnath and Kundu (2001), Taghipour *et al.* (2012), and Rimpika *et al.* (2017), which also reported augmented vegetative growth in nectarine following NAA and urea applications.

In contrast, ethephon exhibited a suppressive effect on vegetative growth, as evidenced by its inhibitory influence on leaf expansion and cell enlargement. Similar results were documented by Hamad and Mohammad (1990), who reported that ethrel application reduced vegetative growth in apple trees.

The study also found that different chemical thinners (ethephon, NAA, and urea), when applied one week after petal fall, significantly impacted fruit set. Ethephon at 150 ppm and NAA at 30 ppm resulted in a lower fruit set (53.38, 52.43%, and 54.21, 53.15%) compared to the control (75.54, 72.42%). This may be due to NAA's stimulation of ethylene production, leading to the abscission of young fruits. Ethephon's fruit thinning effect is attributed to the

activation of specific genes that stimulate cell wall-degrading enzymes like EG (endo- $\beta$ -1,4-glucanase) in the separation zone. Urea also exhibited thinning effects, resulting in a lower fruit set compared to the control, corroborating the findings of Zilkha *et al.* (1988). These outcomes are consistent with prior studies by Basak (2006), Taghipour *et al.* (2012), and Meitei *et al.* (2013), which reported that the application of NAA, ethephon, and urea effectively reduced fruit set in peaches and nectarines.

The study revealed that the application of various thinning treatments significantly reduced the average fruit yield per plant, with values of 12.43 kg and 12.12 kg in 2016 and 2017, respectively, compared to the control, which produced 18.14 kg and 19.16 kg per plant. Among the treatments, the highest yield efficiency (0.72 and 0.55 kg cm<sup>-2</sup>) was recorded in the control (T1), followed by T8 (urea at 2%). Conversely, the lowest yield efficiency (0.29 and 0.22 kg cm<sup>-2</sup>) was observed in T4 (NAA at 30 ppm) during both years of the study. The reduction in overall yield observed with thinning treatments is consistent with the findings of Sharma *et al.* (2003), Rimpika *et al.* (2017), and Rajput *et al.* (2017), who reported that fruit thinning with ethephon and NAA resulted in decreased yield but contributed to the production of superior-quality fruits. This reduction in crop load allows for improved resource allocation to the remaining fruits, thereby enhancing their size, weight, and other quality attributes.

The study demonstrated that thinning treatments significantly influenced fruit size, weight, and firmness. Applications of ethephon at 150 ppm and NAA at 30 ppm notably enhanced fruit length, diameter, and weight. This improvement can be attributed to an increased source-to-sink ratio, greater leaf area, and expanded canopy surface area, facilitating more efficient nutrient partitioning toward the remaining fruits after thinning. These findings are in agreement with earlier studies by Vego *et al.* (2010), Taheri *et al.* (2012), and Rimpika *et al.* (2015), which reported that the application of NAA and ethephon improved fruit size and weight in peaches and nectarines. However, fruit firmness was reduced in plants treated with ethephon, likely due to ethylene-induced activation of cell wall-degrading enzymes, such as pectinases and cellulases, leading to a decline in fruit firmness. These observations are consistent with the results of Sharma *et al.* (2003) and Devlal *et al.* (2017), who also reported reduced fruit firmness following ethephon application in stone fruit crops.

Additionally, the study revealed that thinning treatments improved the skin color of the fruits. Ethephon-treated fruits showed better color development, measured by a lower hue angle, possibly due to enhanced ripening and ethylene-induced anthocyanin pigmentation.

These results align with the observations of Whale *et al.* (2012) and Chandel and Singh (2015), who reported improved fruit color in nectarine and peach.

The chemical thinning treatments also reduced the time from fruit set to maturity, with ethephon (100 ppm and 150 ppm) advancing fruit maturity by 7 to 8 days compared to the control. This earlier maturity could be due to increased ethylene production during the final growth phase. These results are in line with those of Sandhu and Singh (2001), who found that ethephon application advanced peach fruit maturity.

Soluble solid concentration, titratable acidity, and sugar content were influenced by the chemical thinning treatments. Ethephon and NAA significantly increased the soluble solids and sugar content while reducing acidity. This may be due to the enhanced translocation of organic metabolites from leaves to fruits, promoting ripening and sugar accumulation. These findings are consistent with previous research by Sharma *et al.* (2001), Taheri *et al.* (2012), and Chandel and Singh (2015). The increase in soluble solids and sugar content can also be attributed to a reduced fruit load, resulting in more carbohydrates being available for the remaining fruits.

The study demonstrated that the application of different chemical treatments had distinct effects on the macronutrient composition of nectarine fruits over the two-year period. The highest nitrogen content (0.70% and 0.72%) was observed in fruits from trees treated with NAA at 30 ppm, followed closely by urea at 0.6 per cent, whereas the lowest nitrogen content (0.61% and 0.60%) was recorded in the untreated control. Phosphorus content exhibited no statistically significant variation among treatments; however, fruits from trees treated with NAA at 30 ppm recorded the highest phosphorus levels (0.051% and 0.054%). Potassium content was significantly influenced by the treatments, with the highest levels (0.83% and 0.85%) detected in fruits from trees treated with NAA at 30 ppm, while the control group exhibited the lowest potassium levels. Calcium content remained unaffected by the treatments, with control plants showing the highest calcium concentrations (0.068% and 0.069%). Similarly, magnesium content was not significantly impacted by the treatments, although the highest magnesium levels (0.029% and 0.030%) were observed in fruits from NAA-treated trees, and the lowest (0.022% and 0.021%) were found in the control group.

**Table 2: Effect of chemical thinning on growth characteristics of nectarine cv. Silver King during 2016 and 2017**

Treatments	Annual shoot extension growth (cm)		Plant height (cm)		Plant girth (cm)		Trunk Cross sectional area (cm <sup>2</sup> )		Leaf area (cm <sup>2</sup> )	
	2016	2017	2016	2017	2016	2017	2016	2017	2016	2017
T <sub>1</sub> Control	44.21	43.74	260.26	268.12	18.36	20.41	26.84	33.16	32.12	31.68
T <sub>2</sub> Naphthalene acetic acid @ 10ppm	45.44	44.62	262.72	270.79	19.15	21.38	29.20	36.41	32.76	32.24
T <sub>3</sub> Naphthalene acetic acid @ 20ppm	50.14	50.27	271.87	281.68	21.81	24.05	37.87	46.05	35.15	34.51
T <sub>4</sub> Naphthalene acetic acid @ 30ppm	52.21	52.72	273.78	283.61	23.76	26.51	44.95	55.97	36.54	37.36
T <sub>5</sub> Ethephon @ 50ppm	46.79	45.81	265.24	272.60	19.77	21.82	31.11	37.91	33.54	33.34
T <sub>6</sub> Ethephon @ 100ppm	47.61	46.33	266.64	274.45	20.12	22.46	32.23	40.20	33.71	33.66
T <sub>7</sub> Ethephon @ 150ppm	48.17	47.05	267.21	276.67	20.44	22.83	33.30	41.53	33.83	33.78
T <sub>8</sub> Urea @ 0.2%	45.27	44.23	262.37	270.42	18.81	20.84	28.18	34.58	32.44	32.14
T <sub>9</sub> Urea @ 0.4%	49.15	49.26	268.15	278.13	20.77	23.18	34.37	42.78	34.39	34.28
T <sub>10</sub> Urea @ 0.6%	50.67	50.82	270.84	280.63	21.46	24.32	36.67	47.09	35.76	35.62
<b>CD<sub>(p≤0.05)</sub></b>	<b>1.29</b>	<b>1.15</b>	<b>1.72</b>	<b>1.61</b>	<b>0.54</b>	<b>0.61</b>	<b>1.76</b>	<b>2.18</b>	<b>0.66</b>	<b>0.61</b>

**Table 3: Effect of chemical thinning on days to flowering characteristics of nectarine cv. Silver King during 2016 and 2017**

Treatments		*Date of initial bloom (About 10%)		*Date of full bloom (Above 80%)	
		2016	2017	2016	2017
T <sub>1</sub>	Control	17.00	24.33	23.33	33.00
T <sub>2</sub>	Naphthalene acetic acid @ 10ppm	16.00	23.33	22.00	33.33
T <sub>3</sub>	Naphthalene acetic acid @ 20ppm	16.67	23.67	22.67	33.33
T <sub>4</sub>	Naphthalene acetic acid @ 30ppm	16.00	24.67	24.00	34.00
T <sub>5</sub>	Ethephon @ 50ppm	15.00	24.00	21.67	33.33
T <sub>6</sub>	Ethephon @ 100ppm	15.33	22.67	22.67	33.67
T <sub>7</sub>	Ethephon @ 150ppm	15.33	23.67	23.00	33.33
T <sub>8</sub>	Urea @ 0.2%	15.67	23.67	22.33	34.00
T <sub>9</sub>	Urea @ 0.4%	15.00	23.00	23.33	33.67
T <sub>10</sub>	Urea @ 0.6%	15.33	23.67	22.33	33.33
<b>CD<sub>(p≤0.05)</sub></b>		<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>

**\*Reference Date: 1<sup>st</sup> Week of March**

**Table 4: Effect of chemical thinning on fruit set (%), fruit maturity and yield characteristics of nectarine cv. Silver King during 2016 and 2017**

Treatments	Initial fruit set (%)		Fruit maturity (DAFB to Harvesting)		Fruit yield (kg tree <sup>-1</sup> )		Yield efficiency (Kg cm <sup>-2</sup> )	
	2016	2017	2016	2017	2016	2017	2016	2017
T <sub>1</sub> Control	75.54	72.42	85.33	87.33	19.16	18.14	0.72	0.55
T <sub>2</sub> Naphthalene acetic acid @ 10ppm	72.23	69.37	84.00	86.33	17.48	16.72	0.60	0.46
T <sub>3</sub> Naphthalene acetic acid @ 20ppm	62.75	60.06	82.00	83.67	15.03	14.25	0.40	0.31
T <sub>4</sub> Naphthalene acetic acid @ 30ppm	54.21	53.15	79.33	81.00	13.02	12.34	0.29	0.22
T <sub>5</sub> Ethephon @ 50ppm	63.06	60.67	80.33	82.67	14.76	14.36	0.46	0.36
T <sub>6</sub> Ethephon @ 100ppm	58.26	56.44	78.33	79.67	13.24	13.04	0.40	0.31
T <sub>7</sub> Ethephon @ 150ppm	53.38	52.43	77.33	78.33	12.43	12.12	0.36	0.28
T <sub>8</sub> Urea @ 0.2%	73.12	70.61	84.33	86.67	18.76	17.04	0.67	0.49
T <sub>9</sub> Urea @ 0.4%	69.41	66.29	83.00	85.00	17.17	16.55	0.55	0.44
T <sub>10</sub> Urea @ 0.6%	66.73	63.21	82.33	84.33	16.28	15.58	0.45	0.33
<b>CD<sub>(p≤0.05)</sub></b>	<b>1.79</b>	<b>1.71</b>	<b>1.14</b>	<b>1.01</b>	<b>1.52</b>	<b>1.60</b>	<b>0.05</b>	<b>0.03</b>

**Table 5: Effect of chemical thinning on physical fruit characteristics of nectarine cv. Silver King during 2016 and 2017**

Treatments		Fruit length (cm)		Fruit diameter (cm)		Fruit weight (g)		Fruit firmness (kg cm <sup>-2</sup> )	
		2016	2017	2016	2017	2016	2017	2016	2017
T <sub>1</sub>	Control	4.12	4.07	3.91	3.83	44.59	45.01	9.17	9.06
T <sub>2</sub>	Naphthalene acetic acid @ 10ppm	4.31	4.26	4.12	4.03	49.63	50.04	9.02	8.91
T <sub>3</sub>	Naphthalene acetic acid @ 20ppm	4.92	5.01	4.74	4.78	61.15	62.12	8.63	8.55
T <sub>4</sub>	Naphthalene acetic acid @ 30ppm	5.23	5.22	5.00	5.01	72.62	71.45	8.22	8.12
T <sub>5</sub>	Ethephon @ 50ppm	4.94	4.82	4.68	4.59	62.65	63.26	8.53	8.42
T <sub>6</sub>	Ethephon @ 100ppm	5.06	5.01	4.90	4.83	67.44	68.37	8.41	8.33
T <sub>7</sub>	Ethephon @ 150ppm	5.25	5.24	5.05	5.03	73.62	72.61	8.16	8.04
T <sub>8</sub>	Urea @ 0.2%	4.27	4.23	4.08	4.01	48.24	47.63	9.08	8.96
T <sub>9</sub>	Urea @ 0.4%	4.48	4.45	4.25	4.24	51.02	50.42	8.91	8.84
T <sub>10</sub>	Urea @ 0.6%	4.68	4.66	4.44	4.38	53.40	52.56	8.77	8.66
<b>CD<sub>(p≤0.05)</sub></b>		<b>0.21</b>	<b>0.24</b>	<b>0.15</b>	<b>0.16</b>	<b>4.94</b>	<b>5.31</b>	<b>0.17</b>	<b>0.15</b>

**Table 6: Effect of chemical thinning on fruit colour of nectarine cv. Silver King during 2016 and 2017**

Treatments	Fruit colour (L* a H)					
	2016			2017		
	L*	A	°H	L*	A	°H
T <sub>1</sub> Control	34.25	23.72	32.06	33.54	23.68	32.19
T <sub>2</sub> Naphthalene acetic acid @ 10ppm	32.27	25.07	30.06	31.36	24.65	30.51
T <sub>3</sub> Naphthalene acetic acid @ 20ppm	28.27	29.58	24.82	27.38	29.43	25.17
T <sub>4</sub> Naphthalene acetic acid @ 30ppm	25.66	32.12	22.52	24.58	31.39	23.51
T <sub>5</sub> Ethephon @ 50ppm	27.34	30.53	24.27	25.29	30.33	24.56
T <sub>6</sub> Ethephon @ 100ppm	26.65	31.64	23.30	25.67	32.33	23.07
T <sub>7</sub> Ethephon @ 150ppm	24.35	32.50	22.01	23.37	33.16	22.14
T <sub>8</sub> Urea @ 0.2%	33.35	24.44	30.77	32.42	24.23	31.21
T <sub>9</sub> Urea @ 0.4%	31.26	26.35	28.15	30.29	25.29	29.35
T <sub>10</sub> Urea @ 0.6%	30.21	27.58	26.54	29.34	27.33	27.25
<b>CD<sub>(p≤0.05)</sub></b>	<b>1.15</b>	<b>2.15</b>	<b>2.16</b>	<b>2.32</b>	<b>1.86</b>	<b>1.79</b>

**Table 7: Effect of chemical thinning on fruit chemical characteristics of nectarine cv. Silver King during 2016 and 2017**

Treatments	SSC (%)		Titratable acidity (%)		SSC/Acidity ratio		Total sugars (%)	
	2016	2017	2016	2017	2016	2017	2016	2017
T <sub>1</sub> Control	11.15	11.07	0.60	0.61	18.48	18.25	8.33	8.30
T <sub>2</sub> Naphthalene acetic acid @ 10ppm	11.32	11.23	0.58	0.59	19.41	19.14	8.48	8.42
T <sub>3</sub> Naphthalene acetic acid @ 20ppm	11.88	11.83	0.53	0.53	22.46	22.47	9.03	9.05
T <sub>4</sub> Naphthalene acetic acid @ 30ppm	12.71	12.81	0.50	0.50	25.76	25.80	9.37	9.49
T <sub>5</sub> Ethephon @ 50ppm	12.06	12.03	0.54	0.54	22.51	22.42	9.08	9.12
T <sub>6</sub> Ethephon @ 100ppm	12.27	12.26	0.51	0.52	24.07	23.73	9.17	9.21
T <sub>7</sub> Ethephon @ 150ppm	12.77	13.11	0.50	0.49	26.04	25.72	9.45	9.53
T <sub>8</sub> Urea @ 0.2%	11.26	11.17	0.59	0.60	19.21	18.63	8.42	8.38
T <sub>9</sub> Urea @ 0.4%	11.47	11.38	0.57	0.58	20.02	19.74	8.62	8.56
T <sub>10</sub> Urea @ 0.6%	11.76	11.69	0.55	0.54	21.28	21.70	8.77	8.68
<b>CD<sub>(p≤0.05)</sub></b>	<b>0.40</b>	<b>0.41</b>	<b>0.021</b>	<b>0.022</b>	<b>1.11</b>	<b>1.09</b>	<b>0.19</b>	<b>0.24</b>

**Table 8: Effect of chemical thinning on fruit nutrient status of nectarine cv. Silver King during 2016 and 2017**

Treatments		N (%)		P (%)		K (%)		Ca (%)		Mg (%)	
		2016	2017	2016	2017	2016	2017	2016	2017	2016	2017
T <sub>1</sub>	Control	0.61	0.60	0.042	0.041	0.71	0.70	0.068	0.069	0.022	0.021
T <sub>2</sub>	Naphthalene acetic acid @ 10ppm	0.62	0.61	0.043	0.042	0.73	0.72	0.067	0.066	0.022	0.022
T <sub>3</sub>	Naphthalene acetic acid @ 20ppm	0.66	0.67	0.047	0.049	0.78	0.79	0.063	0.064	0.027	0.028
T <sub>4</sub>	Naphthalene acetic acid @ 30ppm	0.70	0.72	0.051	0.054	0.83	0.85	0.061	0.060	0.029	0.030
T <sub>5</sub>	Ethephon @ 50ppm	0.63	0.64	0.046	0.047	0.76	0.77	0.063	0.062	0.026	0.025
T <sub>6</sub>	Ethephon @ 100ppm	0.66	0.67	0.048	0.049	0.77	0.78	0.062	0.061	0.027	0.028
T <sub>7</sub>	Ethephon @ 150ppm	0.68	0.69	0.049	0.051	0.82	0.84	0.060	0.059	0.028	0.029
T <sub>8</sub>	Urea @ 0.2%	0.62	0.62	0.042	0.042	0.72	0.71	0.067	0.068	0.023	0.023
T <sub>9</sub>	Urea @ 0.4%	0.65	0.64	0.044	0.045	0.74	0.73	0.066	0.067	0.024	0.025
T <sub>10</sub>	Urea @ 0.6%	0.69	0.71	0.050	0.053	0.79	0.80	0.065	0.064	0.025	0.026
<b>CD<sub>(p≤0.05)</sub></b>		<b>0.017</b>	<b>0.013</b>	<b>NS</b>	<b>NS</b>	<b>0.027</b>	<b>0.034</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>

## Conclusion

The results indicate that chemical thinning, particularly with NAA and ethephon, holds potential for optimizing fruit quality and marketability in the 'Silver King' nectarine cultivar.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## Reference

Basak, A. 2006. The effect of fruitlet thinning on fruit quality parameters in the apple cultivar 'Gala'. *Journal of Fruit and Ornamental Plant Research*. **14** (2): 143-150.

Chandel, J. S. and Singh, J. 2015. Effect of chemical and hand thinning on growth, yield and fruit quality of nectarine (*Prunus persica* Batsch var. *nucipersica*). *Indian Journal of Horticulture* **72** (1): 28-32.

Devlal, S., Bhardwaj, V. K. and Uniyal, S. 2017. Response of fruit thinning on yield and quality of peach cv. Red June. *Journal of Hill Agriculture* **8** (3): 297-301.

Devnath, S. and Kundu, S. 2001. Effect of bioregulators and nutrients on growth and differentiation of mango (*Mangifera indica* L.) shoots. *Environment and Ecology* **19** (4): 829-832.

Hamad, W. J. and Mohammad, A. A. 1990. Effect of Ethrel on vegetative growth, flowering and fruit characteristics of Ahmer Sayfi apple cultivar. *Annals-of-Agricultural-Science-Cairo* **35** (1): 607-622.

Jackson, D. L. and Looney, N. E. 1999. Use of Bioregulators in Fruit Production. In: Temperate and Sub Tropical Fruit production, Jackson, D.I. and N.E. Looney, (Eds.) CAB International Oxford, pp: 101-106.

Meitei, S. B., Patel, R. K., Deka, B. C., Deshmukh, N. A. and Singh, A. 2013. Effect of chemical thinning on yield and quality of peach cv. Flordasun. *African Journal of Agricultural Research* **8** (27): 3558-3565.

Meitei, S. B., Patel, R. K., Deka, B. C., Deshmukh, N. A. and Singh, A. 2013. Effect of chemical thinning on yield and quality of peach cv. Flordasun. *African Journal of Agricultural Research* **8** (27): 3558-3565.

Rajiv, K., Rimpika, N. Shylla, B., Thakur, A. and Sharma D.P. 2017. Influence of manual and chemical thinning on yield and quality of nectarine (*Prunus persica* (L.) Batch var. *nucipersica*) cv. Snow Queen. *International Journal of Bio-resource and Stress management* **8** (5): 601-604.

Rajput, V., Bhatia, S.K., Kumatkar, R.B. and Sharma, S. 2017. Study of chemical blossom thinning on fruiting and quality in Japanese Plum (*Prunus salicina* Lindl.) cv. Kala Amritsari. *Environment and Ecology* **35** (4A): 2831-2835.

Rimpika, Sharma, N. and Sharma, D.P. 2017. Effect of chemical thinning, gibberellic acid and pruning on growth and production of nectarine ( *Prunus persica* (L.) Batch var. *nucipersica*) cv. May fire. *Journal of Applied and Natural Science* **9** (1): 332-337.

Sandhu A S and Singh Z. 2001. Effect of ethephon on maturation and fruit quality of peach (*Prunus persica* Batsch.). *Punjab Horticulture Journal* **23**(1-4): 172-175.

Sefick, H. J. and Ridley, J. D. 1988. Fruit thinning. **In:** The peach, world cultivars to marketing.

Sharma, N., Singh, R. P. and Singh, B. 2001. Influence of chemical and hand thinning on

maturity, quality and colour of fruits in Redhaven peaches. *Horticulture Journal* **14**(3): 6-10.

Sharma, N., Singh, R. P. and Singh, B. 2003. Effect of chemical and manual thinning on productivity and fruit size of Redhaven peach. *Indian Journal of Horticulture* **60** (3): 239-243.

Shukla, A.K., Singh, D., Shukla, A.K. and Meena, S.R. (2007). Pruning and training of fruit crops. In: Fruit production technology, P.K. Yadav (ed.). International Book Distributing Company Publishing Division, pp.135-148.

Taghipour, L. and Rahemi, M. 2012. The influence of fruit thinning on the apricot cultivar Gerdi. *International Research Journal of Environmental Sciences* **7** (3): 321-324.

Taghipour, L. and Rahemi, M. 2012. The influence of fruit thinning on the apricot cultivar Gerdi. *International Research Journal of Environmental Sciences* **7** (3): 321-324.

Taheri A, Cline J A, Jayasankar S and Pauls P K. 2012. Ethephon induced abscission of "redhaven" peach. *American Journal of Plant sciences* **3**(2): 295-301.

Vego, D., Saravanja, P. and Knezovic, Z. 2010. Fruit thinning of peach and nectarine. *Acta Horticulturae* **884**: 695-699.

Whale, S., Singh, J. and Janes, J. 2012. Ethylene biosynthesis and fruit color development in 'Pink Lady' apples during growth and maturation. *Hort. Science* **39**(4): 762.

Yadav, P.K. 2007. Fruit Production Technology. International Book Distributing Company, Lucknow (U.P.) India, pp.372

Zilkkha, S. Klein, I. and David, I. 1989. Thinning peaches and nectarines with urea. *Journal of Horticultural Sciences* **63**: 2019-216.