

Review Article

A Comprehensive Review on Cucurbit Grafting: A Sustainable Approach to Boost Crop Performance


Abstract: Grafting, which unites the rootstock and scion of a plant, is crucial for combating soil-borne diseases in cucurbitaceous crops such as watermelon, melon, bitter melon, summer squash and cucumber. Essential due to limited arable land and off-season demand, this technique enhances plant vigour by improving nutrient uptake and providing resistance to soil pathogens, salinity, drought and temperature variations, thereby improving overall yield and fruit quality. Initially implemented in Japan in the late 1990s, grafting significantly affects fruit pH, flavour, sugar content, carotenoids, and texture. Globally, it has been adopted to prevent diseases like fusarium wilt and root-knot nematode. This practice mitigates production losses under adverse conditions and promotes organic, safer vegetables by reducing pesticide residues. Recent advancements in grafting methods and versatile rootstocks further support its widespread use.

Keywords: Grafting, abiotic stress, biotic stress, cucurbits and rootstocks.

1. Introduction: Vegetables are crucial for global food and nutritional security, but their production is challenged by pests, diseases and environmental stresses (Tirupathamma *et al.*, 2019). Farmers often rely on agrochemicals, which can be harmful and costly (Colla *et al.*, 2010). Grafting, a technique long used in fruit cultivation, is now gaining popularity among vegetable growers for its effectiveness in reducing agrochemical use and managing soil-borne problems. The first vegetable grafting attempt in Japan in the 1920s targeted Fusarium wilt using watermelon grafted onto pumpkin (Leonardi, 2017). Today, grafting is common in cucurbit cultivation, especially in greenhouses in Japan and Korea (Kumar *et al.*, 2018). Despite its benefits, grafting is not suitable for all vegetables due to variations in species characteristics. Cucurbits and solanaceous crops are the most commonly grafted (Schwarz *et al.*, 2010). Methods such as hole insertion, tongue approach, splice and cleft grafting are used, and automation is being explored to reduce costs (Comba *et al.*, 2016). This review examines the role of grafting in improving cucurbitaceous crops.

2. Grafting method: Selection of grafting is done on crop with the help of the farmers experience, the number of grafts required, the motive of grafting and available facilities required for machinery and infrastructure (Lee *et al.*, 2010).


2.1 Tongue / approach grafting: Grafting involves using rootstocks and scions of equal size. Although this method requires more labour and space, it results in a high seedling survival rate and is most



commonly practiced by small nursery farmers. However, it is not suitable for rootstocks with hollow hypocotyls.

2.2 Cleft grafting: This method, also known as apical or wedge grafting, involves making a slant cut in the lower stem of the rootstock. The scion, pruned to have 1-3 true leaves, is then placed into the split, and a clip is attached between the scion and rootstock (Johnson *et al.*, 2019). This technique is primarily used for solanaceous crops.

2.3 Hole insertion: This method, also known as top insertion grafting, is commonly practiced in China. It is primarily used for watermelon, as the seedlings are smaller than the rootstocks of bottle gourd or squash. For successful transplanting, an optimal temperature range of 21-36°C is required (Shalaby *et al.* 2022).



2.4 Splice grafting: This method is widely used by growers and can be performed manually or with machines for most cucurbit and solanaceous vegetable crops.

2.5 Pin grafting: In this method, specially designed pins are used to secure the grafted position instead of using grafting clips. It is similar to splice grafting.

3. Basic Requirements for Vegetable Grafting:

3.1 Selection of Rootstock and Scion: Grafting is typically performed when plants are at the 2-3 true leaf stage. The selection of rootstock and scion should be based on matching stem diameters.

3.2 Compatibility of Graft: Ensuring graft compatibility reduces the mortality rate even at later growth stages. Successful grafting depends on callus formation between the scion and rootstock, which leads to the formation of vascular bundles (Sanwala *et al.* 2022).

3.3 Grafting Aids: Grafting is facilitated using tools such as grafting clips, tubes, pins, and a grafting blade.

3.4 Screening House: A screening house, constructed with 60-mesh nylon netting, is used to grow seedlings prior to grafting.

3.5 Healing of Grafts: The healing chamber should maintain a temperature of 28-29°C and a relative humidity of 95 per cent for 5-7 days to promote callus formation. This environment reduces light intensity and transpiration, aiding in better attachment of the scion and rootstock.

3.6 Acclimatization of Grafted Plants: After callus formation is complete and the wounded surfaces have healed, the grafted plants should be transferred to a mist chamber or covered with clean plastic in a greenhouse. This helps prevent leaf burning and wilting during the acclimatization process.

4. Current status of vegetable grafting:

Due to the high preference and production of cucurbits and other grafted vegetables in East Asia, this region remains the largest market for vegetable grafting. Watermelon is grown using grafted transplants to varying degrees in Korea (99%), Japan (94%), and China (40%). For solanaceous vegetables, approximately 60-65% of tomatoes and eggplants, and 10-14% of peppers, are cultivated using grafted transplants. In the Netherlands, grafted tomato transplants are used for all soilless tomato production. The popularity of vegetable grafting is rapidly increasing globally, especially in Eastern Europe, North and South America, India, and the Philippines. In China, more than 1,500 industrial nurseries produce grafted transplants, and the international trade of these grafted vegetable transplants is on the rise (Bieet *et al.*, 2017). Worldwide, cucurbitaceous crops dominate vegetable grafting. However, this technology is still in its infancy in India (Aydin *et al.* 2024).

5. Grafting in cucurbitaceous crops:

Grafting is done to control Fusarium wilt and tolerant to drought and flooding. Currently, in this family watermelon is one crop in which grafting is practiced in the world. There are different grafting methods used in cucurbits are listed in the below Table.1

Table 1: Grafting methods and rootstocks used in cucurbit crops

Scion plant	Rootstock	Method
Cucumber	<i>Cucurbita moschata</i> , <i>Cucurbita maxima</i>	Tongue and top insertion method
Watermelon	<i>Benincasa hispida</i> , <i>Cucurbita moschata</i> , <i>Cucurbita melo</i> , <i>Cucurbita moschata</i> × <i>Cucurbita maxima</i> , <i>Lagenaria siceraria</i>	Top insertion and cleft method and slice grafting
Bitter gourd	<i>Cucurbita moschata</i> , <i>Lagenaria siceraria</i>	Top insertion and tongue method
Bottle gourd	<i>Cucurbita moschata</i> , <i>Luffa sp.</i>	Top insertion and tongue method.

5.1 Graft compatibility and survival rate:

Graft compatibility is defined as the genetic closeness between rootstock and scion required for a successful graft union, provided other factors such as technique, timing, and temperature is optimal. For instance, inter-generic grafting of highly nematode-susceptible watermelon cultivars 'Congo' and 'Charleston Gray' onto highly nematode-resistant wild watermelon (*Cucumis africanus*) and wild cucumber (*C. myriocarpus*) rootstocks resulted in a 36% survival rate of the grafts (Pofu and Mashela).

Table 2. Different rootstocks used for biotic and abiotic stress in cucurbit crops

Rootstock	Usedfor
Cucumber	
Fig leaf Gourd	Good Fusarium resistance and low temperature resistance
Burr cucumber (<i>Sicyosangulatus</i>)	Southern Root-knot Nematode resistance
Interspecific hybrid squash (<i>Cucurbita maximax C. moschata</i>)	Fusarium wilt and low temperature resistance
African horned cucumber (<i>Cucumis metuliferus</i>)	Excellent fusarium resistance and good nematode tolerance
<i>Cucurbita moschata</i>	"Blooming-less" cucumber fruit.
<i>Cucurbita moschata</i>	Improves stress tolerance
TRC11550 (Intraspecific Hybrid)	Nematode resistance
Ardito/TRC1401	Very strong and uniform rootstock with highly vigorous root system.
TRC 15NTB4	Resistance to soil-borne diseases and abiotic stress factors.
Watermelon	
Bottle Gourd (<i>Lagenaria siceraria</i>)	Fusarium wilt and chilling tolerance
Shintoza	Fusarium wilt and chilling tolerance
Wax Gourd	Greater drought tolerance
Squash (<i>Cucurbita moschata</i>)	Resistant to fusarium and low temperature
Pumpkins (<i>Cucurbita pepo</i>)	Vigorous root system, resistant to fusarium and low temperature
Watermelon (<i>Citrullus lanatus</i>)	Fusarium tolerance, but not resistance
Citron melon	Root knot Nematode
African horned cucumber (<i>Cucumis metuliferus</i>)	Excellent fusarium resistance and good nematode tolerance
Interspecific hybrid squash (<i>Cucurbita maximax C. moschata</i>)	Resistant to fusarium wilt and vigorous root system ,high temperature tolerance
Ash gourd (<i>Benincasa hispida</i>)	Good disease resistance
Summer squash	
Figleaf Gourd (<i>Cucurbita ficifolia</i>)	Low soil temperature tolerance
Bitter gourd	
Sponge gourd (<i>Luffa cylindrica</i>)	Resistant to fusarium wilt, good tolerance to heat and flooding
Melon	
Squash	Fusarium and low temperature tolerance

(<i>Cucurbita maxima</i> x <i>C. moschata</i>)	Fusarium resistance, low and high soil temperature tolerance, and high soil moisture tolerance
<i>Cucurbita pepo</i>	Fusarium resistance, low and high soil temperature tolerance, and high soil moisture tolerance
Melon (<i>Cucumis melo</i>)	Fusarium tolerance and good fruit quality
African horned cucumber (<i>Cucumis metuliferus</i>)	Fusarium tolerance, low and high soil moisture tolerance and nematode tolerance

5.2 Potential rootstocks for different cucurbitaceous crops:

Watermelon is typically grafted onto rootstocks like *Cucurbita* species, *Citrullus* species and bottle gourd, with primary rootstocks being bottle gourd, interspecific hybrids between *Cucurbita moschata* and *Cucurbita maxima* and wild watermelon. *Cucurbita* spp. rootstocks, particularly interspecific hybrids of *C. maxima* and *C. moschata*, are effective against soil-borne pathogens and abiotic stresses (Sakata *et al.*, 2008). Fig leaf gourd (*C. ficifolia*) is favoured for cucumber due to its compatibility, cold tolerance, and Fusarium wilt resistance, while *Cucurbita* interspecific hybrids are preferred for summer cucumbers for their Fusarium resistance and heat tolerance (King *et al.*, 2010).

5.3 Influence of cucurbit rootstocks to biotic stresses

5.3.1 Grafting against verticillium wilt:

Grafting is commonly used in tomato and brinjal to combat Verticillium wilt, but it's less applied in cucurbits. Grafting improves Verticillium wilt management by boosting root vigor and nutrient uptake (Bletsos *et al.*, 2008). Klosterman (2009) found that reduced infestation is due to the pathogen's poor and slow colonization of vascular tissues and decreased wilt severity is linked to less pathogen colonization. Root exudates from both resistant and susceptible rootstocks inhibit Verticillium growth (Liu *et al.*, 2009). "Crimson Sweet" watermelon grafted onto "Emphasis" and "Strongtoza" rootstocks showed increased resistance to *V. dahliae* (Buller *et al.*, 2013). Grafted watermelon plants had 3.5 times less disease severity compared to non-grafted 'Secretariat' plants (Devi *et al.*, 2020). Effective rootstocks like Pelops, Round, Super Shintoza, and Tetsukabuto also significantly mitigated Verticillium wilt.

5.3.2 Grafting against fusarium wilt:

Grafting is more effective for managing Fusarium wilt in cucurbits compared to solanaceous crops. In watermelon cultivation, interspecific hybrids and rootstocks like "Shintoza" and "Super Shintoza" (*C. maxima* x *C. moschata*) improve tolerance to Fusarium and Verticillium wilt, enhancing fruit size and yield (Gaionet *et al.*, 2018). For cucumbers, rootstocks such as fig leaf gourd and ISHc'Shintoza' offer resistance to Fusarium and environmental stress. Bitter gourd grafted onto *C. colocynthis*, *C. metuliferus*, and *C. moschata* exhibits reduced Fusarium wilt, with these rootstocks being the most effective (Gaionet *et al.*, 2018). In 2021, Reyad *et al.* conducted an experiment at Cairo

University in Giza, Egypt, and found that cucumbers grafted onto *Cucurbita maxima* and *C. moschata* exhibited greater resistance to Fusarium wilt.

5.3.3 Grafting against root-knot nematode:

Commercial cucurbits lack natural resistance to root-knot nematodes (RKN), necessitating resistant rootstocks (Cohen *et al.*, 2007). Root-knot nematodes cause galls that impair water and nutrient uptake, leading to poor plant and fruit quality. Grafting cantaloupes onto *C. moschata* and *C. metuliferus* rootstocks reduced galling, with *C. metuliferus* showing moderate resistance (Pradhan *et al.*, 2017). "Strongtosa" and "Shintosa" rootstocks significantly increased yields (Al-Debeiet *al.*, 2011). *Citrullus colocynthis* and *Cucumis metuliferus* were identified as resistant rootstocks for cucumbers (Punithaveniet *al.*, 2015; Thangamani *et al.*, 2018). Despite their tolerance, some common rootstocks are vulnerable to nematode buildup and should be used cautiously.

5.4 Influence of cucurbit rootstocks to abiotic stresses




Figure 1. Impact of abiotic stresses on plant physiology, growth and productivity

5.4.1 Cold/heat:

Temperature extremes can hinder vegetable production by delaying flowering and fruit ripening. Using low-temperature tolerant rootstocks for grafting superior cultivars offers a quick alternative to lengthy breeding processes (Schwarz *et al.*, 2010). For example, watermelon is grafted onto "Shintosa" rootstocks to hasten planting in cool weather, while fig leaf gourd and bur cucumber are used to enhance cold resistance in cucumbers (Padilla *et al.*, 2023). Grafted cucumbers on

squash rootstocks can better endure adverse conditions, and luffa rootstock-grafted cucumbers can tolerate heat stress (Gaionet *et al.*, 2018). This was reflected in better growth parameters, higher chlorophyll and proline levels, reduced ROS and lipid peroxidation, increased antioxidant enzyme activities (especially CAT and GPX), and elevated expression of Calvin cycle enzymes (Lu *et al.*, 2021).

5.4.2 Drought:

Drought significantly impacts global crop productivity and quality. Grafting high-yielding varieties onto drought-tolerant rootstocks can mitigate water stress effects and improve water use efficiency. For example, mini-watermelons grafted onto the commercial rootstock PS 1313 (*Cucurbita maxima* × *Cucurbita moschata* hybrid) yielded 60% more marketable fruit under drought conditions (Gaionet *et al.*, 2018). In cucumbers, grafting drought-tolerant rootstocks increased commercial yield, enhanced defense gene expression, and improved various physiological parameters such as chlorophyll content, photosystem II efficiency, and antioxidant enzyme activity, while reducing electrolyte leakage and stress markers (Shehata *et al.*, 2022).

5.4.3 Soil salinity:

Soil salinity affects up to 20% of irrigated land and 7% of global land (Shahid *et al.*, 2018). For instance, grafting watermelon onto bottle gourd significantly improves salt tolerance (Yang *et al.*, 2013). Grafting cucumber onto pumpkin (*Cucurbita moschata*) enhances salt tolerance, with grafting onto "Chaojiquanwang" rootstock reducing Na concentration in the scion by limiting Na transport (Huang *et al.*, 2014). Additionally, grafting cucumber onto highly salt-tolerant Luffa rootstocks improves salinity tolerance and fruit quality (Guo *et al.*, 2023). Rootstocks improved salinity tolerance in crops, as demonstrated by Usanmaz *et al.* (2019) in North Cyprus, Turkey. Using four commercial cucumber rootstocks (TZ148 F1, RS841 F1, Nun9075 F1 and Avar F1) and two local *Cucurbita moschata* landraces (Local-1 and Local-3) under three different salinity levels, the study found that grafted crops had higher growth and yield than non-grafted ones. A study by Usanmaz and Abak (2019) found that cucumbers grafted on *Cucurbita moschata* L. rootstocks (Nun 9075, TZ148, and Local-3) showed increased salinity tolerance under an EC of 5.0 dSm⁻¹. The RS841 rootstock enhanced tolerance at both 5.0 and 7.5 dSm⁻¹ by reducing Na⁺ accumulation in leaves.

5.4.4 Flooding:

Many vegetable crops are highly sensitive to flooding caused by heavy rainfall, with some unable to tolerate waterlogged conditions (Singh *et al.*, 2020). Grafting has been shown to improve flooding tolerance in various crops (Bhatt *et al.*, 2015). For example, watermelon grafted onto landrace *Lagenaria siceraria* showed reduced chlorosis under flood conditions. In contaminated soils, watermelon cv. Arava grafted onto *cucurbita* rootstocks exhibited lower levels of harmful heavy metals. Interspecific hybrids (*C. maxima* × *C. moschata*) and certain melon cultivars like Proteo grafted onto hybrids (Dinero, Magnus, RS841, Shintoza, Strongtosa) have been found to reduce heavy metal uptake, with RS841 being particularly effective in maintaining productivity and reducing arsenic transfer to fruits (Allevato *et al.*, 2019).

5.5 Influence of cucurbit rootstocks on qualitative and quantitative characters:

Grafting is widely used by farmers to enhance fruit yield and plant vigour by improving water and nutrient uptake (Lee *et al.*, 2010). Rafting enhances yield by improving root systems, photosynthesis, and disease resistance. For instance, cucumber cv Kalaam grafted onto *Lagenaria siceraria* rootstocks showed superior vegetative growth, fruit yield, and quality compared to other rootstocks (Noor *et al.*, 2019). Similarly, grafted cucumbers on pumpkin rootstocks produced 27 per cent more marketable fruit per plant and yielded 8.4 kg per vine under copper toxicity (Rouphae *et al.*, 2008). Pugalendhi *et al.* (2019) found that grafted bitter melon plants yielded 63.2% more than self-rooted plants. Similarly, Tamilselvi and Pugalendhi (2017) and Tamilselvi *et al.* (2015) reported that *Palee F1* grafted onto *Cucurbita moschata* produced more fruit and higher fruit weight per vine compared to other combinations. Watermelon grafted onto *Lagenaria* and *C. maxima* rootstocks resulted in 24 per cent and 27 per cent firmer fruits, respectively, and had rind thicknesses 21 per cent and 17 per cent higher than ungrafted plants (Proietti *et al.*, 2008). This increased rind thickness, while generating more waste, improved postharvest handling and injury prevention. Melons grafted onto interspecific hybrid rootstocks (*Cucurbita maxima* × *Cucurbita moschata*) experienced severe premature internal decay.

Watermelon grafted onto bottle gourd rootstock produces female flowers early in the crop's lifecycle. In contrast, pumpkin, bottle gourd, watermelon, and wax gourd grafted onto 'Shintosa'-type rootstock showed delayed flowering. Flowering time significantly affects harvest timing and produce quality (Sanket *et al.*, 2017). Lecholocholo *et al.* (2020-21) conducted an experiment in South Africa to study the effect of rootstocks on the quality and volatile constituents of cantaloupe and honeydew melons. Using Kickstart and Carnivor rootstocks (a cross between sweet melon and watermelon) grafted with four melon cultivars, they found that grafted fruits had higher soluble sugar-to-titratable acidity ratio, higher pH, increased fruit firmness, and higher ascorbic acid content compared to non-grafted controls.

6. Certain challenges associated with grafting:

Grafting poses challenges such as the need for skilled labour and post-graft care for 7 to 10 days. Many rural farmers lack knowledge about rootstock-scion compatibility, which is essential for successful grafting. Achieving good synchronization, germination rates, and graft success is crucial but often difficult (Sen *et al.*, 2018). Maintaining ideal conditions for temperature, humidity, and light is challenging in open fields, and grafting increases disease risks, especially from nursery infections (Maurya *et al.*, 2019). Additionally, consistent supply and high cost of rootstocks and grafting tools are significant issues for large-scale production.

7. Future prospects:

Grafting boosts plant performance and supports off-season vegetable production, and it can be a key element in Integrated Pest Management for organic farming. Despite its benefits, grafting is labour-intensive and time-consuming, and the availability of planting materials for widespread use is

limited. However, the growing focus on organic and sustainable farming presents a significant opportunity for this eco-friendly technique. There is a need to develop more efficient grafting methods, both manual and mechanical, and to identify, collect, and conserve indigenous and wild rootstocks that are better suited to local conditions, rather than relying solely on exotic sources.

8. Conclusion:

Many countries have adopted this technique, achieving higher profits through healthy rootstocks. Rootstock-scion compatibility is crucial for successful grafting, but the mechanisms behind graft union formation need further study. While grafting addresses issues like pest and disease susceptibility, soil salinity and drought, and promotes organic farming, it remains labour-intensive and requires skilled handling. Recent advances in automated grafting robots have improved efficiency, but challenges such as disease transmission and limited tool availability still need to be addressed. Developing better breeding techniques to enhance rootstock-scion compatibility is essential.

References

1. Al-Debei, H.S., Makhadmeh, I., Abu-Al Ruz, I., Al-Abdallat, A.M., Ayad, J.Y. and Al Amin, N. 2011. Influence of different rootstocks on growth and yield of cucumber (*Cucumis sativus* L.) under the impact of soil-borne pathogens in Jordan. *J. Food Agri. Environ.* **10**: 343-349.
2. Allevato, E., Mauro, R.P., Stazi, S.R., Marabottini, R., Leonardi, C., Ierna, A. and Giuffrida, F. 2019. Arsenic accumulation in grafted melon plants: Role of rootstock in modulating root-to-shoot translocation and physiological response. *Agronomy.* **9**: 828.
3. Aydin, A. 2024. Effects of grafting with wild tomato (*Solanum pimpinellifolium* and *Solanum habrochaites*) rootstocks on growth and leaf mineral accumulation in salt stress. *Horticulture, Environment and Biotechnology.***10**:1-17.
4. Bhatt, R.M., Upreti, K.K., Divya, M.H., Bhat, S., Pavithra, C.B. and Sadashiva, A.T. (2015). Interspecific grafting to enhance physiological resilience to flooding stress in tomato (*Solanum lycopersicum* L.). *Scientia Horticulturae.* **182**: 8-17.
5. Bie, Z.M.A., Nawaz, Y, Huang, J.M, Lee Colla ,G. 2017. Introduction to Vegetable Grafting. **18**: 1-21.
6. Bletsos, F.A. and Olympios, C.M. 2008. Rootstocks and grafting of tomatoes, peppers and eggplants for soil-borne disease resistance, improved yield and quality. *The European J. Plant Sci. Bio.* **2**: 62-73.
7. Buller, S., Inglis, D. and Miles, C. 2013. Plant growth, fruit yield and quality and tolerance to verticillium wilt of grafted watermelon and tomato in field production in the Pacific Northwest. *Hort Science.* **48**: 1003-1009.
8. Colla, G., Roupshael, Y., Leonardi, C. and Bie, Z. 2010. Role of grafting in vegetable crops grown under saline conditions. *Scientia Horticulturae.* **127**: 147-155.
9. Comba, L., Gay, P. and Aimonino, D. R. 2016. Robot ensembles for grafting herbaceous crops. *Bio. Eng.***146**: 227-239.

10. Crino, P., Bianco, C., Roupshael, Y., Colla, G., Saccardo, F., Paratore, A. 2007. Evaluation of rootstock resistance to fusarium wilt and gummy stem blight and effect on yield and quality of a grafted 'inodorus' melon. *Hort. Science*.**42**:521-525.
11. Davis, A.R., Perkins-Veazie, P., Sakata, Y., Lopez-Galarza, S., Maroto, J.V., Lee, S.G. 2008. Cucurbit Grafting. *Crit. Rev. Plant Sci*.**27**:50-74.
12. Devi, P., Perkins-Veazie, P. and Miles, C.A. 2020. Rootstock and plastic mulch effect on watermelon flowering and fruit maturity in a verticillium dahliae-infested field. *Hort Science*. **55**: 1438-1445.
13. Gaion, L.A., Braz, L.T. and Carvalho, R.F. 2018. Grafting in vegetable crops: A great technique for agriculture. *Int. J. Veg. Sci*. **24**: 85-102.
14. Guo, Z., Qin, Y., Lv, J., Wang, X., Dong, H., Dong, X. and Piao, F. 2023. Luffa rootstock enhances salt tolerance and improves yield and quality of grafted cucumber plants by reducing sodium transport to the shoot. *Environmental Pollution*. **316**: 120521.
15. Huang, Y., Kong, Q.S., Chen, F. and Bie, Z.L. 2014. The History, Current Status and Future Prospects of Vegetable Grafting in China. In: I International Symposium on Vegetable Grafting. *Acta Horticulturae*.**1086**: 31-39.
16. Johnson, S., Kreider, P., Miles, C. 2019. Vegetable Grafting Eggplants and Tomatoes Washington State University.
17. Klosterman, S.J., Atallah, Z.K., Vallad, G.E. and Subbarao, K.V. 2009. Diversity, pathogenicity and management of Verticillium species. *Annual Review of Phytopathology*. **47**: 39-62.
18. Kumar, S., Bharti, N. and Saravaiya, S.N. 2018. Vegetable Grafting: A Surgical Approach to combat biotic and abiotic stresses-A review. *Agricultural Reviews*. **39**: 1-11.
19. Lee, J.M., Kubota, C., Tsao, S.J., Bie, Z., Echevarria, P.H., Morra, L. and Oda, M. 2010. Current status of vegetable grafting: Diffusion, grafting techniques, automation. *Scientia Horticulturae*. **127**: 93-105.
20. Leonardi, C.H.E.R.U.B.I.N.O., Kyriacou, M., Gisbert, C.A.R.MI. N.A., Oztekin, G.B., Mourão, I.S. A.B.E.L. and Roupshael, Y.O.U.S.S.E.F. 2017. Quality of Grafted Vegetables. Vegetable Grafting: Principles and Practices. [(Eds.) Colla, G., Pérez-Alfocea, F. and Schwarz, D.] *CAB International, Oxfordshire, UK*, 216-244.
21. Liu, N., Zhou, B., Zhao, X., Lu, B., Li, Y. and Hao, J. 2009. Grafting eggplant onto tomato rootstock to suppress Verticillium dahliae infection: The effect of root exudates. *Hort Science*.**44**: 2058-2062.
22. Lu, K., Sun, J., Li, Q., Li, X. and Jin, S. 2021. Effect of cold stress on growth, physiological characteristics and calvin-cycle related gene expression of grafted watermelon seedlings of different gourd rootstocks. *Horticulturae*.**7**: 391.
23. Maurya, D., Pandey, A.K., Kumar, V., Dubey, S. and Prakash, V. 2019. Grafting techniques in vegetable crops: A review. *Int. J. Chem. Stud*. **7**: 1664-1672.
24. Noor, R.S., Wang, Z., Umair, M., Yaseen, M., Ameen, M., Rehman, S.U. and Sun, Y. 2019. Interactive effects of grafting techniques and scion-rootstocks combinations on vegetative growth, yield and quality of cucumber (*Cucumis sativus* L.). *Agronomy*. **9**: 288.

25. Padilla, Y. G., Gisbert-Mullor, R., Lopez-Galarza, S., Albacete, A., Martinez-Melgarejo, P. A., & Calatayud, A. 2023. Short-term water stress responses of grafted pep per plants are associated with changes in the hormonal balance. *Frontiers in Plant Science*, **14**: 1170021.
26. Pradhan, S.R., Sahu, G.S., Tripathy, P., Dash, S.K., Mishra, B., Jena, R. and Sahoo, T.R. 2017. Vegetable grafting: A multidimensional approach for crop management in vegetables. *International J. Current Microbiol. Applied Sci.***6**: 3332-3345.
27. Proietti, S., Roupael, Y., Colla, G., Cardarelli, M., De Agazio, M., Zacchini, M., Moscatello, S. and Battistelli, A. 2008. Fruit quality of mini-watermelon as affected by grafting and irrigation regimes. *J. Sci. Food Agri.* **88**: 1107-1114.
28. Pugalendhi, L., Tamilselvi, N.A. and Thangamani, C. 2019. Vegetable grafting-a boon for soil-borne pest and disease management A review. *Madras Agricultural Journal.* **106**: 561-570.
29. Punithaveni, V., Jansirani, P. and Sivakumar, M. 2015. Screening of cucurbitaceous rootstocks and cucumber scions for root knot nematode resistance (*Meloidogyne incognita* Kofoid and White). *Electronic Journal of Plant Breeding.***6**: 486-492.
30. Reyad, N. E. H. A., El-Sayed, S. F., & Azoz, S. N. 2021. Evaluation of grafting using cucurbit interspecific hybrids to control fusarium wilt in cucumber. *Plant Cell Biotechnol. Mol. Biol.* **5**: 50-63.
31. Roupael, Y., Cardarelli, M., Colla, G. and Rea, E. 2008. Yield, mineral composition, water relations and water use efficiency of grafted mini-watermelon plants under deficit irrigation. *Hort Sci.***43**: 730-736.
32. Sakata, Y., Ohara, T., Sugiyama, M. 2008. The history of melon and cucumber grafting in Japan. *Acta Horticult.***767**: 217-228.
33. Sanket, K.B. and Kumar, A. 2017. Grafting of vegetable crops as a tool to improve yield and tolerance against diseases-A review. *Int. J. J. of Agri Sci.***9**:4050-4056.
34. Sanwal, S.K., Mann, A., Kumar, A., Kesh, H., Kaur, G., Rai, A.K., Kumar, R., Sharma, P.C., Kumar, A., Bahadur, A., Singh, B. and Kumar, P. 2022. Salt tolerance eggplant rootstocks modulate sodium partitioning in tomato scion and improve performance under saline conditions. *Agriculture.* **12**:1-15.
35. Schwarz, D., Roupael, Y., Colla, G. and Venema, J.H. 2010. Grafting as a tool to improve tolerance of vegetables to abiotic stresses: Thermal stress, water stress and organic pollutants. *Sci. Horti.***127**: 162-171.
36. Sen, A., Chatterjee, R., Bhaisare, P. and Subba, S. 2018. Grafting as an alternate tool for biotic and abiotic tolerance with improved growth and production of Solanaceous vegetables: Challenges and scopes in India. *International J. Current Micro. Applied Sci.***7**: 121-35.
37. Shahid, S.A., Zaman, M. and Heng, L. 2018. Soil Salinity: Historical perspectives and a world overview of the problem. In: guideline for salinity assessment, mitigation and adaptation using nuclear and related techniques. *Springer, Cham.***8**:43-53
38. Shehata, S.A., Omar, H.S., Elfaidy, A.G., El-Sayed, S.S., Abuarab, M.E. and Abdeldaym, E.A. 2022. Grafting enhances drought tolerance by regulating stress-responsive gene expression and antioxidant enzyme activities in cucumbers. *BMC Plant Biol.* **22**: 1-17.

39. Singh, H., Sethi, S., Kaushik, P. and Fulford, A. 2020. Grafting vegetables for mitigating environmental stresses under climate change: A review. *Journal of Water and Climate Change*. **11**: 1784-1797.
40. Tamilselvi, N.A. and Pugalendhi, L. 2017. Graft compatibility and anatomical studies of bitter gourd (*Momordica charantia* L.) scions with cucurbitaceous rootstocks. *International Journal of Current Microbiology and Applied Sci.* **6**: 1801-1810.
41. Tamilselvi, N.A., Pugalendhi, L. and Sivakumar, M. 2015. Response of cucurbitaceous rootstocks and bitter gourd scions against root knot nematode *Meloidogyne incognita* Kofoid and White. *Annals of Plant and Soil Research*. **17**: 175-177.
42. Thangamani, C., Pugalendhi, L. and Punithaveni, V. 2018. Screening wild and cultivated cucurbits against root knot nematode to exploit as rootstocks for grafting in cucumber. *J. Hort. Sci.* **13**: 32-41.
43. Tirupathamma, T.L., Ramana, C.V., Naidu, L. and Sasikala, K. 2019. Vegetable grafting: A multiple crop improvement methodology. *Current J. Applied Sci. Technol.* **33**: 1-10.
44. Usanmaz, S. and Abak, K. 2019. Plant growth and yield of cucumber plants grafted on different commercial and local rootstocks grown under salinity stress. *Saudi J. Bio. Sci.* **26**: 1134-1139.
45. Yang, Y., Lu, X., Yan, B., Li, B., Sun, J., Guo, S. and Tezuka, T. 2013. Bottle gourd rootstock-grafting affects nitrogen metabolism in NaCl-stressed watermelon leaves and enhances short-term salt tolerance. *J. of Plant Physio.* **170**: 653-661.