

Review Article

Pharmacogenomics: Generalities and Applications in the Pharmaceutical Sciences

ABSTRACT

Although the main objective of drugs is to improve the person's health, not all people present the same response since it can be exacerbated, adequate, or absent. This variability is associated with gender, age, lifestyle, and therapeutic adherence. Pharmacogenomics studies the genetic variability that influences patients' response to treatment, focusing mainly on the analysis of drug metabolism. This science involves numerous factors. They constitute a network of linked aspects that must be considered to analyze and evaluate the drug response. They can be divided into interindividual variability, pharmacokinetics variability, pharmacodynamic variability, and biomarkers. This knowledge has purposes in various therapeutic areas (immunological, neurological, cardiovascular, and cancer disorders). Therefore, pharmacogenomics has been a key tool in implementing personalized medicine to improve pharmacological response and minimize unwanted effects. Its implementation will increase in the short and medium term, always putting the patient's quality of life first.

Keywords: pharmacogenetics, interindividual variability, pharmacokinetics, pharmacodynamics, biomarkers.

1. INTRODUCTION

A drug is defined as a substance capable of preventing, controlling, or curing the effects that a disease causes in the patient body. This substance can be chemical or biological synthesis [1]. The discovery of drugs has been one of the most relevant historical events for society since they control symptoms and even cure illnesses, which, 100 years ago, caused many deaths worldwide [2]. As a complement, the vaccines have eradicated infections, including polio and smallpox. All this has made it possible to improve society's quality of life [3].

Although the main objective is to improve the person's health, not all people present the same response since it can be exacerbated, adequate, or absent. This variability is associated with gender, age, lifestyle, and therapeutic adherence. However, much of the action generated in the body is provoked by the expression or suppression of genes that encode proteins involved in the drug's pharmacokinetics, pharmacodynamics, and immunological processes [4]. Some products where this phenomenon is observed are:

- Warfarin: a polymorphism in the CYP2C9 gene participates in its metabolism and the S-enantiomer inactivation [5].
- Thiazide diuretics: studies have shown that the presence of polymorphisms in the genes of the Afro-descendant race allows a better response of thiazide diuretics, including hydrochlorothiazide, against arterial hypertension [6].

Through these outcomes, pharmacogenomics was born. This field is a strategy to individualize the selection and pharmacological employment to avoid adverse effects and maximize its efficacy [7], based on identifying genetic variants that influence its pharmacokinetics and pharmacodynamics [8]. Rapid advances in this science have improved the

understanding of adverse reactions, made prescribing more accurate, and reduced unnecessary costs to address such side effects [9].

Pharmacogenomics has become an essential component of personalized medicine [10], which utilizes knowledge of the genetic and molecular basis of health and disease brought on by human genome sequencing to guide decisions regarding disorder prediction, prevention, diagnosis, and treatment [11]. Additionally, it ponders the environment and lifestyle to determine the best way to prevent or treat the condition [12].

Thus, this review aims to identify the application fields of pharmacogenomics in pharmaceutical sciences.

2. PHARMACOGENOMICS CONCEPT, HISTORY, AND EVOLUTION

Pharmacogenomics is an area of pharmaceutical sciences that studies genetic variability in patients' treatment responses, focusing on drug metabolism analysis. The metabolic capacity depends on the production of enzymes, which varies from the polymorphisms found in individuals [13]. Such mutations are genetic deoxyribonucleic acid (DNA) variants in more than 1 % of the world population [14].

Polymorphisms can increase or decrease enzyme synthesis, affecting two essential pharmaceutical processes: pharmacokinetics (the processes that the body performs on a drug) and pharmacodynamics (the effects that the drug has on the body) [13, 15]. Therefore, the principal objective of pharmacogenomics is to predict an individualized drug treatment based on each person's genetic profile based on a genome study [16].

Likewise, it is necessary to distinguish between pharmacogenetics and pharmacogenomics. The first refers to interindividual variability in drugs under genetic factors' influence. The second concept encompasses the interindividual variability of such drugs under the influence of certain gene expressions, which may also be fundamental in the physiopathological manifestation [17].

The historical pharmacogenomics bases arise from James Watson and Francis Crick's elucidation of DNA structure in the 1950s [18]. This finding laid the foundations of modern molecular biology and the branches of science associated with genetic material study.

Subsequently, another landmark event was the Human Genome Project in the 2000s. The aim was to complete a mapping of human genes to obtain their functional, physical, and chemical composition [19].

In this way, in the post-genomic era, research began with the contribution of omics. These sciences allow the study of diverse molecules participating in the organism's functions. They include proteomics, metabolomics, and genomics. Proteomics studies many proteins in a sample, covering their functions, post-translational modifications, and interactions with other proteins or substances [20].

Metabolomics analyzes the concentration changes of specific metabolites according to the body's responses to genetic variation. It also allows for obtaining a sample metabolic profile, both qualitatively and quantitatively [20].

Finally, genomics is dedicated to investigating the complete genome. It allows studies of gene identification, characterization, interaction, and function [20].

Thanks to human genomics studies, interest in pharmacogenomics awoke to explain the illness from a genetic basis. Besides, it intends to search for new therapeutic targets and drugs [19].

Pharmacogenomics has two clinical methodologies: candidate gene analysis and genome-wide association studies. The first strategy is identifying the genes responsible for drug metabolism, transport, or therapeutic targets and ruling out possible polymorphisms that affect these processes. Regarding the second, a comparison is executed of the total genomic profile of two groups of patients phenotypically classified into the one that receives the drug and the one to which the placebo is administered. Both are genotyped, and the frequencies are compared to uncover polymorphisms affecting pharmacological response [21].

The candidate gene analysis has made it possible to understand the pathologies. In an investigation, genetic mapping was done to identify a mutation that causes autoimmune myasthenia gravis in seven family members: four affected, two unaffected, and one with an uncertain diagnosis. The results indicated that a sequence variant in the ENOX1 gene is related to the probability of suffering from it in said individuals [22].

Furthermore, genome-wide association studies have identified genetic variations that influence cancer risk. It has been determined that specific alleles increase the hazard. Many are located in non-coding regions, which influence gene expression. Such information has led to the discovery of new drugs and the repositioning of existing ones [23].

3. PHARMACOGENOMICS IMPORTANCE

Pharmacogenomics is gaining relevance in pharmacological research due to non-invasive genetic engineering and molecular biology techniques, recombinant DNA, and the need to explain drug response variations [16, 24]. Researchers aim to find biomarkers associated with the diagnosis, prognosis, and treatment response [24].

The identification of genes with polymorphic variants has had a significant clinical impact. An example is warfarin, an anticoagulant indicated to prevent and treat thromboembolic events. Its correct dosage is difficult because of the narrow therapeutic index, causing bleeding and response variability in patients [25]. The polymorphism in the CYP2C9 and vitamin K epoxide reductase complex subunit 1 (VKORC1) genes influences their therapeutic effects, demonstrating that they are associated with an increased risk of excessive anticoagulation and bleeding events [25, 26].

The major histocompatibility complex (MHC) comprises a group of cell surface proteins that bind to foreign molecules to be recognized by the corresponding T lymphocytes, thus inducing an immune response. This protein structure is the human leukocyte antigen (HLA) [27].

Variation in HLA genes defines susceptibility to autoimmune diseases, infections, and immune reactions [27]. HLA-B*57:01 gene carriers have a high-risk factor for hypersensitivity to abacavir [24, 26, 28]. It is a first-line antiretroviral, classified as a nucleoside reverse transcriptase inhibitor, against human immunodeficiency virus (HIV), capable of terminating DNA synthesis and, therefore, viral replication [28]. Its administration in this group can trigger Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) [29].

Both are hypersensitivity reactions (HSR), which are excessive or inappropriate immune responses produced by the immune system to a self or foreign antigen, damaging body tissues. According to the mechanism of tissue injury, HSRs are divided into four types [30, 31], as shown in **Table 1**.

Table 1. Types of hypersensitivities and their main characteristics [30, 31].

Hypersensitivity type	Characteristics
Type 1 HSP	-It is an immediate immune response mediated by specific immunoglobulin E (IgE) against the triggering agent. -This reaction can be acute, resulting in severe systemic complications, or chronic, comprising recurrent processes, but not so severe.
Type 2 HSP	-It is mediated by IgM and IgG antibodies, which use cell surface antigens. -Cytotoxic reactions that produce cell destruction, opsonization, and phagocytosis are developed.
Type 3 HSP	-IgM and IgG antibodies bind to the antigen, forming immune complexes. -The complexes are deposited in various tissues, causing an inflammatory reaction.
Type 4 HSP	-Delayed type reactions are observed (the response is reflected 48 to 72 hours after the immune reaction). -They are mediated by T cells.

SJS and TEN are considered type 4 HSR to drugs and their metabolites. They represent distinct grades of the same severe skin adverse reaction, characterized by widespread keratinocyte death. It results in the denudation of the skin and mucosa to total thickness, making the person susceptible to sepsis. Mortality is around 30 % [32, 33, 34].

Given this situation, the United States Food and Drug Administration (FDA) established the mandatory exam for HLA-B*57:01 in all people to be treated with abacavir. Similarly, it indicated its prohibition in those with positive results for this genotype [35].

The same happens with carbamazepine, a tricyclic antidepressant against epilepsy and other neurological and psychiatric disorders. Its mechanism is unknown but associated with binding to sodium channels and interactions with calcium channels. Moreover, it potentiates the inhibitory effects of gamma-aminobutyric acid (GABA) and decreases the glutamate excitatory [36].

The presence of the HLA-B*15:02 allele is associated with an augmented risk of HSRs with severe cutaneous involvement, including SJS and TEN, particularly in Asian persons [37, 38]. The FDA changed the drug's label, incorporating information on allele genomics as a marker, and recommended genotyping in this population [37].

In transplant patient care, polymorphisms in the ABCB1, CYP3A4, and CYP3A5 genes are recognized and considered to optimize tacrolimus therapy [39]. This therapeutic option has favored patient survival after transplantation. It forms a complex with the immunophilin FK binding protein 12 (FKBP-12), interfering with the transduction pathway of the intracellular calcium-dependent signal essential for T lymphocyte activation [40]. Its indication has spread as a first-line option for intestine, kidney, heart, lung, and bone marrow transplantation [41] since adequate immunosuppressive therapy is essential to maintain high allograft viability and prevent acute rejection [39]. Nevertheless, genetic factors influence absorption, metabolism, excretion, and drug response [41].

Because of these findings, the FDA recognized that pharmacogenomics plays a critical role. Now, pharmaceutical products can contain information on genetic biomarkers [39]. Labeling aims to provide information to patients and health personnel on the proper use of the product and to facilitate its understanding. It is a tool in health education and is referenced every day in practice by health professionals [42].

An extensive list of therapeutic medications with pharmacogenomic information on their respective biomarkers has been created [43]. Thus, this science has found an application field in therapeutic areas such as psychiatry [16, 24], analgesia [24], neurology [16, 24, 45, 46], oncology [16, 46], and cardiology [16, 24, 47].

4. FACTORS AFFECTING PHARMACOGENOMICS

Pharmacogenomics involves diverse factors. They constitute a linked network, which must be analyzed and evaluated for the drug response. They can be divided into interindividual variability, pharmacokinetics variability, pharmacodynamic variability, and biomarkers. Each is detailed below.

4.1 Interindividual variability

Everyone is a complex organism involving factors and aspects of relevance in clinical practice. Each individual responds singularly to various treatments or the same disease. This differentiated response is mainly the product of intrinsic drug, genetic, epigenetic, environmental, and personal factors. Together, they affect the proteins that metabolize or transport substances, their therapeutic targets (receptors), and variations in relevant metabolic enzymes. The result is the influence on efficacy and pharmacological safety. Nonetheless, the relevance or contribution of each one varies for the substance under study [24].

According to the above, there are factors related to the drug. Some are physicochemical characteristics, administration routes, dosage, excipients, and interactions with other medicines [24, 48].

Concerning genetic factors, metabolic enzymatic activity, transporters' efficiency, and some receptors' sensitivity must be considered [24, 49, 50]. The interindividual variability in the expression of the genes of drug-metabolizing enzymes such as cytochromes (CYPs) and some transporters is more remarkable than other types [51].

Another component is epigenetic factors. One of them involved histones, proteins that, when undergoing modifications such as methylation or acetylation, favor the alteration of gene expression [52].

Expression levels are associated in the same way with micro-RNAs (of ribonucleic acid) since their role in regulating the expression of drug metabolism and transport genes has been identified [53]. As a complement, DNA alterations such as methylation in regions rich in guanine and cytosine are frequently found in the gene promoter region, causing its silencing [54].

Likewise, environmental aspects affect the differentiated response between patients. They include smoking, alcohol consumption, and diet [24].

Finally, the elements of the person must be pondered. They incorporate age, sex, pregnancy, lactation, renal and hepatic functions, and pathologies [24]. A decline of up to 3.5 % in the ratio of CYP enzymes has been seen for each life decade, as well as a reduction in renal function. Additionally, delayed gastric and colonic emptying has been reported in women, together with a higher stomach pH [55]. Together, these characteristics significantly vary the pharmaceutical product response.

4.2 Pharmacokinetics and pharmacodynamic variability

The genetic variability of populations and individuals significantly affects other pharmaceutical processes in the human body, such as pharmacokinetics and pharmacodynamics [56]. These sciences oversee studying the interactions and mechanisms of action of drugs within the organism [57]. Regarding pharmacokinetics, the variability of the drug absorption, distribution, metabolism, and excretion (ADME) process is being investigated [58].

Metabolism is interesting because polymorphisms in the genotypic sequence encoding for metabolizing enzymes or transporter proteins can cause unequal therapeutic responses among individuals. When this happens, problems in the safety and efficacy of pharmaceuticals can arise [57, 59].

The genetic variants involved in pharmacogenomics can be somatic or germinal. The somatic ones occur in the individual after birth and are not hereditary. They optimize the most effective therapeutic choice for the patient [57, 60]. In contrast, the germ cells are hereditary, acquired from the parents, with the consequent progeny transmission. They make it possible to predict therapeutic efficacy and toxicity [57, 60]. These variants are notable because the genes have all the information to synthesize proteins, and, in case of changes in the sequences, they can decrease or increase their function, such as drug transport or metabolism [57, 61].

Pharmacogenomics aims to create a relationship with pharmacokinetics for drug dosage personalization. This synergy seeks to improve its benefits and reduce consumption risks [60].

The CYP450 family is among the metabolizing enzymes responsible for 90 % of the hepatic metabolism of endogenous and exogenous substances. Thanks to pharmacogenomic studies, four metabolic phenotypes responsible for the medicine concentration in circulation have been identified [62], detailed in **Table 2**. This classification has helped to identify people who suffer from substance poisoning, even when administered at regular doses [63].

Table 2. Definition of drug metabolic phenotypes [62, 64].

Type of metabolizer	Definition
Ultrarapid	Due to two active copies of the CYP450 gene, it has a high metabolizing capacity. Therefore, the minimum effective concentration and therapeutic response are never reached.
Rapid or normal	It has two phenotypically active alleles, one active and one partially defective, or a null allele and a duplication. It has an adequate response because it reaches the desired concentrations.
Intermediate	It has two alleles in diverse combinations: defective-defective, defective-null, or active-null. The defective one causes the enzyme to reduce its metabolizing function, and the null one does not work. These conditions reduce metabolic capacity, and the plasmatic concentrations slightly exceed the minimum toxic concentration, adversely affecting the organism.
Poor	It has both null alleles or a deleted gene (loss of one or more nucleotides of the genetic material). Its metabolic capacity is nil, and the plasmatic concentrations greatly exceed the minimum toxic concentration, predisposing it to side effects on health by drug accumulation.

For its part, pharmacodynamic variability depends on how the drug-receptor interaction is carried out. Therefore, genes that encode for receptors and functional proteins related to pharmacological actions after receptor binding are studied. If a subject has polymorphisms, the therapeutic response may be improved by encoding more therapeutic targets or diminished by encoding less. In addition, studies indicate that there are more receptors than usual, and there is a greater possibility of developing adverse reactions. The drawback is that the phenotypic manifestations of polymorphisms in genes related to pharmacodynamics are only observed when the patient is exposed to the substance [13, 62].

4.3 Biomarkers

Pharmacogenetic or pharmacogenomic biomarkers are genes responsible for encoding proteins involved in the action of drugs, either by participating in the mechanism of action or their metabolism, often related to pharmacological toxicity. Its variations have allowed the description of interindividual and interpopulation responses, transforming as a vital tool in implementing personalized medicine to improve the pharmacological response and minimize unwanted effects [16].

Genetic variation is the product of structural differences in particular genes or variations in portions with regulatory activity, the product of insertions, deletions, duplications, inversions, or substitutions of nitrogenous bases modifying the DNA sequence. This situation generates a change in the gene functionality, with the consequent synthesis of modified proteins [65].

Each gene occupies a characteristic location on a specific chromosome and has components necessary for transcription regulation. They include exons, introns, promoter sequences, and regulatory regions [66].

The exons are segments with information that encodes the macromolecule, while the introns comprise sections that do not appear in the mature mRNA because they are removed. Regulatory regions can silence or amplify gene expression, and the promoter indicates the transcription initiation [66].

Polymorphisms can occur as single nucleotide polymorphisms (SNPs), representing about 90 % of all variations found in the human genome. Other more complex differences are copy number variations (CNVs) [65, 67].

SNPs occur when a paired base pair is substituted within the nucleotide sequence. If the variation is in a regulatory area or gene, it can alter the encoded protein synthesis [65].

For its part, CNV occurs when DNA sections are repeated or deleted several times. As with SNPs, they become transcendental when they affect regulatory or encoding regions. The result could be increased protein synthesis, which could lead to an augmentation in enzymatic activity, with the consequent affectation of drug metabolism and alteration of pharmacokinetic parameters. On the contrary, it could cause gene deletion, and since there are fewer protein copies, the enzymatic activity could be diminished [68].

Studies have indicated that allelic variations of pharmacogenetic biomarkers occur with different population frequencies according to ethnicity, age, and gender. Such research has generated a more personalized medicine capable of tailoring treatment to individual needs and patients' characteristics. A current advantage is the technological improvements in determining pharmacogenetic biomarkers [69, 70, 71, 72]. **Table 3** lists some of those studied together with drugs associated with each other and the clinical area where these medications are used.

Table 3. Pharmacogenomic biomarkers associated with different drugs utilized in clinical practice for distinct therapeutic areas.

Biomarker	Drug	Therapeutic area
CYP2C19	Formoterol [73]	Pulmonary
	Carisoprodol [74]	Rheumatology
	Citalopram [74]	Psychiatry
	Clopidogrel [74]	Cardiology
	Diazepam [74]	Neurology
	Voriconazole [74]	Infectious diseases
CYP2C9	Celecoxib [75]	Rheumatology
	Meloxicam [76]	Anesthesiology
	Phenytoin [76]	Neurology
	Piroxicam [76]	Rheumatology
	Warfarin [77]	Hematology
	CYP2D6	Tamoxifen [78]
Venlafaxine [79]		Psychiatry
Tramadol [80]		Anesthesiology
Propranolol [81]		Cardiology
TPMT		Azathioprine [82]
	Cisplatin [83]	Oncology
	Mercaptopurine [84]	Oncology
	Tioguanine [84]	Oncology

VKORC1	Warfarin [77]	Hematology
CYP1A2	Rucaparib [85]	Oncology
UGT1A6	Valproic acid [86]	Neurology
EGFR	Cetuximab [87]	Oncology
HER2	Trastuzumab [88]	Oncology
HLA-B	Carbamazepine [89]	Neurology

5. CLINICAL APPLICATION OF PHARMACOGENOMICS

Table 3 illustrates the importance and variety of application areas of biomarkers in understanding the response of numerous drugs on the market. This information has caused important official agencies such as the FDA to show interest in data collection that serves as a guide and highlights the importance of biomarker analysis prior to decision-making associated with the therapeutic indication and its respective justification [90, 91]. Some of these areas are explained in more detail.

5.1 Immunological disorders

HSRs and autoimmune problems affect a significant percentage of the world's population. HSRs occur from an exacerbated immune response against allergens from the environment, while autoimmune pathologies develop when the immune system triggers a response against autoantigens. Both arise from a loss of clinical tolerance and reaction against innocuous foreign or self-antigens. For this reason, at the therapeutic level, the aim is to broaden the activation threshold of the cells and to modify the function of antigenic memory T and B lymphocytes to restore tolerance to some degree and avoid symptoms for an extended period [92].

Conventional therapy included nonsteroidal anti-inflammatory drugs, corticosteroids, and immunosuppressants. These molecules are non-specific and have serious adverse effects, which can even worsen the patient's health [93]. For example, glucocorticoids can provoke Cushing's syndrome by increasing cortisol in the blood. Its manifestations comprise edema, weight gain, and striae [94].

These drawbacks changed with the advent of antigen-specific immunotherapy. It involves drugs capable of decreasing the immune response against a particular antigen or a set of antigens associated with an immune disorder [95].

This therapy has shown promising results, and its effectiveness against rhinitis, conjunctivitis, and asthma has been appreciated. Gradually increasing amounts of the allergen responsible for the clinical picture are administered subcutaneously until the patient reaches a dose that promotes immunological tolerance against this element [96].

In addition, genetic factors associated with HLA confer a greater risk of autoimmune liver affectations. These molecules induce and regulate the immune response through antigen presentation and T-cell recognition. For this reason, a therapy capable of promoting tolerance through the **gradual administration of an antigenic peptide of synthetic origin is being studied** [97].

Antigen-specific immunotherapy can change the natural course of immunological illnesses. Moreover, pharmacogenomics can monitor clinical evolution and predict treatment-effective responses, select patients who respond to them, facilitate the study of drug combinations, investigate optimal doses, and discover the mechanism of action in detail [92].

Therefore, biological therapy, targeting immune system components, is attempted as a better replacement. The problem is that not all patients respond adequately. Some do not have a response or lose it after having responded [93].

Therefore, pharmacogenomics locates genetic determinants involved with the efficacy or toxicity of the therapeutic options available against these conditions, making finding the best therapeutic route possible. One of the investigations regarding genetic polymorphisms and the clinical response to treatment of systemic lupus erythematosus (characterized by the generation of antibodies against organs and tissues) revealed their presence in genes concerned with pharmacokinetics such as CYP450, the organic-anion transporter polypeptide (OATP), and ABC transporters, and pharmacodynamics, specifically the Fc gamma receptor (FcγR) and toll-like receptors (TLR) [98].

These variations affect the therapeutic target and the patient's sensitivity to the drugs, respectively. For this reason, its knowledge is key to reducing the risk of suffering secondary reactions and even favoring the treatment efficacy [98].

On the other side, identifying the genes corresponding to disease susceptibility, therapeutic targets, drug metabolism, and genetic predictors with similar pathogenic signaling pathways are excellent candidates for improving therapeutics [93]. One case refers to the expectation of the pharmacological response against tumor necrosis factor alpha (TNF- α) through evaluating the genetic variants of its signaling pathways [99].

In a bibliographic study, several polymorphisms involved in the mechanism of action of anti-TNF- α therapies for rheumatoid arthritis were described, mainly six associated with infliximab, five with etanercept, and three with adalimumab. They are responsible for preventing the blockade of the interaction between TNF- α and the binding sites of its cell surface receptors (p55 and p75) [100].

Also, it is possible to identify genetic factors that stimulate the probability of triggering enzyme activity descent, leukopenia development, and HSR generation when consuming some anti-inflammatory or immunosuppressive drugs [101]. Even associations with HLA-encoded MHC genes have been accomplished [102].

Candidate gene and genome-wide association studies have been valuable in identifying elements related to allergic reactions' susceptibility to various medications since the HLA genes encode proteins presenting antigens to T lymphocytes, which may be predictors of HSR. It was demonstrated through a study on a population from Taiwan, which experienced carbamazepine-induced SJS. Candidate gene analysis embraced the genotyping of HLA alleles and CYP450 polymorphisms. A strong association of class I allele B*15:02 was specified. For this reason, genotyping is now done in Asian ethnic groups before starting treatment with said medication [103].

Despite the above, more information must be available for personalized therapy against autoimmune diseases and HSR. The cumulative knowledge of physiopathology, their better-defined classifications, and the identification of autoantigens, together with their corresponding antibodies, lead to the generation of increasingly safe and effective therapeutic strategies [93, 104].

5.2 Neurological diseases

They affect the body's autonomic, peripheral, and central nervous systems. The most common are epilepsy, Alzheimer's, and Parkinson's. However, others, such as migraines, can be located within this area [105]. According to data from the World Health Organization (WHO) for 2019, Alzheimer's and other dementias were the second cause of death for high-income nations, responsible for 814,000 deaths [106].

On the other part, according to the Centers for Disease Control and Prevention (CDC), in 2021, 119,399 deaths from Alzheimer's were reported in the United States [107]. Regarding the American region, the Pan American Health Organization (PAHO) reported 533,172 deaths from neurological disorders throughout 2019 [105].

Epilepsy is characterized by recurrent and unpredictable interruptions of normal brain function called epileptic seizures. Brain dysfunction may result from distinct causes [108].

Genetic polymorphisms present in phase I biotransformation enzymes, specifically CYP2C9, CYP2C19, and CYP3A4/3A5, and phase II (uridine diphosphate glucuronyltransferase or UDP-UGT), are of clinical relevance for its treatment. Phenytoin, one of the primary drugs available for epileptic seizures, is metabolized mainly by CYP2C9 (about 90 %) and a little by CYP2C19. If there is function loss in some variants of the alleles that encode for these enzymes, there is an elimination decrease, with the consequent rise in blood and a greater neurotoxicity risk. More than 50 variants have been identified in the gene that encodes for CYP2C9, confirming its high polymorphism [44, 109]. Hence, their role is to identify necessary adaptations regarding their employment.

As a complement, in an observational study with 23 drug-resistant epilepsy patients, a genomic DNA analysis was performed. It was identified that the relevant SNPs were CYP2D6*2, CYP2D6*4, CYP2C19*2, and CYP3A4*1B, which were related to poor metabolizers. These variations can affect the response to antiepileptic drugs and generate therapy resistance [110].

In another study involving 98 children with epilepsy in China, a relationship was found between the valproic acid plasma concentration and SNPs involving UDP-UGT, specifically UGT1A6 and UGT2B7, and CYP2C9. Individuals with specific polymorphisms in UGT1A6 exhibited lower plasma concentrations and required higher doses. In this way, these mutations affect the metabolism of said substance in epileptic people [111].

On the other hand, Alzheimer's is the most common neurodegenerative disorder and the most common cause of dementia, accounting for approximately half of all cases. The prevalence is approximately 30 % among people 85 years

and older. Its primary clinical manifestation is accelerated cognitive function loss. Alterations in mood and behavior are seen, followed by memory loss, disorientation, and aphasia. The hippocampus and cerebral cortex are the most frequently affected areas [112].

Besides, senile plaques and neurofibrillary tangles are characteristic lesions in affected tissues. Senile plaques in hippocampal blood vessels and neurons mainly comprise $\text{A}\beta$. The protein is produced due to a proteolytic process mediated by α , β , and γ secretases, which break down the amyloid precursor protein (APP) and divide it into components, including amyloid- β [113, 114].

The tangles are filamentous bundles of abnormal tau proteins, which accumulate in the cytoplasm of affected neurons. These macromolecules mainly promote and maintain the structure of microtubules through the C-terminal domain union. Both amyloid- β and tau proteins form insoluble clumps widely associated with this illness [113, 114].

This health problem has an autosomal dominant inheritance pattern. Three mutations in genes that code for proteins involved in amyloid plaque formation (APP, presenilin-1, and presenilin-2) cause early-onset Alzheimer's. A non-familial condition has been associated with the apolipoprotein E $\epsilon 4$ allele (ApoE- $\epsilon 4$) gene. A very low-density lipoprotein transporter gene is required for amyloid- β deposition [112, 115].

In a meta-analysis with 1266 patients with Alzheimer's who were treated with donepezil, polymorphisms in CYP2D6 or ApoE were associated with the effectiveness of treatment. A significant descent in the drug response was achieved for those with variations in both genes [116].

Another study analyzed DNA from the brain region of 71 individuals with the pathology and 81 controls. Variants were searched for different amyloid- β transporters, specifically ABCA1, ABCA7, ABCB1, ABCC2, and ABCG2, some of which were related to the Apo $\epsilon 4$ presence or absence. It was found that an ABCA7 polymorphism is related to the disorder and that ABCB1 is involved in the amyloid- β accumulation process in the brain [117].

Parkinson's (second most common neurodegenerative) is a movement condition whose main clinical characteristics are substantial movement absence, tremors at rest, rigidity, bradykinesia, and postural instability. The cause behind this is a loss of dopamine-producing neurons in the midbrain substantia nigra [112, 118]. Additionally, processes such as altered mitochondrial function and dopamine metabolism, oxidative stress, abnormal protein aggregation, inflammation, necrosis, and accelerated apoptosis are presented. Intracellular deposits of α -synuclein, ubiquitin, and other protein aggregates (Lewy bodies) have also been discovered in many neurons and are medical features [112].

The gene that encodes the α -synuclein protein and alterations in dopamine receptors are involved in its early and rapid onset. Likewise, mutations in leucine-rich repeat kinase 2 (LRRK2 or PARK8 gene), parkin-2 (PARK2), and PTEN-induced kinase I (PINK1 or PARK6) have been associated with late-onset, contributing to the augmented risk of illness developing [112, 115, 119].

A study with 199 patients who received levodopa assessed the relationship between dyskinesia and polymorphisms in the DRD2/ANKK1 dopamine receptor gene. The results showed that its variants influence the induction of involuntary movements (dyskinesia) in drug administration [120].

In another study of 228 patients with idiopathic Parkinson's, polymorphisms in the regions for DRD1 and DRD3 were analyzed. A possible relationship between genetic variants and motor complications due to levodopa use was analyzed. It was concluded that a particular polymorphism (DRD1 A48G) may influence dyskinesia [121].

5.3 Cardiovascular pathologies

Cardiovascular events are one of the leading causes of morbidity and mortality worldwide. According to the WHO, around 18 million lives are lost yearly. Even a third of these deaths occur in people under 70 years of age, which is alarming [122].

Another associated problem is that drugs, such as anticoagulants and antiarrhythmics, intended for its treatment have shown highly variable metabolic response ranges because of patients' genetic characteristics. This problem exemplifies the need to find instruments to define adequate doses to maximize drug therapy sensitivity and specificity [123].

Pharmacogenomic studies have focused on existing and commonly employed pharmaceutical products at the clinical level. Guidelines have even been created with therapeutic dosage recommendations based on the genotype of the biomarkers. One of the most studied has been warfarin. It is administered orally as a racemic mixture of S- and R-warfarin

and is indicated to prevent and treat thromboembolic events. Its mechanism of action is based on the VKOR enzyme inhibition, which leads to vitamin K depletion in its reduced form and the synthesis inhibition of the coagulation factors II, VII, IX, and X, and the anticoagulant proteins S and C. The result is a decrease in prothrombin levels and the consequent reduction in the blood clots' thrombogenicity [123, 124, 125].

The interindividual variability of its response has been associated with SNPs, mainly those of the CYP450 family, like CYP2C9, which participates in metabolizing its S enantiomer [123, 124]. The polymorphisms related to these metabolic enzymes promote varied catalytic activity. The allele with total enzymatic activity has been designated CYP2C9*1, and its allelic variants are CYP2C9*2 and CYP2C9*3. They present the arginine change for cysteine at position 144 of the protein (Arg144Cys) and isoleucine for leucine at position 359 (Ile359Leu), respectively. They are the most common in world populations. Thanks to pharmacogenomic studies, it has been established that the *2 allele is linked with a 30 % decrease in enzyme activity. In comparison, the *3 allele is correlated with a reduction of up to 95 % in warfarin metabolism compared to that of CYP2C9*1. Therefore, it is mandatory to make dose adjustments according to the person's metabolizing condition [123, 124, 126].

Furthermore, VKOR can be mentioned as part of drug metabolism pathways. It is characterized by a vital polymorphism (VKORC1) linked with changes regarding the metabolizing phenotype of individuals against said drug [124, 127].

A study evaluated the effects of the presence of CYP2C9 and VKORC1 polymorphisms regarding the cardiovascular patients' sensitivity towards warfarin and the response capacity. Both variables were analyzed during the treatment stabilization phase. Three CYP2C9 and four VKORC1 polymorphisms were studied. The work helped determine that persons with more than one VKORC1 variant were at augmented risk of warfarin sensitivity (excessive anticoagulation) compared to those with one or no polymorphisms. This result was similar to those with more than one CYP2C9 polymorph. They exhibited a higher sensitivity risk to the molecule than persons with one or no polymorphs [124].

Another interesting detail was that volunteers with a single CYP2C9 or VKORC1 polymorph required significantly lower doses than patients without such variants. In summary, the presence of polymorphisms is associated with increased sensitivity to warfarin during therapeutic stabilization [124].

The FDA currently recommends that genetic modifications related to CYP2C9 and VKORC1 be determined before warfarin treatment starts to adjust the dose correctly. Pharmacogenomics' importance is evident in generating more effective and safer therapies and solving patients' problems regarding an adequate response through a suitable and supported dose adjustment [123, 128, 129].

Likewise, clopidogrel belongs to the ticlopidine family and has antiplatelet action. It is indicated for inhibiting blood clot synthesis at the peripheral, coronary, or cerebral arteries. Also, it is a standard treatment for acute coronary syndrome, acute myocardial infarction, and cerebrovascular attacks. It must first be transformed into the active metabolite to exert its therapeutic effect, corresponding to about 5 % of the prodrug. Numerous enzymes, including CYP2C19, mediate this transformation. This metabolite inhibits the adenosine diphosphate receptor P2Y12, expressed on platelets [123, 130, 131].

Through studies, it has been possible to establish polymorphs responsible for interindividual differences. It is estimated that 30 allelic gene variations are responsible for protein synthesis. Some of the most common are CYP2C19*2, CYP2C19*3, CYP2C19*4, and CYP2C19*17 [123, 131, 132].

With CYP2C19*2, a decrease in the metabolite in blood has been reported, reducing antiplatelet activity and promoting the risk of cardiovascular accidents [123]. Besides, CYP2C19*2, CYP2C19*3, and CYP2C19*4 characterize intermediate and poor metabolizers, while CYP2C19*17 carriers are classified as extensive metabolizers [132].

Biochemical drug monitoring tests, such as measuring serum prothrombin concentration or platelet aggregometry tests, are routinely executed. These assays only show the response to the indicated dose but do not function to predict the reaction of a specific patient prior to the treatment beginning, as pharmacogenomic assays can do [123]. Through them, it is possible to find gene polymorphs dependable for the pharmacological action associated with the medication [133].

5.4 Oncology

Cancer involves a complex carcinogenesis process that encompasses genetic and epigenetic factors [49]. Historically, it has been one of the leading death causes. In 2020, there were nearly 10 million fatalities [134]. However, mortality has declined in recent decades thanks to reduced smoking, early diagnosis, minimization of toxicities, and improved treatments through specific biomarkers [135].

Pharmacogenomics is leading in this area, with implications for therapeutic selection, treatment, dosing, and risk prediction [39]. For clinical decision-making, a better understanding of how drugs and adjuvant agents used for this pathology are metabolized is essential [136].

One of the therapies is chemotherapy. It is highly toxic and is usually administered in high doses. The combination of chemotherapeutic agents provokes side effects in 50 % of individuals. In addition, they have a narrow therapeutic index, challenging the management of toxicities [137]. Serious side effects include myelosuppression, renal failure, elevated transaminases, heart failure, tumor lysis syndrome, diarrhea, thrombosis, pulmonary fibrosis, secondary tumors, constipation, pneumonitis, and seizures [135].

Thiopurine S-methyltransferase (TPMT) is a relevant biomarker responsible for 6-mercaptopurine inactivation, an FDA-approved chemotherapeutic to treat leukemia [137]. It inhibits purine synthesis and acts as an antiproliferative agent, interfering with the synthesis of proteins, DNA, and RNA and promoting proliferative T lymphocyte apoptosis [138].

TPMT polymorphisms vary up to ten times the steady-state concentration in patients with the same dose. Those with elevated 6-MP concentrations may induce severe myelosuppression [136]. Treatment in children with acute lymphocytic leukemia depends on the maximum tolerable drug dose [139].

Slow metabolizers can tolerate the full dose for only 7 % of the total treatment time, while intermediate or normal metabolizers do so for 65 and 84 %, respectively. For this reason, the Clinical Pharmacogenetics Implementation Consortium (CIPC) formulated a guideline recommending the usual dose for normal metabolizers, a 30 to 70 % reduction for intermediate metabolizers, and a 90 % reduction for poor metabolizers [139].

Breast cancer is the most common in women and ranks second in mortality. The therapeutic options are surgery, chemotherapy, radiotherapy, and hormonal therapy [140]. The presence of hormone receptors is a determining factor in choosing a therapeutic regimen. Generally, estrogen receptors are expressed in most tumors. Several studies have demonstrated the usefulness of selective modulators of such receptors, such as tamoxifen, as an effective therapy and preventive agent against tumor recurrence [141].

Tamoxifen is the preferred drug for treating estrogen receptor α -positive premenopausal patients [142]. It is a selective estrogen receptor modulator. Its anti-estrogenic effect blocks the hormone action, which stimulates the development of the tumor cells by competing for binding to the estrogen receptor α [142, 143]. The substance requires metabolic activation by the CYP2D6 enzyme, generating the active metabolites 4-hydroxytamoxifen and endoxifen, managers of therapeutic effects [141]. CYP2D6 alleles can confer normal, decreased, or no activity and cause a wide activity range among the population. Such variations are associated with reduced endoxifen metabolite concentration, the product of an enzyme-reduced activity [140].

To relate the predicted CYP2D6 phenotype and serum endoxifen concentrations with disease-free survival, a multicenter trial was performed in patients with early breast cancer who received adjuvant tamoxifen. Individuals were genotyped for genetic variants in the CYP2D6 gene and classified as ultrarapid, extensive, intermediate, and poor metabolizers. Through a dose escalation of tamoxifen, there was a significant increase in serum concentrations in poor and intermediate metabolizers. In the case of the slow ones, the endoxifen mean level augmented from 24 to 81 % compared to the mean concentration in extensive metabolizers [144].

For its part, 5-fluorouracil (5-FU) is a prevalent and effective chemotherapeutic agent for treating head and neck, breast, pancreas, and gastrointestinal tract cancer [145]. Furthermore, it is prescribed for advanced colon cancer in adjuvant chemotherapy [146]. Its cytotoxicity mechanism has been attributed to misincorporating its metabolites into RNA and DNA and the thymidylate synthase (TS) enzyme inhibition [145]. This molecule catalyzes the deoxyuridine monophosphate (dUMP) to thymidine monophosphate (dTMP) conversion, one of the nucleotides that form thymine [147]. By its inhibition, there is a drop in DNA replication and repair, with the consequent arrest of tumor cell growth [146]. The most common adverse effects comprise dose-dependent hematologic and gastrointestinal toxicities and skin reactions like hand-foot syndrome [148].

Another important enzyme is dihydropyrimidine dehydrogenase (DPD). Individuals with diminished activity are at high risk of supratherapeutic concentrations with standard dosing since the drug half-life is increased. This situation can lead to severe toxicity [145, 148]. DPD activity is highly heterogeneous in the population, partly attributable to its encoding gene CPYD variability [148].

Given the above, the Netherlands National guideline for colorectal carcinoma recommends DPD testing before therapy to optimize efficacy and avoid side effects [149]. Additionally, two guidelines have been created to help clinicians interpret

DPYD genotypes and adjust the 5-FU dose published by CPIC and the Dutch Pharmacogenetics Working Group (DPWG) [148, 150, 151].

Finally, when put into clinical practice in a schematic and orderly manner, pharmacogenomics represents a central opportunity for advancement in the medical area to improve the health conditions of the entire population. The value given to this branch is reflected in the COVID-19 pandemic. Options have been sought to enhance treatments, elucidating markers and genetic variants of interest in treatments with chloroquine and hydroxychloroquine [152] and other medications considered for this infection [153, 154, 155,156].

6. CONCLUSIONS

Research on the relationships between genomic sciences and pharmacological therapy led to the birth of pharmacogenomics as a pharmaceutical science. Its implementation allows individualizing drug selection and administration to avoid adverse consequences and maximize effectiveness.

This knowledge has a wide range of purposes. Nonetheless, it is necessary to understand better aspects related to interindividual variability, pharmacodynamics, pharmacokinetics, and biomarkers associated with pharmacological treatments.

There are different pathologies where this knowledge has been put into practice, including immunological, neurological, and cardiovascular diseases and cancer. Therefore, it is inevitable that its utilization will increase in the short and medium term, for which the work of each nation's health authorities is indispensable, who would oversee assessing its advantages and disadvantages, always putting the patients' quality of life first.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Herrero Sáenz J. (2019). The Pharmacology of Care: A deductive careological approach from the health and fear paradigm of Avedis Donabedian. *Jan*, 13(4), 1348.
2. De Anca Escudero, A. (2009). The importance of Pharmacy in History. *Cuadernos del Tomás*, 1, 173-191.
3. Valenzuela, M. T. (2020). Importance of vaccines in Public Health: milestones and new challenges. *Revista Médica Clínica Las Condes*, 31(3-4), 233-239.
4. Gil del Valle, L., Rabeiro Martínez, C. L., Gravier Hernández, R., Hernández González-Abreu, M. C., & Bermudez Alfonso, Y. (2017). Current issues on pharmacogenetics and the molecular basis of variable drug response. *Cuban Journal of Pharmacy*, 51(1).
5. Vélez Gómez, S., Torres, I., Manrique, R. D., Duque, M., & Gallo, J. E. (2018). Pharmacogenomic application of the CYP2C19, CYP2C9 and VKORC1 genes involved in the metabolism of clopidogrel and warfarin drugs. *Colombian Journal of Cardiology*, 25(6), 396-404.
6. Brewster, L. M., & Seedat, Y. K. (2013). Why do hypertensive patients of African ancestry respond better to calcium blockers and diuretics than to ACE inhibitors and β -adrenergic blockers? A systematic review. *BMC Medicine*, 11, 141.
7. Weinshilboum, R. M., & Wang, L. (2017). Pharmacogenomics: Precision Medicine and Drug Response. *Mayo Clinic Proceedings*, 92(11), 1711-1722.
8. Kaye, A. D., Mahakian, T., Kaye, A. J., Pham, A. A., Hart, B. M., Gennuso, S., et al. (2018). Pharmacogenomics, Precision Medicine, and Implications on Anesthesia care. *Best Practice & Research Clinical Anaesthesiology*, 32(2), 61-81.
9. Cacabelos, R., Cacabelos, N., & Carril, J. C. (2019). The role of pharmacogenomics in adverse drug reactions. *Expert Review of Clinical Pharmacology*, 12(5), 407-442.
10. Dunnenberger, H. M., Biszewski, M., Bell, G. C., Sereika, A., May, H., Johnson, S. G., et al. (2016). Implementation of a multidisciplinary pharmacogenomics clinic in a community health system. *American Journal of Health-System Pharmacy*, 73(23), 1956-1966.
11. Di Sanzo, M., Cipolloni, L., Borro, M., La Russa, R., Santurro, A., Scopetti, M., et al. (2017). Clinical Applications of Personalized Medicine: A New Paradigm and Challenge. *Current Pharmaceutical Biotechnology*, 18(3), 194-203.
12. Sisodiya, S. M. (2021). Precision medicine and therapies of the future. *Epilepsia*, 62(Suppl 2), S90-S105.
13. Ortiz, L., & Tabak, R. (2012). Pharmacogenomics in Clinical Practice. *Revista Médica Clínica Las Condes*, 23(5), 616-621.
14. Rosero, C. Y., Corredor, M., & Mejía, L. (2017). Polymorphisms in genes involved in the development of gastric cancer: review. *Colombian Journal of Gastroenterology*, 31(4), 391-402.
15. Tafur-Betancourt, L. A., Lema, E., Milla, M. M., Londoño, A., & Navarro Vargas, J. R. (2017). From pharmacokinetics to pharmacodynamics, are we ready for 3D software? *Colombian Journal of Anesthesiology*, 45(4), 335-339.
16. Rodríguez Duque, R., & Miguel Soca, P. E. (2020). Pharmacogenomics: principles and applications in medical practice. *Havana Journal of Medical Sciences*, 19(6), e3128.
17. Prior-González, O. A., Garza-González, E., Fuentes-de la Fuente, H. A., Rodríguez-Leal, C., Maldonado-Garza, H. J., & Bosques-Padilla, F. J. (2011). Pharmacogenetics and its clinical importance: towards a safe and efficient personalized therapy. *University Medicine*, 13(50), 41-49.

18. Watson, J. D., & Crick, F. H. C. (1953). Genetical Implications of the Structure of Deoxyribonucleic Acid. *Nature*, 171, 964-967.
19. Daudén Tello, E. (2006). Pharmacogenetics I. Concept, history, objectives and areas of study. *Actas Dermo-Sifiliográficas*, 97(10), 623-629.
20. Frigolet, M. E., & Gutiérrez-Aguilar, R. (2017). "Omic" sciences, how do they help health sciences? *Revista Digital Universitaria*, 18(7).
21. Pierna Álvarez, M., Marcos-Vadillo, E., García-Berrocal, B., & Isidoro-García, M. (2019). Pharmacogenomics: personalized medicine. *Journal of the Clinical Laboratory*, 12(3), 147-154.
22. Landouré, G., Knight, M. A., Stanescu, H., Taye, A. A., Shi, Y., Diallo, O., et al. (2012). A candidate gene for autoimmune myasthenia gravis. *Neurology*, 79(4), 342-347.
23. Sud, A., Kinnersley, B., & Houlston, R. S. (2017). Genome-wide association studies of cancer: current insights and future perspectives. *Nature Reviews Cancer*, 17(11), 692-704.
24. Quiñones, L., Roco, A., Cayún, J. P., Escalante, P., Miranda, C., Varela, N., et al. (2017). Pharmacogenomics as a fundamental tool for personalized medicine: applications in clinical practice. *Medical Journal of Chile*, 145(4), 483-500.
25. Li, J., Wang, S., Barone, J., & Malone, B. (2009). Warfarin Pharmacogenomics. *P T*, 34(8), 422-427.
26. Relling, M. V., & Evans, W. E. (2015). Pharmacogenomics in the clinic. *Nature*, 526(7573), 343-350.
27. Fan, W. L., Shiao, M. S., Hui, R. C. Y., Su, S. C., Wang, C. W., Chang, Y. C., et al. (2017). HLA Association with Drug-Induced Adverse Reactions. *Journal of Immunology Research*, 2017, 3186328.
28. Rodríguez Carranza, R. (2013). *Vademécum Académico de Medicamentos* (5. ed.). McGraw Hill: Mexico City.
29. Wake, D. T., Smith, D. M., Kazi, S., & Dunnenberger, H. M. (2022). Pharmacogenomic Clinical Decision Support: A Review, How-to Guide, and Future Vision. *Clinical Pharmacology & Therapeutics*, 112(1), 44-57.
30. Vyas, S., Bansal, A., Murugan, N., Bhalla, A. S., Naranje, P., & Manchanda, S. (2023). Hypersensitivity Reactions and the Respiratory System: Imaging Based Review. *Current Problems in Diagnostic Radiology*, 52(1), 56-65.
31. Dispenza, M. C. (2019). Classification of hypersensitivity reactions. *Allergy and Asthma Proceedings*, 40(6), 470-473.
32. Lerch, M., Mainetti, C., Terziroli Beretta-Piccoli, B., & Harr, T. (2018). Current Perspectives on Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis. *Clinical Reviews in Allergy & Immunology*, 54(1), 147-176.
33. Charlton, O. A., Harris, V., Phan, K., Mewton, E., Jackson, C., & Cooper, A. (2020). Toxic Epidermal Necrolysis and Steven-Johnson Syndrome: A Comprehensive Review. *Advances in Wound Care*, 9(7), 426-439.
34. Dodiuk-Gad, R. P., Chung, W. H., Valeyrie-Allanore, L., & Shear, N. H. (2015). Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis: An Update. *American Journal of Clinical Dermatology*, 16(6), 475-493.
35. Roden, D. M., McLeod, H. L., Relling, M. V., Williams, M. S., Mensah, G. A., Peterson, J. F., et al. (2019). Pharmacogenomics. *Lancet*, 394(10197), 521-532.
36. Fricke-Galindo, I., Llerena, A., Jung-Cook, H., & López-López, M. (2018). Carbamazepine adverse drug reactions. *Expert Review of Clinical Pharmacology*, 11(7), 705-718.
37. Klein, M. E., Parvez, M. M., & Shin, J. G. (2017). Clinical Implementation of Pharmacogenomics for Personalized Precision Medicine: Barriers and Solutions. *Journal of Pharmaceutical Sciences*, 106(9), 2368-2379.
38. Smith, T. R., Kearney, E., Hulick, P. J., & Kisor, D. F. (2016). History repeats itself: the family medication history and pharmacogenomics. *Pharmacogenomics*, 17(7), 669-678.

39. García-González, X., Cabaleiro, T., Herrero, M. J., McLeod, H., & López-Fernández, L. A. (2016). Clinical implementation of pharmacogenetics. *Drug Metabolism and Personalized Therapy*, 31(1), 9-16.
40. Provenzani, A., Santeusano, A., Mathis, E., Notarbartolo, M., Labbozzetta, M., Poma, P., et al. (2013). Pharmacogenetic considerations for optimizing tacrolimus dosing in liver and kidney transplant patients. *World Journal of Gastroenterology*, 19(48), 9156-9173.
41. Yu, M., Liu, M., Zhang, W., & Ming, Y. (2018). Pharmacokinetics, Pharmacodynamics and Pharmacogenetics of Tacrolimus in Kidney Transplantation. *Current Drug Metabolism*, 19(6), 513-522.
42. Ramírez-Telles, M., & Argotti-Rodríguez, U. (2022). Regulation of Drug Prescribing Information in Latin America and the Caribbean. *Therapeutic Innovation & Regulatory Science*, 56(4), 536-551.
43. U.S. Food & Drug Administration. (2024). Table of Pharmacogenomic Biomarkers in Drug Labeling. Retrieved from <https://www.fda.gov/drugs/science-and-research-drugs/table-pharmacogenomic-biomarkers-drug-labeling>
44. Balestrini, S., & Sisodiya, S. M. (2018). Pharmacogenomics in epilepsy. *Neuroscience Letters*, 667, 27-39.
45. Gilman, C., McSweeney, C., & Mao, Y. (2014). The applications of pharmacogenomics to neurological disorders. *Current Molecular Medicine*, 14(7), 880-890.
46. Carr, D. F., Turner, R. M., & Pirmohamed, M. (2021). Pharmacogenomics of anticancer drugs: Personalising the choice and dose to manage drug response. *British Journal of Clinical Pharmacology*, 87(2), 237-255.
47. Scibona, P., Angriman, F., Simonovich, V., Heller, M. M., & Bellosso, W. H. (2014). Farmacogenómica cardiovascular. *Archivos de Cardología de México*, 84(1), 25-31.
48. Turner, R. M., Park, B. K., & Pirmohamed, M. (2015). Parsing interindividual drug variability: an emerging role for systems pharmacology. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*, 7(4), 221-241.
49. Soni, N., Soni, N., Maheshwari, R., Thakkar, S., Sharma, D., Tekade, R. K., et al. (2020). Pharmacogenomics and pharmacoepigenomics: Impact on therapeutic strategies. In R. K. Tekade (Ed.), *The Future of Pharmaceutical Product Development and Research*. San Diego: Academic Press.
50. Lewis, L., Crawford, G. E., Furey, T. S., & Rusyn, I. (2017). Genetic and epigenetic determinants of inter-individual variability in responses to toxicants. *Current Opinion in Toxicology*, 6, 50-59.
51. Yang, L., Price, E. T., Chang, C. W., Li, Y., Huang, Y., Guo, L. W., et al. (2013). Gene expression variability in human hepatic drug metabolizing enzymes and transporters. *PLoS One*, 8(4), e60368.
52. Alvarado-González, A., & Arce Jiménez, I. (2013). Mecanismos de acción y resistencia a glucocorticoides en asma y enfermedad pulmonar obstructiva crónica. *Acta Médica Costarricense*, 55(4), 162-168.
53. Swathy, B., & Banerjee, M. (2022). Understanding Pharmaco-Epigenomic Response of Antipsychotic Drugs Using Genome-Wide MicroRNA Expression Profile in Liver Cell Line. *Frontiers in Molecular Neuroscience*, 15, 786632.
54. Ahmed, S. A. H., Ansari, S. A., Mensah-Brown, E. P. K., & Emerald, B. S. (2020). The role of DNA methylation in the pathogenesis of type 2 diabetes mellitus. *Clinical Epigenetics*, 12(1), 104.
55. Trenaman, S. C., Bowles, S. K., Andrew, M. K., & Goralski, K. (2021). The role of sex, age and genetic polymorphisms of CYP enzymes on the pharmacokinetics of anticholinergic drugs. *Pharmacology Research & Perspectives*, 9(3), e00775.
56. Guio, H., Levano, K. S., Sánchez, C., & Tarazona, D. (2015). Role of Pharmacogenomics in the Treatment Regimen of Tuberculosis. *Peruvian Journal of Experimental Medicine and Public Health*, 32(4), 794-800.
57. Llerena, A. (2021). *Pharmacogenomics: The path to personalized treatment*. Roche Institute Foundation: Madrid.

58. Doogue, M. P., & Polasek, T. M. (2013). The ABCD of clinical pharmacokinetics . *Therapeutic Advances in Drug Safety*, 4(1), 5-7.
59. Arrieta-Bolaños, E., Alvarado-Ulate, P., Baudrit-Carrillo, O., & Salazar-Sánchez, L. (2012). Pharmacogenetics: towards individualization of drug therapy in Costa Rica. *Acta Médica Costarricense*, 54(4), 207-216.
60. Moore, L., Cagan, A., Coorens, T. H. H., Neville, M. D. C., Sanghvi, R. , Sanders, M. A., et al. (2021). The mutational landscape of human somatic and germline cells. *Nature*, 597(7876), 381-386.
61. Valenzuela Jiménez, B., González Sales, M., Escudero Ortiz, V., Martínez Navarro, E., Pérez Ruixo, C., Rebollo Liceaga, J., et al. (2013). Influence of genetic polymorphisms in UGT1A1, UGT1A7 and UGT1A9 on the pharmacokinetics of irinotecan, SN-38 and SN-38G. *Farmacia Hospitalaria*, 37(2), 111-127.
62. Belmonte Campillo, C. (2018). Impact of Genetic Polymorphisms in the Pharmacokinetics, Pharmacodynamics and Safety Profile of Aripiprazole [undegraduate thesis]. Madrid: Universidad Autónoma de Madrid.
63. Schulz, M., Iwersen-Bergmann, S., Andresen, H., & Schmoltdt, A. (2012). Therapeutic and toxic blood concentrations of nearly 1,000 drugs and other xenobiotics. *Critical Care*, 16(4).
64. National Human Genome Research Institute. (2024). Deletion. Retrieved from <https://www.genome.gov/genetics-glossary/Deletion>
65. Betcher, H. K., & George, A. L. Jr. (2020). Pharmacogenomics in pregnancy. *Seminars in Perinatology*, 44(3), 151222.
66. Ferrer, J., Maestro, M. A., Fernández-balsells, M. (2009). Genetic Bases of Endocrine Diseases. In M. Pombo, L. Audí, R. Bueno, R. Calzada, F. Cassorla, C. Diéguez et al (Eds.), *Treatise on Pediatric Endocrinology*. Spain: McGraw Hill.
67. Hosseini, S. S., Jebelli, A., Vandghanooni, S., Jahanban-Esfahlan, A., Baradan, B., Amini, M., et al. (2022). Perspectives and trends in advances DNA biosensors for the recognition of single nucleotide polymorphisms. *Chemical Engineering Journal*, 441, 135988.
68. Orrico, K. B. (2019). Basic Concepts in Genetics and Pharmacogenomics for Pharmacists. *Drug Target Insights*, 13, 1177392819886875.
69. Alshabeeb, M. A., Deneer, V. H. M., Khan, A., & Asselbergs, F. W. (2019). Use of Pharmacogenetic Drugs by the Dutch Population. *Frontiers in Genetics*, 10, 567.
70. Díaz Fernández, U., & Rodríguez Ferreiro, A. O. (2016). Applications of biotechnology in the development of personalized medicine. *MEDISAN*, 20(5), 678.
71. Aguirre Fernández, R. E., Serra Valdés, M. A., & Aguirre Posada, M. E. (2017). Holistic vision of new challenges: technological paradigms and bioethical foundations in futuristic medicine. *Havana Journal of Sciences Médicas*, 16(5), 839-849.
72. Litman, T. (2019). Personalized medicine-concepts, technologies, and applications in inflammatory skin diseases. *APMIS*, 127(5), 386-424.
73. Martínez- Matilla, M., Blanco-Verea, A., Santori, M., Ansede-Bermejo, J., Ramos-Luis, E., Gil, R., et al. (2019). Genetic susceptibility in pharmacodynamic and pharmacokinetic pathways underlying drug-induced arrhythmia and sudden unexplained deaths. *Forensic Science International: Genetics*, 42, 203-212.
74. Pratt, V. M., Del Tredici, A. L., Hachad, H., Ji, Y., Kalman, L. V., Scott, S. A., et al. (2018). Recommendations for Clinical CYP2C19 Genotyping Allele Selection: A Report of the Association for Molecular Pathology. *The Journal of Molecular Diagnostics*, 20(3), 269-276.
75. Siu, Y. A., Hao, M. H., Dixit, V., & Lai, W. G. (2018). Celecoxib is a substrate of CYP2D6: Impact on celecoxib metabolism in individuals with CYP2C9*3 variants. *Drug Metabolism and Pharmacokinetics*, 33(5), 219-227.

76. Cottrill, E., Pennington, Z., Ahmed, A. K., Jiang, B., Ehresman, J., Zhu, A., et al. (2021). First Report of Pharmacogenomic Profiling in an Outpatient Spine Setting: Preliminary Results from a Pilot Study. *World Neurosurgery*, 145, e21-e31.
77. Kaye, A. D., Koress, C. M., Novitch, M. B., Jung, J. W., Urits, I., Viswanath, O., et al. (2020). Pharmacogenomics, concepts for the future of perioperative medicine and pain management: A review. *Best Practice & Research Clinical Anaesthesiology*, 34(3), 651-662.
78. Kiyotani, K., Mushiroda, T., Nakamura, Y., & Zembutsu, H. (2012). Pharmacogenomics of Tamoxifen: Roles of Drug Metabolizing Enzymes and Transporters. *Drug Metabolism Pharmacokinetics*, 27(1), 122-131.
79. Ahmed, A. T., Biernacka, J. M., Jenkins, G., Rush, A. J., Shinozaki, G., Veldic, M., et al. (2019). Pharmacokinetic-Pharmacodynamic interaction associated with venlafaxine-XR remission in patients with major depressive disorder with history of citalopram / escitalopram treatment failure. *Journal of Affective Disorders*, 246, 62-68.
80. Zhu, Y., Lopes, G. S., Bielinski, S. J., Borah, B. J., Larson, N. B., Moyer, A. M., et al. (2021). Impact of Pharmacogenomic Information on Values of Care and Quality of Life Associated with Codeine and Tramadol-Related Adverse Drug Events. *Mayo Clinic Proceedings: Innovations, Quality & Outcomes*, 5(1), 35-45.
81. Geng, Y. J., Madonna, R., Hermida, R. C., & Smolensky, M. H. (2021). Pharmacogenomics and circadian rhythms as mediators of cardiovascular drug-drug interactions. *Current Research in Pharmacology and Drug Discovery*, 2, 100025.
82. Taha, N., Hosein, K., Grant-Orser, A., Lin-Shaw, A., & Mura, M. (2021). TPMT and HLA-DQA1-HLA-DRB genetic profiling to guide the use of azathioprine in the treatment of interstitial lung disease: First experience. *Pulmonary Pharmacology & Therapeutics*, 66, 101988.
83. Olgun, Y., Aktaş, S., Altun, Z., Kırkım, G., Kızmaçoğlu, D. Ç., Erçetin, A. P., et al. (2016). Analysis of genetic and non genetic risk factors for cisplatin ototoxicity in pediatric patients. *International Journal of Pediatric Otorhinolaryngology*, 90, 64-69.
84. Strohbuscha, A., & Kator, S. (2020). Pharmacogenomics in Practice: Guidance for Thiopurine Dosing Using Thiopurine Methyltransferase (TPMT) and Nudix Hydrolase 15 (NUDT15). *The Journal for Nurse Practitioners*, 16(3), 238-239.
85. Konecny, G. E., Oza, A. M., Tinker, A. V., Oaknin, A., Shapira-Frommer, R., Ray-Coquard, I., et al. (2021). Population exposure-efficacy and exposure-safety analyses for rucaparib in patients with recurrent ovarian carcinoma from Study 10 and ARIEL2. *Gynecologic Oncology*, 161(3), 668-675.
86. Mani, B., Nair, P. P., Sekhar, A., Kamalanathan, S., Narayan, S. K., & Kesavan, R. (2021). CYP2C19 & UGT1A6 genetic polymorphisms and the impact on Valproic acid-induced weight gain in people with epilepsy: Prospective genetic association study. *Epilepsy Research*, 177, 106786.
87. Hirsch, F. R., Redman, M. W., Moon, J., Agustoni, F., Herbst, R. S., Semrad, T. J., et al. (2022). EGFR High Copy Number Together with High EGFR Protein Expression Predicts Improved Outcome for Cetuximab-based Therapy in Squamous Cell Lung Cancer: Analysis from SWOG S0819, a phase III trial of Chemotherapy with or without Cetuximab in advanced NSCLC. *Clinical Lung Cancer*, 23(1), 60-71.
88. Hertz, D. L., McLeod, H. L., & Hoskins, J. M. (2009). Pharmacogenetics of breast cancer therapies. *Breast*, 18(Suppl 3), S59-S63.
89. Min, F., Fan, C., Zeng, Y., He, N., Zeng, T., Qin, B., et al. (2022). Carbamazepine-modified HLA-A*24:02-bound peptidome: Implication of CORO1A in skin rash. *International Immunopharmacology*, 109, 108804.
90. Hikino, K., Fukunaga, K., & Mushiroda, T. (2018). Gap between the US and Japan in coverage of pharmacogenomic biomarkers by health insurance programs: More coverage is needed in Japan. *Drug Metabolism and Pharmacokinetics*, 33(6), 243-249.

91. Vivot, A., Boutron, I., Ravaud, P., & Porcher, R. (2015). Guidance for pharmacogenomic biomarker testing in labels of FDA-approved drugs. *Genetics in Medicine*, 17(9), 733-738.
92. Mainet-González, D. (2018). La farmacogenómica en el desarrollo de la inmunoterapia específica de alérgenos y autoantígenos. *Bionatura*, 3(3), 673-682.
93. Tavakolpour, S., Darvishi, M., & Ghasemiadl, M. (2018). Pharmacogenetics: A strategy for personalized medicine for autoimmune diseases. *Clinical Genetics*, 93(3), 481-497.
94. Lima-Martínez, M. M., Zerpa, J., Guerrero, Y., Rivera, J., Vielma, M., & Endocrinology Group of Merida (ENDOMER). (2013). Management of Patients with Cushing's Syndrome: Protocol of the Endocrinology Service of the Autonomous Institute of the University Hospital of Los Andes. *Venezuelan Journal of Endocrinology and Metabolism*, 11(3), 147-156.
95. Wraith, D. (2016). Autoimmunity: Antigen-specific immunotherapy. *Nature*, 530(7591), 422-433.
96. Moote, W., Kim, H., & Ellis, A. K. Allergen-specific immunotherapy. (2018). *Allergy Asthma & Clinical Immunology*, 14(Suppl 2), 53.
97. Richardson, N., Ng, S. T. H., & Wraith, D. C. (2020). Antigen-Specific Immunotherapy for Treatment of Autoimmune Liver Diseases. *Frontiers in Immunology*, 11, 1586.
98. Barliana, M. I., Afifah, N. N., Amalia, R., Hamijoyo, L., & Abdulah, R. (2022). Genetic Polymorphisms and the Clinical Response to Systemic Lupus Erythematosus Treatment Towards Personalized Medicine. *Frontiers in Pharmacology*, 13, 820927.
99. Walczak, M., Lykowska-Szuber, L., Plucinska, M., Stawczyk-Eder, K., Zakerska-Banaszak, O., Eder, P., et al. (2020). Is Polymorphism in the Apoptosis and Inflammatory Pathway Genes Associated With a Primary Response to Anti-TNF Therapy in Crohn's Disease Patients? *Frontiers in Pharmacology*, 11, 1207.
100. Puentes Osorio, Y., Amariles, P., Aristizábal Bernal, B. H., Pinto Peñaranda, L. F., & Calleja Hernández, M. A. (2018). Farmacogenómica de etanercept, infliximab, adalimumab y metotrexato en artritis reumatoide. Revisión estructura. *Revista Colombiana de Reumatología*, 25(1), 22-37.
101. Khan, D. A., & Phillips, E. J. (2020). Pharmacogenomic biomarkers in allergy and immunology practice. *Journal of Allergy and Clinical Immunology*, 146(3), 509-512.
102. Sukasem, C., Puangpetch, A., Medhasi, S., & Tassaneeyakul, W. (2014). Pharmacogenomics of drug-induced hypersensitivity reactions: challenges, opportunities and clinical implementation. *Asian Pacific Journal of Allergy and Immunology*, 32, 111-123.
103. Daly, A. K. (2013). Pharmacogenomics of adverse drug reactions. *Genome Medicine*, 5, 5.
104. Arbitrio, M., Scionti, F., Di Martino, M. T., Caracciolo, D., Pensabene, L., Tassone, P., et al. (2021). Pharmacogenomics Biomarker Discovery and Validation for Translation in Clinical Practice. *Clinical and Translational Science*, 14(1), 113-119.
105. Pan American Health Organization. (2021). Burden of Neurological Conditions. Retrieved from <https://www.paho.org/en/enlace/burden-neurological-condition>
106. World Health Organization. (2024). The top 10 causes of death. Retrieved from <https://www.cdc.gov/nchs/fastats/deaths.htm>
107. National Center of Health Statistics. (2020). Deaths and Mortality. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>
108. Fisher, R. S., van Emde Boas, W., Blume, W., Elger, C., Genton, P., Lee, P., et al. (2005). Epileptic Seizures and Epilepsy: Definitions Proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia*, 46(4), 470-472.

109. Božina, N., Sporiš, I. Š., Božina, T., Klarica-Domjanović, I., Tvrdeić, A., & Sporiš, D. (2019). Pharmacogenetics and the treatment of epilepsy: what do we know? *Pharmacogenomics*, 20(15), 1093-1101.
110. López-García, M. A., Feria-Romero, I. A., Serrano, H., Rayo-Mares, D., Fagiolino, P., Vázquez, M., et al. (2017). Influence of genetic variants of CYP2D6, CYP2C9, CYP2C19 and CYP3A4 on antiepileptic drug metabolism in pediatric patients with refractory epilepsy. *Pharmacological Reports*, 69(3), 504-511.
111. Guo, Y., Hu, C., He, X., Qiu, F., & Zhao, L. (2012). Effects of UGT1A6, UGT2B7, and CYP2C9 genotypes on plasma concentrations of valproic acid in Chinese children with epilepsy. *Drug Metabolism and Pharmacokinetics*, 27(5), 536-542.
112. Checkoway, H., Lundin, J. I., & Kelada, S. N. (2011). *Neurodegenerative diseases*. IARC Scientific Publications, (163), 407-419.
113. Carvajal Carvajal, C. (2016). *Biología Molecular de la Enfermedad de Alzheimer*. *Medicina Legal de Costa Rica*, 33(2).
114. López-Camacho, P. Y., Guzmán Hernández, R. N. H., Hernández González, V. H., Díaz Muñoz, J. E., García-Sierra, F., & Basurto-Islas, G. (2017). Investigación y terapias en la enfermedad de Alzheimer basadas en beta amiloide y tau. *Archivos de Neurociencias*, 22(2), 72-88.
115. Cacabelos, R. (2020). Pharmacogenomics of Alzheimer's and Parkinson's diseases. *Neuroscience Letters*, 726, 133807.
116. Xiao, T., Jiao, B., Zhang, W., Tang, B., & Shen, L. (2016). Effect of the CYP2D6 and APOE Polymorphisms on the Efficacy of Donepezil in Patients with Alzheimer's Disease: A Systematic Review and Meta-Analysis. *CNS Drugs*, 30(10), 899-907.
117. Cascorbi, I., Flüh, C., Remmler, C., Haenisch, S., Faltraco, F., & Grumbt, M., et al. (2013). Association of ATP-binding cassette transporter variants with the risk of Alzheimer's disease. *Pharmacogenomics*, 14(5), 485-494.
118. Hurtado, F., Cardenas, M. A., Cardenas, F., & León, L. A. (2016). La Enfermedad de Parkinson: Etiología, Tratamiento y Factores Preventivos. *Universitas Psychologica*, 15(5).
119. Payami, H. (2017). The Emerging Science of Precision Medicine and Pharmacogenomics for Parkinson's Disease. *Movement Disorders*, 32(8), 1139-1146.
120. Rieck, M., Schumacher-Schuh, A. F., Altmann, V., Francisconi, C. L. M., Fagundes, P. T. B., Monte, T. L., et al. (2012). DRD2 haplotype is associated with dyskinesia induced by levodopa therapy in Parkinson's disease patients. *Pharmacogenomics*, 13(15), 1701-1710.
121. Dos Santos, E. U. D., Duarte, E. B. C., Miranda, L. M. R., Asano, A. G. C., Asano, N. M. J., Maia, M. M. D., et al. (2019). Influence of DRD1 and DRD3 Polymorphisms in the Occurrence of Motor Effects in Patients with Sporadic Parkinson's Disease. *NeuroMolecular Medicine*, 21(3), 295-302.
122. World Health Organization. (2024). Cardiovascular diseases. Retrieved from https://www.who.int/health-topics/cardiovascular-diseases#tab=tab_1
123. Vélez Gómez, S., Torres, I., Manrique, R. D., Duque, M., & Gallo, J. A. (2018). Pharmacogenomic application of CYP2C19, CYP2C9 and VKORC1 genes involved in the metabolism of clopidogrel and warfarin drugs. *Colombian Journal of Cardiology*, 25(6), 396-404.
124. Al-Eitan, L. N., Almasri, A. Y., & Khasawneh, R. H. (2019). Effects of CYP2C9 and VKORC1 polymorphisms on warfarin sensitivity and responsiveness during the stabilization phase of therapy. *Saudi Pharmaceutical Journal*, 27(4), 484-490.
125. Pratt, V. M., Cavallari, L. H., Del Tredici, A. L., Hachad, H., Ji, Y., Kalman, L. W., et al. (2020). Recommendations for Clinical Warfarin Genotyping Allele Selection: A Report of the Association for Molecular Pathology and the College of American Pathologists. *The Journal of Molecular Diagnostics*, 22(7), 847-859.

126. Kuzikov, A. V., Filippova, T. A., Masamrekh, R. A., & Shumyantseva, V. V. (2022). Electrochemical determination of (S)-7-hydroxywarfarin for analysis of CYP2C9 catalytic activity. *Journal of Electroanalytical Chemistry*, 904, 115937.
127. Li, S., Liu, S., Liu, X. R., Zhang, M. M., & Li, W. (2020). Competitive tight-binding inhibition of VKORC1 underlies warfarin dosage variation and antidotal efficacy. *Blood Advances*, 4(10), 2202-2212.
128. Al-Saikhan, F. I. (2020). Genetic risk assessment towards warfarin application: Saudi Arabia study with a potential to predict and prevent side effects. *Saudi Journal of Biological Sciences*, 27(1), 456-459.
129. Zhu, Y., Swanson, K. M., Rojas, R. L., Wang, Z., St Sauver, J. L., Visscher, S. L., et al. (2020). Systematic review of the evidence on the cost-effectiveness of pharmacogenomics-guided treatment for cardiovascular diseases. *Genetics in Medicine*, 22(3), 475-486.
130. Kuszynski, D. S., Christian, B. D., Dorrance, A. M., & Lauver, D. A. (2021). Clopidogrel Treatment Inhibits P2Y2-Mediated Constriction in the Rabbit Middle Cerebral Artery. *European Journal of Pharmacology*, 911, 174545.
131. Kuo, F. Y., Lee, J. H., Lan, W. R., Su, J. H., Lee, W. L., Wang, Y. C., et al. (2022). Effect of CYP2C19 status on platelet reactivity in Taiwanese acute coronary syndrome patients switching to prasugrel from clopidogrel: Switch Study. *Journal of the Formosan Medical Association*, 121(9), 1786-1797.
132. Montazid, M. S., Sajib, A. A., Hassan, K. N., Khaleque, A., Rahman, M., Sufian, A., et al. (2021). Multiplex allele-specific detection of clinically important CYP2C19 variants associated with clopidogrel metabolism in a Bangladeshi population sample. *Meta Gene*, 27, 100830.
133. Pratt, V. M., Turner, A., Broeckel, U., Dawson, D. B., Gaedigk, A., Lynnes, T. J., et al. (2021). Characterization of Reference Materials with an Association for Molecular Pathology Pharmacogenetics Working Group Tier 2 Status: CYP2C9, CYP2C19, VKORC1, CYP2C Cluster Variant, and GGCX: A GeT-RM Collaborative Project. *The Journal of Molecular Diagnostics*, 23(8), 952-958.
134. Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., et al. (2021). Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA: A Cancer Journal for Clinicians*, 71(3), 209-249.
135. Zhang, Y., Somtakoune, S. D., Cheung, C., Listiawan, M., & Feng, X. (2016). Therapeutic Application of Pharmacogenomics in Oncology. *AAPS J*, 18(4), 819-829.
136. Patel, J. N., Olver, I., & Ashbury, F. (2021). Pharmacogenomics in cancer supportive care: key issues and future directions. *Support Care Cancer*, 29(11), 6187-6191.
137. Elzagallaai, A. A., Carleton, B. C., & Rieder, M. J. (2021). Pharmacogenomics in Pediatric Oncology: Mitigating Adverse Drug Reactions While Preserving Efficacy. *Annual Review of Pharmacology and Toxicology*, 61, 679-699.
138. Fernández-Ramos, A. A., Marchetti-Laurent, C., Poindessous, V., Antonio, S., Laurent-Puig, P., Bortoli, S., et al. (2017). 6-mercaptopurine promotes energetic failure in proliferating T cells. *Oncotarget*, 8(26), 43048-43060.
139. Mlakar, V., Huezo-Díaz Curtis, P., Satyanarayana Upgunduri, J. R., Krajinovic, M., & Ansari, M. (2016). Pharmacogenomics in Pediatric Oncology: Review of Gene-Drug Associations for Clinical Use. *International Journal of Molecular Sciences*, 17(9), 1502.
140. Reis, S. S., Carvalho, A. S., & Fernandes, R. (2019). Pharmacogenomics, CYP2D6, and Tamoxifen: A Survey of the Reasons Sustaining European Clinical Practice Paradigms. *Medicine*, 55(7), 344.
141. Chan, C. W. H., Law, B. M. H., So, W. K. W., Chow, K. M., & Waye, M. M. Y. (2020). Pharmacogenomics of breast cancer: highlighting CYP2D6 and tamoxifen. *Journal of Cancer Research and Clinical Oncology*, 146(6), 1395-1404.
142. Cronin-Fenton, D. P., & Damkier, P. (2018). Tamoxifen and CYP2D6: A Controversy in Pharmacogenetics. *Advances in Pharmacology*, 83, 65-91.

143. Ariza Márquez, Y. V., Briceño Balcázar, I., & Ancízar Aristizábal, F. (2016). Breast cancer treatment and pharmacogenetics. *Revista Colombiana Biotecnología*, 28(1), 121-134.
144. Dezentjé, V. O., Opdam, F. L., Gelderblom, H., Hartigh den, J., Van der Straaten, T., Vree, R., et al. (2015). CYP2D6 genotype- and endoxifen-guided tamoxifen dose escalation increases endoxifen serum concentrations without increasing side effects. *Breast Cancer Research and Treatment*, 153(3), 583-590.
145. Matsusaka, S., & Lenz, H. J. (2015). Pharmacogenomics of fluorouracil-based chemotherapy toxicity. *Expert Opinion on Drug Metabolism & Toxicology*, 11(5), 811-821.
146. Xie, P., Mo, J. L., Liu, J. H., Li, X., Tan, L. M., Zhang, W., et al. (2020). Pharmacogenomics of 5-fluorouracil in colorectal cancer: review and update. *Cellular Oncology*, 43(6), 989-1001.
147. Nelson, B., Carter, J. V., Eichenberger, M. R., Netz, U., & Galandiuk, S. (2016). Genetic polymorphisms in 5-Fluorouracil-related enzymes predict pathologic response after neoadjuvant chemoradiation for rectal cancer. *Surgery*, 160(5), 1326-1332.
148. Hamzic, S., Aebi, S., Joerger, M., Montemurro, M., Ansari, M., Amstutz, U., et al. (2020). Fluoropyrimidine chemotherapy: recommendations for DPYD genotyping and therapeutic drug monitoring of the Swiss Group of Pharmacogenomics and Personalised Therapy. *Swiss Medical Weekly*, 150, w20375.
149. Martens, F. K., Huntjens, D. W., Rigter, T., Bartels, M., Bet, P. M., & Cornel, M. C. (2020). DPD Testing Before Treatment With Fluoropyrimidines in the Amsterdam UMCs: An Evaluation of Current Pharmacogenetic Practice. *Frontiers in Pharmacology*, 10, 1609.
150. Amstutz, U., Henricks, L. M., Offer, S. M., Barbarino, J., Schellens, J. H. M., Swen, J. J., et al. (2018). Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for Dihydropyrimidine Dehydrogenase Genotype and Fluoropyrimidine Dosing: 2017 Update. *Clinical Pharmacology & Therapeutics*, 103(2), 210-216.
151. Lunenburg, C. A. T. C., van der Wouden, C. H., Nijenhuis, M., Crommentuijn-van Rhenen, M. H., de Boer-Veger, N. J., Buunk, A. M., et al. (2020). Dutch Pharmacogenetics Working Group (DPWG) guideline for the gene-drug interaction of DPYD and fluoropyrimidines. *European Journal of Human Genetics*, 28(4), 508-517.
152. Babayeva, M., & Loewy, Z. (2020). Repurposing Drugs for COVID-19: Pharmacokinetics and Pharmacogenomics of Chloroquine and Hydroxychloroquine. *Pharmacogenomics and Personalized Medicine*, 13, 531-542.
153. Al-Taie, A., Büyükc. A. Ş., & Sardas, S. (2022). Considerations into pharmacogenomics of COVID-19 pharmacotherapy: Hope, hype and reality. *Pulmonary Pharmacology & Therapeutics*, 77, 102172.
154. Badary, O. A. (2021). Pharmacogenomics and COVID-19: clinical implications of human genome interactions with repurposed drugs. *The Pharmacogenomics Journal*, 21, 275-284.
155. Stevenson, J. M., Alexander, G. C., Palamuttam, N., & Mehta, H. B. (2021). Projected Utility of Pharmacogenomic Testing Among Individuals Hospitalized With COVID-19: A Retrospective Multicenter Study in the United States. *Clinical and Translational Science*, 14(1), 153-162.
156. EDEKI, CHARLES. 2019. "the emerging focus on pharmacogenomics". *International journal of biochemistry research & review* 27 (2):1-6. [HTTPS://DOI.ORG/10.9734/IJBCRR/2019/V27I230114](https://doi.org/10.9734/IJBCRR/2019/V27I230114).