

PATTERN OF SKIN TUMOURS SEEN IN RIVERS STATE UNIVERSITY TEACHING HOSPITAL
(RSUTH), PORT HARCOURT, NIGERIA:
A 12 YEAR RETROSPECTIVE STUDY

ABSTRACT

- Background:** Cancers are known to be on the increase in Nigeria. This has been attributed to the rise in oil exploration in the nation. The Niger Delta region faces a lot of oil spillage and is highly vulnerable to the effects of this spillage which includes health disorders like cancers.
- Objective:** To document the incidence of skin tumours (**benign and malignant**) over a 12-year period in the Dermatology Out-patient clinic of the Rivers State University Teaching Hospital, Port-Harcourt, Nigeria.
- Materials and Methods :** The data was taken from new patient registers in the dermatology out-patient clinic and analysed.
- Results:** Skin Tumours consists of 5.3% of all dermatological diagnoses within the period. The M: F ratio was 1:1. Benign tumours made up 81.4% of cases and pre-malignant tumours 0.98%. The commonest benign tumour is keloids (28.9%). Malignant skin cancers constituted 17.6% of the cases with Kaposi sarcoma (9.8%), metastatic skin cancer (2.9%) and squamous cell carcinoma (2%) being the commonest skin cancers. The young adult age group (20-44) constituted majority (65.7%) of those affected with skin tumours.
- Conclusion:** There is a relative low prevalence of skin tumours among patients attending the skin clinic in RSUTH (5.3%). The Commonest benign tumor is Keloids while the commonest malignant tumour is Kaposi sarcoma.
- Key words:** Benign, Cancer, Skin, Tumour, RSUTH

Introduction

Skin tumors are abnormal growths on the skin that are caused by various factors such as genetic mutations, viruses, UV radiation, viral infections etc.

Skin tumours can be classified as benign, premalignant and malignant. Benign skin tumors can further be classified into the following:

Epidermal tumors-Seborrheic keratosis, Epidermal nevus, Epidermoid cyst, Viral warts

Dermal tumors-Dermatofibroma, Lipoma, Dermatofibrolipoma,

Viral tumours-Molluscum contagiosum,

Vascular tumors-Haemangioma,, Glomus tumour, Pyogenic granuloma

Hairfollicletumors-Trichoepithelioma, Trichofolliculoma

Sweatglandtumors-Syringoma, Hidroacanthoma

Connectivetissuetumours-Fibroma, Fibrolipoma

Neuraltumours-Neurofibroma, Schwannoma

Pre-malignantskintumours- Actinic keratosis, Keratoacanthoma, Bowen's disease

Malignantskintumours- Squamous cell carcinoma, Basal cell carcinoma, Kaposi sarcoma, Rhabdomyosarcoma, melanoma etc. (Ref. G) They are characterized by uncontrolled growth and spread of abnormal cells.

Symptoms of skin tumours may include obvious growth, change in skin color or texture, bleeding, itching or pain. Tumours, especially the malignant types, have been on the increase in Nigeria. This has been attributed to the increasing prevalence of HIV in the country (1,2). Albinism is another major risk factor(3,4). Other possible risk factors include UV radiation, weakened immune system from various causes, family history, exposure to chemicals, etc.

This increase has also been attributed to the rise in oil exploration in the nation. **And subsequent oil spillage. Oil spillage is the unintentional release of crude oil or petrochemical products into the environment. Toxic substances emitted include benzene, toluene, naphthalene, heavy metals, etc. (Egbe et al, 2010).** The Niger Delta region faces a lot of oil spillage and is highly vulnerable to the effects of this spillage which includes health disorders like cancers (5,6).

OBJECTIVES

The main objective of this study is to document the incidence and types of skin tumours (including skin cancers) over a 12 year period in the Dermatology Out-patient clinic of The Rivers State University Teaching Hospital, PortHarcourt, Nigeria.

METHODOLOGY

This 12 year retrospective study was performed in the Dermatology out-patient clinic (DOPC) of Rivers State University Teaching Hospital, Port-Harcourt, Nigeria.

Medical records of patients with various types of skin tumours that attended the clinic between January 2012 and December 2023 were retrieved and relevant clinicopathologic data were collated, entered into Microsoft Excel spreadsheet and analysed. These records include biodata, clinical information and relevant histopathological diagnosis.

RESULTS

A total of 1,931 patients attended the clinic during the period studied. There was a steady increase in the number of patients seen over the years: **From just 85 patients in 2012 to 325 patients in 2023.** Out of this number, 102 (5.3%) presented with various skin tumours. There was an equal number of males and females with skin tumours **51 males and 51 females.** (M:F = 1:1).

The age range of the patient with skin tumours is between 11 and 88 years. Majority of affected patients (65.7%) fall into the 20 – 44 years age group followed by those above 60 years of age that constituted 14.7%. The mean age of the patients is 35.2 ± 16.6 years.

Most of the patients with skin tumours were students (34.3%).

Total number of patients with skin tumours were 102, out of which 83 (81.4%) were benign, 1 (0.98%) was premalignant and 18 were malignant (17.64%).

Table1:AgeDistributionofthepatients

Agegroup 0-9 10-19 20-44 45-59 ≥60 Total

Year	0-9	10-19	20-44	45-59	≥60	Total
2012	0	1	2	0	3	6
2013	0	0	1	0	1	2
2014	0	0	0	1	0	1
2015	0	0	4	1	0	5
2016	0	1	2	1	1	5
2017	0	0	2	0	1	3
2018	0	0	1	0	0	1
2019	0	0	0	0	1	1
2020	0	0	7	1	2	10
2021	0	4	9	5	3	21
2022	0	2	10	0	1	13
2023	0	2	29	1	2	34
Total	0	10	67	10	15	102
Frequency(%)	0	9.8	65.7	9.8	14.7	100

Table 2: Frequency Of Benign Tumours (Total N=83)

Tumour	Frequency	Tumour	Frequency
Acne Keloidalis Nuchae	6(7.22)	Lipoma	1(1.20)
Acrochordon (skin tags)	2(2.40)	Mulloscum Contagiosum	2(2.40)
Dermal Naevus	1(1.20)	Prurigo nodularis	1(1.20)
Dermatofibroma	1(1.20)	Pyogenic granuloma	1(1.20)
Dermatosis Papulosa nigra	4(4.81)	Sarcoidosis	1(1.20)
Eccrine Hidrocystoma	1(1.20)	Sebaceous Cyst	2(2.40)
Eruptive Xanthoma	1(1.20)	Seborrhoeic Keratoses	1(1.20)
Haemangioma	1(1.20)	Steatocystoma Multiplex	1(1.20)
Keloids	24(28.91)	Syringoma	4(4.81)

Neurofibromatosis

8(9.64)

Warts

20(24.09)

Total	83(100)		
--------------	----------------	--	--

The most frequent benign tumour is keloid(28.91%)followed by viral warts (24.09%) and neurofibromatosis (9.64%)(see Table 2 above).Among the cases of warts, genital warts consisted of 60% (12) of the total warts, plantar warts were 15% (3) andthe rest 25% were common warts (verruca vulgaris). One of the verruca warts were disseminating occurring in multiple sites.

Out of the Neurofibromatosis cases,2 of them (25%) wereplexiformNeurofibromatosis.

The most frequentmalignant tumouris Kaposi sarcoma followed by metastatic skin cancer as shown in table 3 below.RVD and Post-transplant immunosuppression was associated with Kaposisarcoma

SKIN TUMOURS IN RSUTH PORTHARCOURT

Table3:FrequencyofMalignantTumours(N=18)

Malignant Tumour	FrequencyN(%)
Basal cell Carcinoma	1(5.6)
Kaposi Sarcoma	10(55.5)
Melanoma	1(5.6)
Metastatic skin cancer	3(16.7)
Rhabdomyosarcoma	1(5.6)
Squamous cell carcinoma	2(11.0)
Total	100

DISCUSSION

Skin tumours constituted 5.3% of cases seen in the DOPC of RSUTH. This is in contrast to a study done in Lagos where skin tumours constituted 19.8% of all dermatological consultations and in South Africa 23.73% of all dermatological consultations (7,8).

The reason for this difference may be attributed to the higher number of dermatologists present in these centres compared to our study centre and hence a higher number of patients seeking specialist care. Cancer awareness and fear of cancer is another possible reason.

Male to female ratio of those with skin tumors is 1:1, this is comparable to a studies done in the South Western as well as the South Eastern part of the country (1:1.06) and (1:1.14) respectively (9,10).

The equal affectation of both sexes is not surprising because both sexes are equally exposed to UV radiation and other environmental factors. Both sexes have similar immune systems and also, genetic mutations that increase cancer risk affect both sexes.

Benign skin tumours accounted for 82.3 % of the skin tumor cases seen in our study. This high figure is comparable to other studies done elsewhere in Burkina Fasso, 96.5%, and Lagos, Nigeria, 68.7 %. (11,7). This high value is expected because benign skin tumours are generally more common than the malignant types due to genetic predisposition, environmental factors like viruses, cellular differentiation, growth regulation, immune surveillance and early clinical detection.

The commonest benign tumours in this study was keloids(28.91%), followed by viral warts(24.09%) and then neurofibromatosis(9.64%) . A similar study in South Eastern Nigeria also listed soft tissue fibromas like keloids and dermatofibromas as the commonest benign skin tumours(10). This is in contrast to another study in the subregion where the commonest benign lesions were found to be cutaneous papilloma(7). The reason for the high prevalence of keloids may be due to the relatively young age group of patients studied(mean age of patients with tumours was 35.2 years) since keloids are known to occur in younger persons. Other possible associated factors are race and genetic predisposition.

The only premalignant lesion in this study was actinic keratosis. Actinic keratosis may progress to squamous cell carcinoma if left untreated. This was found in a patient with albinism. This is not surprising because it is rare in blacks due to the protective effect of melanin. It is commoner in Caucasians and albinos. Squamous cell carcinoma was also found in another albino patient in this study which may have arisen from a previous Actinic keratosis.

17.6% of all skin tumours in this study was found to be malignant. This is comparable to a study done in Benin where 26.9% of skin biopsy samples were found to be malignant (12). Another study in Lagos Nigeria revealed a prevalence of 31.2% in (7). These low figures can be explained by the fact that blacks have a lower prevalence of skin cancers due to the protection by melanin (13,14). **This is not the case in Western , overseas countries where skin cancers are more prevalent and for this reason, one out of every 3 diagnosed cancer is a skin cancer. Skin cancers are the commonest diagnosed cancers in Caucasians with a rising prevalence. This may partly be attributed to the continuous depletion of the ozone layer (L) Australia and New Zealand are said to have the highest prevalence of skin cancers.**

The commonest malignant skin tumor in this study is Kaposi sarcoma followed by metastatic skin cancer. This is comparable to another study done in the South –South region of Nigeria where Kaposi Sarcoma was also found to be the commonest skin malignancy (38.8%)(15). This is in contrast to other studies in the Southwestern part of the country where the commonest skin malignancy was Squamous cell carcinoma followed by Melanoma (7,9). It is also in contrast to studies done in the US and other parts of the western world that revealed Basal cell carcinoma as the commonest skin malignancy. The high incidence of Kaposi Sarcoma in this study may be attributable to the increasing prevalence of HIV infection in the country and the fact that the south- south zone of the country has the highest prevalence of HIV nationwide-3.1 % (16).

Another risk factor for Kaposi Sarcoma identified in the study is post renal transplant immunosuppression. This may be attributed to the increasing number of transplant patients in the country due to the establishment of renal transplant centres and increasing number of renal transplant recipients returning home after transplant carried out overseas.

CONCLUSION

There is a relative low prevalence of skin tumours among patients attending the skin clinic in RSUTH(5.3%).The Commonest benign tumor is Keloids while the commonest malignant tumour is Kaposi sarcoma.

Consent

As per international standards, parental written consent has been collected and preserved by the author(s).

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.

3.

REFERENCES

1. Mandong B M, Chirdan L B, Anyebe A O, Mannaseh A N. -Histopathological study of KS in Jos: A 16 year review. *Annals of Afr Med* 2004; 3(4): 174-176.
- 2 . Onunu A N, Okoduwa C, Eze E U, Adeyekan A A, Kubeyinje E PA .
3. Yakubu A, Mabogunje O A. - Skin cancer in African albinos. *Acta Oncol* 1993; 2: 621-622.
R. A. Kaposi sarcoma in Nigeria. *Int J Dermatology* 2007; 46: 246-267.
4. Kromberg J G, Castle D, Zwane E M, Jenkins T. - Albinism and skin cancer In South Africa. *Clin. Genet.* 1989; 43-52.
.Egbe RE, Thompson D Environmental challenges of oil spillage for families in oil producing communities of the Niger Delta region. *JHER* [Internet]. 2010 Dec [cited 2018 Jun 19]; 13:24–34. Available from: <http://www.heran.org/html/jher13/2egbe.pdf>
E. Overton, T. Wade, J. Radovic, B. Meyer, M.S. Miles, S. Larter
Chemical composition of Macondo and other crude oils and compositional alterations during oil spills
Oceanography, 29 (2016), pp. 50-63

[5. International Agency For Research In Cancer/World Healthorganisation Population FactSheet](https://gco.iarc.fr/today/fact-sheets-) : Nigeria.<https://gco.iarc.fr/today/fact-sheets->
Accessed 09/06/2024

6. Hurtig AK, San Sebastián M. Geographical differences of cancer incidence in the Amazon basin of Ecuador in relation to residency near oil fields. *Int J Epidemiol* 2002; 31: 1021– 1027.

7. Ayanlowo, O.; Daramola, A.O.; Akinkugbe, A.; Olumide, Y.M.; Banjo, A.A; Abdulkareem F. Skin tumors at the Lagos University Teaching Hospital, Nigeria *West African Journal of Medicine* 2013; 32(4), : 286-290.

8. Whiting D,A. Skin Tumours in White South Africans Part I. Patients, Methods and Incidence. *SA Mediese Tydskrif*. 1978. pp98-10

9. Adedayo I. S., Olakunle F. B., Oladipo Omoseebi, Olagoke Erinomo, Tope M. I., Shiyanbola A,C.et al Patternof Skin Cancers in a Tertiary Medical Center in Southwest NigeriaAnnals ofAfrican Surgery 2022;Vol. 19 No. 2. doi.org/10.4314/aas.v19i2.3
- 10.NnadozieU.U., OkekeU.V., Maduba C.C., Ugbala A,Ogbuanya U.A., Oguonu C.A.,et al. Morphological Pattern of Benign Skin Tumors in a Teaching Hospital in SoutheastNigeria. Ann Afr Surg. 2024; 21(3).DOI: <http://dx.doi.org/10.4314/aas.v21i3.2>
- 11.Xavier-Junior JC, Ocanha □Xavier JP. WHO (2018) classification of skin tumors.
- 12.LeBoit P.E, Burg G, Weedon D, Sarasin A, editors. World Health Organization Classification of Tumors. WHO Pathology and Genetics of Skin Tumors. Lyon: IARC Press, 2006: 9-11.

11. Barro-Traoré F, Traoré A, Konaté I, Traoré SS, Sawadogo NO, Sanou I, et al. Epidemiological features of tumors of the skin and mucosal membranes in the department of dermatology at the Yalgado Ouedraogo National Hospital, Ouagadougou, Burkina Faso. *Sante*. 2003;13(2):101-4. French. PMID: 14530122.11

12. Forae GD, Olu-Eddo AN. Malignant Skin Tumors in Benin City, South-South, Nigeria. *Oman Med J* 2013;28(5):311-315.

13. Bradford PT. Skincancer in skin of color. *Dermatol Nurs*. 2010;21(4):170-8.

14. Chalya PL, Gilyoma JM, Kanumba ES, et al. Dermatological malignancies at a University Teaching Hospital in north-western Tanzania: a retrospective review of 154 cases. 2012;14(1):1-9.

C. Urban K, Mehrmal S, Uppal P, Giese RL, Delost GR. The global burden of skin cancer: a longitudinal analysis from the Global Burden of Disease Study, 1990–2017. *JAAD Int*. 2021;2:98–108. <https://doi.org/10.1016/j.jdin.2020.10.013>.

D. Ali L, Roky AH, Azad AK, et al. Autophagy as a targeted therapeutic approach for skin cancer: evaluating natural and synthetic molecular interventions. *Cancer Pathog Ther*. 1 February 2024, <https://doi.org/10.1016/j.cpt.2024.01.002>.

E. Leiter, U., Eigentler, T. & Garbe, C. Epidemiology of skin cancer. *Adv. Exp. Med. Biol.* 810, 120–140 (2014).

F.) Amdad Hossain Roky, Mohammed Murshedul Islam, Abu Mohammed Fuad Ahasan, Md Saqline Mostaq, Md Zihad Mahmud, Mohammad Nurul Amin, Md Ashiq Mahmud,

Overview of skin cancer types and prevalence rates across continents,

Cancer Pathogenesis and Therapy,2024,ISSN 2949-7132, <https://doi.org/10.1016/j.cpt.2024.08.002>.

(<https://www.sciencedirect.com/science/article/pii/S2949713224000582>)

H.)Marks R. An overview of skin cancer: incidence and causation. *Cancer*. 1995;75(suppl S2):607. doi: 10.1002/1097-0142(19950115)75:2+<607::aid-cnrcr2820751402>3.0.co;2-8. [DOI] [PubMed] [Google Scholar]

I)Sober AJ, Koh HK, Tran NT, Washington CV. New York: McGraw Hill; 1998. Melanomas and other skin cancers, in Harrison's principles of internal medicine; pp. 543–549. [Google Scholar]

J)Asuquo ME, Ebughe G. Cutaneous Cancers in Calabar Southern Nigeria. *Dermatology Online Journal*. 2009;15(4):11. [PubMed] [Google Scholar]

K) Rogers HW, Weinstock MA, Harris AR, Hinckley MR, Feldman SR, Fleischer AB, et al.

L.) Kricker A, Amstrong BK, English DR. Sun exposure and non- melanocytic skin cancer. *Cancer Causes Control*. 1994; 5(4):367-392.

15. Asuquo, M. E, Ebughe, G. Cutaneous cancers in Calabar, Southern Nigeria. *Dermatology Online Journal*,2009 15(4).

<http://dx.doi.org/10.5070/D36jz5v3fg> Retrieved from <https://escholarship.org/uc/item/6jz5v3fg>

M.)Ochicha O, Edino ST, Mohammed AZ and Umar AB. Dermatological malignancies in Kano, Northern Nigeria: A histopathological review. *Annals of African Medicine*. 2004; 3(4):188-191.

N.)Asuquo M, Ugare G, Odio B, Ebughe G. Squamous cell carcinoma of the skin in Calabar. *Nigerian Journal of Surgical Sciences*. 2006; 16(1):35-38.

O.)Rafindadi A H. A study of 1959 solid cancers seen in ABUTH, Zaria. *Nigerian Journal of Surgery*. 1998; 5:45-48.

P.)Yakubu A, Mabogunje A. Skin cancer in Zaria, Nigeria. *Trop Doct*. 1995; 25(suppl 1):63-67.

Q.)Mandong BM, Orkar KS, Sule AZ, Dakum NL. Malignant skin tumours in Jos University Teaching Hospital, Jos, Nigeria (hospital-based study). *Nigerian journal of Surgical Research*. 2000; 3:29-33.

16.Nigeria Hiv/AidsIndicator And Impact Survey March 2019 .

