

Effect of goat breeds on the milk composition under climatic conditions of BhandarejTahsil of Dausa district, Rajasthan, India

Abstract: The aim of the study is to assess milk composition at specific location. The study was designed as per the resources available at local situation. The research has conducted at the Bhandarejtahsil of Rukmani Devi College of Agriculture, Dausa district of Rajasthan during 2021-23. The specific gravity of Jamunapari goat breed's milk was greater than that of Sirohi animals. 05 milk samples of Jamunapari Goat and 05 milk samples were collected from Sirohi goat breed during the lactation at different villages of dausatahsil of dausa district throughout two years. The overall fat per cent in the both breeds during 2021- 23 of all the 1200 samples was found to be 4.68 ± 0.021 , Solids Not Fat (SNF) 8.45 ± 0.030 per cent and total solids percentage was found to be 13.32 ± 0.035 . The protein per cent during 2021 - 22 conditions was significantly ($p < 0.01$) greater than that of farm rearing goat milk in both the breeds. The statistical analysis also revealed that the lactose content in 2021 – 22 has significantly higher than that of 2022 -23 in Sirohi as well as Jamunapari goat breeds at $p < 0.05$. The overall average ash content in both breed's milk during 2021 - 23 of all samples was found to be 0.815 ± 0.007 per cent. Breed had conspicuous effects on milk quality of goats under study.

Keywords: SirohiGoat ,Jamunapari goat,Milk Composition, Dausa, Rajasthan

INTRODUCTION

The productive improvements among dairying animals can be made through proper management, feeding and handling, etc. which may influence expression of productive characters as per its heritability nature. Before identifying the animals for breeding and production purpose screening of animals shall be performed on the basis of physical traits (Singh et al., 2013). The global goat population currently stands at 921 million, of which over 90% are found in developing countries. Asia is home to about 60% of the total world goat population and has the largest goat breed share of 26%. Goats play a vital socioeconomic role in Asian agriculture, particularly for resource-poor people living in harsh environments. Non-cattle milk accounts for approximately 15% of the total milk consumption by humans worldwide (Singh et al 2014). India today, stands first in the area of milk production at the world level, with an annual growth rate of about 4%. The country's milk production in 2010 was estimated to be 110 million tones. A large quantity of milk produced in the country amounting to over 46% is being consumed as liquid milk. The production and use of animal products in the use of human diet is receiving tremendous attention. With this object in view the need for developing Animal Husbandry is recognized very well. The other objects are to provide animal power for farmings and adoption of better land use pattern (Singh et al 2012). Goat population of our country increased from 47.14 million in the year 1951 to 124.5 million during 2005 (Singh and Sharma, 2013a).

Goats are integer part of livestock production and play a vital role in the socio-economic structure of rural poor. The aim of this study was to project the importance and

significance of goat milk with special reference to Indian field and farm rearing conditions. There are adverse ecological and physiological constraints in the Indian system of goat farming. Goats are more often poorly managed and this is attributed to their ability to survive under harsh conditions and also because most people in rural areas rear goats for their subsistence purposes to support their families. This benefit is often not shown in national statistics because of informal trading and slaughtering (Singh et al 2014a). Goat milk contains less lactose than cow's milk, so is less likely to trigger lactose intolerance (Singh et al 2014b). Goat meat being high quality protein source is the choicest meat in domestic market (Singh et al 2014c). Major population of India is primarily depends on agricultural based system for their daily life including goat keeping that constitute an important rural business of small marginal farmers and landless labours (Singh et al 2014d). Reproductive management of an animal is governed through a number of parameters, viz. age at first conception, age at first calving and first gestation length etc. (Singh et al 2014e), (Singh et al 2014f), (Singh et al 2014g) and (Singh et al 2014h). Goat milk contains less lactose than cow's milk, so is less likely to trigger lactose intolerance (Singh and Sharma, 2015).

It has since played a significant socioeconomic role in the evolvement of human civilization around the world (Singh and Sharma, 2015a). The nutritional value of milk is closely related to its composition, which is affected by factors such as breed, diet, stage of lactation, season etc. Goat milk has more calcium (Ca), phosphorus (P), potassium (K), magnesium (Mg) and chloride (Cl) and less sodium (Na) and sulphur (S) contents than cow milk (Singh and Sharma, 2016). Livestock production is backbone of Indian agriculture contributing 7% to national GDP and source of employment and livelihood for 70% population in rural areas. Animals reared in intensive production systems consume a considerable amount of protein and other nitrogen-containing substances in their diets (Singh et al 2017). This benefit is often not shown in national statistics because of informal trading and slaughtering (Singh and Sharma, 2017a).

Goats play a vital socio-economic role in Asian agriculture, particularly for resource-poor people living in harsh environments (Singh et al 2018). Jamnapari (or Jamunapari) is a breed of goat originating from Indian subcontinent. Since 1953 they have been imported to Indonesia (popular as Etawa goat, and its mixture with a local goat called "PE", *peranakan Etawa* or Etawa mix) where they have been a great success (Singh et al 2017b). These breeds or types were distributed across the world as a result of the migration and translocation of humans, usually due to changing climatic conditions and natural resources (Singh and Sharma, 2017c). The Sirohi is an Indian breed of domestic goat. It is named for its area of origin, the Sirohi District of Rajasthan, in north-western India. It may be described as a dual-purpose breed, reared for both meat and milk production, or as meat breed. It is well adapted to the dry tropical climate of Rajasthan.

Milk-secreting tissues and various ducts throughout the udder can be damaged by bacterial toxins, and sometimes permanent damage to the udder occurs (Singh and Singh, 2020). Livestock has become an integral part of all interventions aimed at reducing rural poverty and enhancing food and nutrition security (Singh and Somvanshi, 2020). India is endowed with a significant share of the world's livestock population growing steadily and continuously (Singh, G. 2019). The goat is thought to have been the earliest domesticated ruminant and of all the species of domesticated animals

except dog, has the widest ecological range (Singh, 2024). Man, Animal and Nature are in symbiotic relationship for their survival and sustenance (Singh et al., 2024a).

MATERIALS AND METHODS

The research has conducted at the Bhandarejtahsil of Rukmani Devi College of agriculture, Dausa district of Rajasthan with broad objective, as 'Effect of goat breeds on the milk composition under climatic conditions of Bhandarejtahsil of Dausa district Rajasthan during 2021-23. 05 milk samples of Jamunapari Goat and 05 milk samples were collected from Sirohi goat breed during the lactation at different villages of Dausatahsil of Dausa district throughout two years. A total of 1200 milk samples were collected from both goat breeds in Dausa Tahsil, with 600 milk samples from Jamunapari goats and 600 milk samples from Sirohi goats. All samples were analyzed by electronic milk analyzer. Data were analyzed using the ANOVA procedure of RBD (at 5% and 1% significance levels) for milk samples.

RESULTS AND DISCUSSION

1. Effect of Goat breeds on Specific gravity of milk:-

The data obtained on specific gravity in the present study of Sirohi and Jamunapari goat breed milk are presented in Table 1.

Table 1 :Effect of Goat breeds on Specific gravity of milk

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t)	
						5%	1%
1.	Sirohi	1.0220±0.0003	1.0232±0.0003	1.022±0.0003	3.51 ⁺	1.540	2.145
2.	Jamunapari	1.0286±0.0002	1.0263±0.0003	1.0279±0.0002	3.20 ⁺		
	Overall mean	1.0249±0.0002	1.0247±0.0002	1.0340±0.0002			

Note: +=Significant at 5% level of significance

It is observed from Table 1. that the specific gravity of Sirohi and Jamunapari goat breeds has found to be 1.0220±0.0003 and 1.0286±0.0002, respectively with an average of 1.0249±0.0002. It is observed from above table that specific gravity of Jamunapari goat breed's milk has significant than that of Sirohi animals.

2. Effect of Goat breeds on Fat of milk %:-

The result obtained for the fat percentage of milk of Sirohi and Jamunapari goat breed are presented in Table 2.

Table 2: Effect of Goat breeds on Fat % of milk.

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t)	
						5%	1%
1.	Sirohi	4.33±0.034	4.71±0.033	4.57±0.032	2.987 ⁺	1.547	2.321
2.	Jamunapari	4.71±0.087	4.80±0.023	4.805±0.011	2.354 ⁺⁺		

	Overall mean	4.52±0.060	4.75±0.046	4.68±0.021		
--	---------------------	------------	------------	------------	--	--

Note: +=Significant at p <0.05

++=Significant at p <0.01

The average percentage of fat in the milk of Sirohi and Jamunapari goat breeds has found to be 4.33±0.034 and 4.71±0.087 respectively with an average value of 4.52±0.060 per cent during 2021 - 22. Similarly, fat content in 2022 - 23 of Sirohi and Jamunapari goat breeds has found to be 4.71±0.033 and 4.80±0.023, respectively with an average value of 4.75±0.046 per cent. The overall fat per cent in the both breeds during 2021- 23 of all the 1200 samples has found to be 4.68±0.021. The results of the present investigation on fat content of Sirohi and Jamunapari goat breeds are in consonance with the observation of Singh and Sharma (2015b), However, Prasad et al. (2005) reported a higher fat percentage in goat milk. Lower values have been reported by Agnihotri et al. (2002).

3. Effect of Goat breeds on Protein of milk %:-

The data on Protein percentage in the present investigation of Sirohi and Jamunapari goat breed's milk during 2021 - 23 are recorded in Table 3.

Table 3 : Effect of Goat breeds on Protein % of milk

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t) 1% 5%
1.	Sirohi	3.28±0.028	3.21±0.034	3.24±0.032	5.875 ⁺⁺	1.960 2.576
2.	Jamunapari	3.19±0.027	3.23±0.036	3.21±0.032	4.51 ⁺⁺	
	Overall mean	3.23±0.028	3.22±0.035	3.225±0.32		

Note: ++=Significant at p <0.01

It is observed from the Table 3 that the protein percentage in the Sirohi and Jamunapari goat breeds during 20 – 21 has found to 3.28±0.028 and 3.21±0.034, respectively with an average of 3.23±0.028 per cent. The protein content of aforesaid breeds during 2022 - 23 found to be 3.21±0.034 and 3.23±0.036 per cent respectively with an average value of 3.22±0.035 per cent. It is clear from the above table that protein per cent during 2021 - 22 has significantly (p <0.01) greater than that of 2022 -23 goat milk in both the breeds. The statistical analysis revealed that protein content of Sirohi goat breed has significantly higher than that of Jamunapari goat breed during 2021 - 22. The analysis of variance of these data (Table 3) revealed that protein content variation either Sirohi or Jamunapari goat breeds had significant (p < 0.05) effect during 2021 - 23.

4. Effect of Goat breeds on Lactose of the milk%:-

The lactose per cent in the milk of Sirohi and Jamunapari goat breeds during 2021 - 23 in the present investigation are presented in Table 4.

Table 4 : Effect of Goat breeds on Lactose % of milk

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t) 1% 5%

1.	Sirohi	4.39 ±0.021	4.40±0.022	4.45±0.021	1.859 ⁺	1.960 2.576
2.	Jamunapari	4.47 ±0.020	4.43±0.025	4.41±0.022	1.243 ⁺	
	Overall mean	4.43±0.0205	4.415±0.0235	4.43±0.014		

Note: +=Significant at p <0.05

It is clear from Table 4 that the lactose content in Sirohi and Jamunapari goat breed's milk during 2021 – 22 has found 4.39 ±0.021 and 4.47 ±0.020 per cent, respectively. Similarly the lactose content in above both breed milk during 2022 – 23 has found 4.40±0.022 and 4.43±0.025 per cent, respectively. The overall average lactose per cent in Sirohi and Jamunapari goat breed's milk during 2021 - 23 also calculated and found to be 4.43±0.014 for all 1200 samples. The statistical analysis also revealed that the lactose content in 2021 – 22 has significantly higher than that of 2022 -23 in Sirohi as well as Jamunapari goat breeds at p <0.05. Our results on lactose percentage in above goat breed's milk are in fair agreement with those reported by Prasad et al. (2002) and Agnihotri (2002) have reported higher values on it.

5. Effect of Goat breeds on Ash of milk%:-

The results obtained for ash content of Sirohi as well as Jamunapari goat milk during 2021 - 23 are presented in Table 5.

Table 5: Effect of Goat breeds on Ash % of milk

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t) 5% 1%
1.	Sirohi	0.80±0.008	0.82±0.007	0.81±0.008	3.879 ⁺	1.960 2.576
2.	Jamunapari	0.82±0.008	0.84±0.005	0.82±0.006	3.654 ⁺⁺	
	Overall mean	0.81±0.008	0.82±0.006	0.815±0.007		

Note: +=Significant at p <0.05

++=Significant at p <0.01

The average percentage of ash during 2021 – 22 Sirohi and Jamunapari goat breed's milk has found to be 0.80±0.008 and 0.82±0.008, respectively with an average value of 0.81±0.008 per cent. Similarly ash content in 2022 -23 of aforesaid breed's milk has found to be 0.82±0.007 and 0.84±0.005, per cent with an average of 0.82±0.006 per cent. The overall average ash content in both breed's milk during 2021 - 23 of all samples has found to be 0.815±0.007 per cent. It is also clear from the above table that ash content in the milk of Sirohi or Jamunapari goat breed 2022-23 has significantly greater than that of 2021-22. The analysis of variance table showed that ash content of Sirohi goat breed's milk during 2022- 23 has significantly more than that of Jamunapari goat breed milk. The results of present investigation on the level of ash in Sirohi and Jamunapari goats' milk during 2021 - 23 are slightly higher by the findings (0.78 ± 0.01) of Agnihotri et al. (2002) and Singh and Sharma (2015).

6. Effect of Goat breeds on Total Solids of milk%:-

The results obtained for total solids percentage in milk of Sirohi and Jamunapari goat breed during 2021 - 23 are presented in Table 6.

Table 6 : Effect of Goat breeds on Total solids % of milk

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t) 5% 1%
1.	Sirohi	13.03±0.030	13.16±0.035	13.09±0.032	1.21 ⁺	1.960 2.576
2.	Jamunapari	13.23±0.036	13.87±0.039	13.55±0.037	1.985 ^{NS}	
	Overall mean	13.13±0.033	13.51±0.037	13.32 ± 0.035		

Note: NS=Non Significant

+ =Significant at $p < 0.05$

According to above Table 6, the average total solids percentage in milk of Sirohi and Jamunapari goat breed during 2021 – 22 has found to be 13.03±0.030 and 13.23±0.036, respectively with an average value of 13.13±0.033. Similarly the total solids percentage in the milk of above goat breeds during 2022 – 23 has found to be 13.16±0.035 and 13.87±0.039, respectively with an average value of 13.13±0.037 per cent. The overall average total solids percentage of all 1200 samples has found to be 13.32 ± 0.035. It has observed from above table that Sirohi goat breed's milk total solids were significantly ($p < 0.050$) higher in 2021-22 than 2022-23 whereas in case of Jamunapari goat breed an insignificant difference has observed in total solids content. The analysis of variance in the table revealed that the breed variation during 2021 -22 has non-significant but it has significant during 2022 -23 at 5% level of significance.

The level of total solids percentage in the milk of above goat's breeds as obtained in the present study, Compared favorably with the results of Pal et al. (2011) and Agnihotri (2002). Prasad et al. (2005) have, however, reported higher values of Singh and Sharma (2015).

7. Effect of Goat breeds on Solids-Not-Fat %:-

The data on solids-not-fat percentage in Sirohi and Jamunapari goat breed's milk during 2021 - 23 are recorded in Table 7.

Table 7: Effect of Goat breeds on Solids-Not-Fat % of milk.

Sl. No.	Breeds	2021 - 22	2022 -23	Overall average	Test of significance	Table value (t) 5% 1%
1.	Sirohi	8.42±0.036	8.44±0.024	8.43±0.030	3.659 ⁺⁺	1.960 2.576
2.	Jamunapari	8.46±0.034	8.48±0.026	8.47±0.030	3.854 ⁺⁺	

	Overall mean	8.44±0.035	8.46±0.025	8.45±0.030		
--	---------------------	------------	------------	------------	--	--

Note: ++=Significant at $p < 0.01$

It is observed from the Table 7 that the solids-not-fat percentage in the milk of Sirohi and Jamunapari goat breeds during 2020 – 21 has found to be 8.42 ± 0.036 and 8.46 ± 0.034 , respectively with an average value of 8.44 ± 0.035 per cent. The solids-not-fat content of above breeds during 2022 - 23 samples has also calculated and found to be 8.44 ± 0.024 and 8.48 ± 0.026 per cent, respectively with an average of 8.46 ± 0.025 per cent. The overall average solids-not-fat percentage of above samples in 2021-22 and 2022 - 23 of all 1200 samples has 8.45 ± 0.030 per cent. The statistical analysis showed that the solids-not-fat percentage has significantly ($p < 0.01$) higher in 2022-23 either Sirohi or Jamunapari goat breed than that of 2021-22. It is due to higher fat percentage and lower percentage of protein, lactose and ash in 2022-23 of Sirohi as well as Jamunapari goat breeds. The analysis of variance table on these data also revealed that significantly breed variation on solids-not-fat content has observed either 2021-22 or 2022-23. The results of the present investigation on solids-not-fat content of Sirohi and Jamunapari goat breed's milk 2021-23 are in consonance with slightly higher values of Singh and Sharma, (2015).

CONCLUSION

The specific gravity and fat percentage in the milk samples of Sirohi has significant effects than milk of Jamunapari goat breeds during 2021 - 23. Breed had conspicuous effects on milk quality of goats under study.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.

REFERENCES

- Agnihotri, M.K., Singh, Nawab and Babji, Y. 2002. Milk composition of goat reared under field condition. *Indian J. of Anim. Sci.* **72**(11): 1019-1921.
- Pal, U.K., Mandal, P.K.; Rao, V.K. and Das, C.D. 2011. Quality and Utility of Goat Milk with Special Reference to India: An Overview. *Asian Journal of Animal Sciences*, **5**: 56-63.
- Prasad, H.; Tiwari, H.A. and Sengar, O.P.S. 2005. Milk yield and their crosses with Jamunapari, Barbari and Black Bengal breeds of goat. *Small Rumin. Res.* **58**(2):195-199.
- Singh, G., Dutt, G., Sharma, R.B., Fatima, A., and Singh, R.P. (2012). Study of first gestation length in Gir cows, *The Journal of Rural and Agricultural Research*, **12**(1): 64-65
- Singh, G., Dutt, G., Sharma, R.B., Singh, S.K., Fatima, A. and Chauhan, S.V.S. (2012a)., An analytical study of reproductive performance in Gir cows, *Indian Research Journal of Extension Education, Special Issue*, (2): 203 - 206
- Singh, G. and Sharma, R.B. (2013). Influence of breeds on goat milk composition under field and farm rearing conditions, *Indian Research. Journal of Genetics & Biotechnology*, **5**(4): 258-261
- Singh, G. and Sharma, R.B. (2014). Effect of season on the milk quality of Jamunapari goats under field and farm rearing condition, *Indian Research. Journal of Genetics & Biotechnology*, **6**(1): 335-339
- Singh, G., Thorat, G.N., Trivedi, M.S., Mishra, R. and Sharma, S.K. (2014a). A test to measure knowledge about poultry management practices, *The Journal of Rural and Agricultural Research*, **14**(2): 44-47
- Singh, G., Sharma, R.B., and Mishra, R. (2014b). Seasonal variations in the milk minerals of Jakhrana goats under field and farm rearing conditions, *Journals of community mobilizations and sustainable development*, **9**(2): 120 - 123
- Singh, G., Sharma, R.B., Mishra, R. and Rajput, S. (2014c). Effect of multiple births on Jakhrana goat milk quality under field and farm rearing conditions, *Indian Research. Journal of Genetics & Biotechnology*, **6**(4): 629-635
- Singh, G. Sharma, R.B., and Mishra, R. (2014d). Effect of season on the milk quality of Jakhrana goats under field and farm rearing condition, *Indian Research. Journal of Genetics & Biotechnology*, **6**(3): 571-577
- Singh, G. Sharma, R.B., Mishra, R. and Rjput, S. (2014e). Effect of season on goat meat composition under field and farm rearing conditions, *Indian Research. Journal of Genetics & Biotechnology*, **6**(3): 511-517
- Singh, G. Sharma, R.B., and Mishra, R. (2014f). Effect of multiple births on Jamunapari goat milk quality under field and farm rearing conditions, *Indian Research. Journal of Genetics & Biotechnology*, **6**(2): 453-458
- Singh, G. Sharma, R.B., Kumar, A. and Chauhan, A. 2014g. Effect of Stages of Lactation on Goat Milk Composition under Field and Farm Rearing Condition, *Advances in Animal and Veterinary Sciences*, **2**(5): 287 – 291

Singh, G. Dutt, G. and Rajput, S. (2014h). Study of age at first calving in Gir cows, *Indian Research. Journal of Genetics & Biotechnology*, **6**(1): 362-365

Singh, G. and Sharma, R.B. (2015). Effect of multiple births on Jakhrana goat milk minerals under field and farm rearing Conditions, *Indian Research. Journal of Genetics & Biotechnology*, **7**(2): 227 – 234.

Singh, G. and Sharma, R.B. (2015a). Influence of breed on goat meat composition under field and farm rearing Conditions, International conference on Emerging Trends in Biotechnology and Science with Especial Reference to Climatic Change, 15 – 17 Feb., 2015 held at KVK TonkBanasthaliVidyapith organized by Indian Society of Genetics and biotechnology Research

Singh, G. and Sharma, S.K. (2015b). On Farm Trial (OFT) of pearl millet green fodder at Tonk district, *The Journal of Rural and Agricultural Research*, **15**(2): 28-29

Singh, G. and Sharma, R.B. (2016). Impact of stages of lactation on the minerals of Jakhrana goat milk under field and farm rearing condition, *Research Journal of Animal Husbandry and Dairy Science*, **7**(1): 28-34

Singh, G. and Sharma, R.B. (2016a). Effect of goat breeds on the milk mineral composition under field and farm rearing conditions, *The Bioscan*, **11**(2): 691 – 694

Singh, G. and Sharma, R.B. (2016b). Effect of rearing systems on mineral contents of milk during lactation in Jamunapari goats, *Indian Journal of Small Ruminants*, **22**(2): 270-271

Singh, G., Sharma, R.B. and Singh, M. (2017). Green fodder production potential of Oat cv. Kent under semi-arid climatic conditions of Tonk-Rajasthan in Frontline Demonstration, *International Journal of Current Microbiology and Applied Science*, **6**(3): 2228-2232

Singh, G., Sharma, R.B. Singh, M. and Sharma, S.K. (2017a). Utilisation of agricultural wastes in participatory poultry farming with women under climatic conditions of Tonk district of Rajasthan, *Agriculture Science Digest*, **37**(1): 60-63

Singh, G. and Sharma, R.B. (2017b). Effect of field and farm rearing conditions on the sensory quality of goat meat, *Journal of Community Mobilization and Sustainable Development*, **11**(2): 188-192

Singh, G., Sharma, R.B., Chahal, B.P., Singh, M. and Sharma, S.K. (2018). Effect of multiple births on Jamunapari goat milk minerals under field and farm rearing conditions, *Indian Journal of Animal Research*, **52**(4): 628-631

Singh, G., Sharma, R.B., Singh, M. and Choudhary, R. (2017c). Effect of Season on Jamunapari Goat Meat Composition under Field and Farm Rearing Condition, *International Journal Pure and Applied Bioscience* **5**(2): 563-568 doi: <http://dx.doi.org/10.18782/2320-7051.2714>

Singh, G. and Sharma, R.B. (2017d). Seasonal impact on the minerals of jamunapari goat milk minerals under field and farm rearing condition, *International Journal of Current Microbiology and Applied Science*, **6**(9): 1298-1303.

Singh, G. (2019). Analytical study of Front Line Demonstration (FLD) of Kadaknath poultry farming under climatic conditions of Tonk district of Rajasthan, *The Journal of Rural and Agricultural Research*, **19**(2): 49-52

Singh, G. and Singh, R.P. (2020). An analytical study on mastitis in cows under climatic conditions of Tonk district, *The Journal of Rural and Agricultural Research*, **20**(1): 18-21

Singh, G. and Somvanshi, S.P.S. (2020a). Study on animal rearing practices by dairy owners of district Jaipur rajasthan, *The Journal of Rural and Agricultural Research*, **20**(1): 61-64

Singh, G. (2019a). Effect of area specific mineral mixture on productive performance of murrah buffaloes under climatic conditions of tonk district, *Indian Research. Journal of Genetics & Biotechnology*, **11**(4): 277-281

Singh, G. (2024). Nutrition and feeding management of goats for chevonproduction, *International Journal of Science, Environment and Technology*, **13**(5): 334 – 349

Singh, G., Singh, S., Sharma, K., Sharma L.K. and Kumar, A. (2024a). Effect of goat rearing on environment and rural prosperity in India, *International Journal of Science, Environment and Technology*, **13**(6): 421 – 433

UNDER PEER REVIEW