

***Original Research Article***

**EFFECT OF DIFFERET ORGANIC NUTRIENT SOURCES ON THE GROWTH AND YIELD OF RADISH**

**ABSTRACT**

Indiscriminate use of chemical fertilizers is a common practice which is hazardous both for soil health and environment. It has been more than three decades that global agriculture has been depicted towards organic agriculture owing to sustainability and reduced environmental effects. A field experiment was conducted at IAAS, Lamjung Campus from Poush 2079 to Magh 2079, with the objective to study the effect of various sources of organic manure on the growth and yield of radish. The experiment was conducted in Randomized Complete Design using 7 treatments (Control, FYM, Poultry Manure, Goat Manure, FYM + Poultry Manure, FYM + Goat Manure, Poultry Manure + Goat Manure) with 3 replications of each. Tokinashi variety of radish was used in the experiment. Experiment was done in field with the plot size of (2 x 1)m<sup>2</sup> and plant spacing of 30 cm x 20 cm with the total area of 90m<sup>2</sup>. The effect of different treatments on growth and yield of radish was affected significantly. The highest total root yield (204.33), and biological yield (263.67) was found in Poultry manure. The study suggested that the application of poultry manure was found to be more beneficial and significantly better growth and yield performance of radish.

## **Introduction**

Radish (*Raphanus sativus*L.) is one of the most popular root vegetables of the world which can be grown from tropical to temperate regions. It is grown for the consumption of its young fresh tender tuberous root which can be used to cooked or raw food as a salad(Umar, Ibrahim, Iro, and Obidola 2019). Radish is the dicotyledonous biannual root vegetables of the Brassicaceae family. It was domesticated in the eastern Mediterranean region. (Kitamura 1958). The area, production, and productivity of Radish in Nepal is 16441 hectare, 256,013 metric tonnes and 15.38 tonnes/hectare(Ministry of Finance 2022))

Radish (*Raphanus sativus* L.) is a significant short-duration (70-80 days) root vegetable commonly grown during the winter season(Brintha, and Seran, 2009).The popularity of radish cultivation could be due to its wider adaptability, low cost of production, short crop duration and can be grown in almost all type of soil without much care. Diseases and pest problems are also less in radish as compare to other vegetable crops(Chapagain *et al.* 1970). Radish is a short-duration and quick growing crop, so,the root growth and development should be uninterrupted. For this, optimum nutrition should be provided through organic, inorganic and bio-fertilizer sources.Chemical fertilizers are expensive and result in poor health condition of soil andwater if used repeatedly. So, alternative cheap organic sources of nutrients should be used.(Sanjay Kumar *et al.* 2016)

Organic manures are the types of manures which are derived from animal, human and plant residues which contain plant nutrients in complex organic forms (Kumar *et al.*, 2014). By increasing soil moisture content, pH, bulk density, soil porosity, CEC, and nitrate nitrogen availability while lowering nitrate nitrogen leaching to lower depths, organic manure increases radish root production(Subedi, *et.al*). For improving plant development at 45 and 60 days after sowing (DAS), organic manures, combination applications of vermicompost and chicken manure (50% each), FYM+vermicompost (50%+50%), performed best.(*International journal of plant science*)

In radish plant height, number of leaves, leaf breadth, root length, root diameter, and biomass yield all rose significantly after the application of organic manures (Yadav *et al.*, 2018) The highest yields of biomass and roots were from the application of chicken manure, which was in line with the fertilizer's recommended rate. (Khatri *et al.*, 2019).The application of PM considerably changed the physical characteristics of the soil (p 0.05); it decreased bulk density and increased porosity and moisture content (Kiran *et.al.*, 2016)

## **Materials and Methodology**

The experiment was carried out in the field of the Institute of Agriculture and Animal Science, Lamjung Campus, Sundarbazaar in the western mid hills of the Nepal year 2079/80. This place has a humid tropical climate. The average temperature is 18 °C. The total annual rainfall in this area is reported as 2800mm. The location is at 28.2765 °N, 84.3542 °E with an elevation of 625 meters above sea level. A factorial field experiment at randomized complete block design with three replications was carried out to investigate the effect of organic nutrient sources on the growth and yield of radish. Seven rows with five plants in each row were sown with the spacing of 20 cm\* 30 cm with the total are of 2m<sup>2</sup> plot Tokinashi variety of radish was selected to planted. All the treatment rate were taken according to National Krishi Diary 2021/22.

Table 1: Different treatments and their proportion in plot

S. N	Treatment	Rate (Kg per plot)
T1	Control	0
T2	FYM	8
T3	Poultry Manure	2.67
T4	Goat Manure	4
T5	FYM + Poultry Manure	4 + 1.3
T6	FYM + Goat Manure	4 + 2
T7	Poultry Manure + Goat Manure	1.33+2

Table 2: Experimental design of the plot

	R1		R2		R3	
	T2		T3		T5	
	T1		T2		T2	
	T7		T7		T4	
	T5		T6		T3	
	T4		T1		T7	
	T3		T4		T1	
	T6		T5		T6	

The field was ploughed twice, harrowed and leveled by planking. The weeds and leftover crop residue were removed from the field, and after which it was partitioned into different sections following the experimental layout on Poush 2, 2079 BS. Well decayed organic manure were applied seeding was done days prior to seed sowing, while chemical fertilizers were applied at the time of sowing. Chemical fertilizers were applied in 2 doses, half doses at sowing and half as split dose 30 days after sowing.

There are 21 plots, with 35 plants in each plot. Daily light irrigation was done earlier 10 days. After that, the field was irrigated twice a week, prior to harvest. Manual weeding was done irrespective of the seeding days for each plot to keep them free from the competition with the unwanted plants. Radish was harvested after 60 days of sowing.

### **Data Collection**

Data collection was done four times for the vegetative parameter at 20 days after seeding, 30 days after seeding, 45 days after seeding and 60 days after seeding. And at final harvest data was taken for yield parameters.

### **Vegetative Parameters**

Plant height: Measured up to tallest point (Tip of the longest leaf) using scale.

No. of leaves: Counting of leaves was done from base to top.

Leaf area: Length and breadth of three largest leaves were measured using scale and area was calculated.

Canopy diameter: By averaging the length and breadth of diameter.

### **Yield Parameters**

Root Weight: Weight of the root measured by weighing balance.

Root Length: By using measuring scale.

Root Diameter: Measured using vernier caliper.

Leaf yield: 
$$\frac{\text{Total weight of Leaves(kg)}}{\text{Total area(m}^2\text{)}}$$

Root yield: 
$$\frac{\text{Total weight of roots(kg)}}{\text{Total area(m}^2\text{)}}$$

Biological yield: 
$$\frac{\text{Total weight of roots+Leaves(kg)}}{\text{Total area(m}^2\text{)}}$$

Harvest index(%): 
$$\frac{\text{Root Yield(kg)}}{\text{Biological yield(kg/m}^2\text{)}} * 100$$

### **Data Analysis**

The Data Entry was done using MS- Excel and Data analysis was done using R-Stat(version 4.3.1).

### **Result and Discussion**

#### **Effect of Organic manure on the plant height of Radish**

The highest plant height was recorded from FYM+ Goat at 30 DAS which was similar with all other treatment except control. At 40 DAS higher plant height was obtained from application of poultry manure and its combination with FYM, goat and combination of FYM with goat as compared to control which was similar to individual application of FYM and goat. At 50DAS higher plant height was obtained from poultry and goat alone and its combination and combination of FYM and goat. At 60 DAS higher plant height was obtained from poultry and its combination with FYM as compared to control which was like all the treatments. Overall plant height showed good response to poultry manure. Previous studies also confirm that poultry manure was appropriate to enhance the seedling growth at different stages (Shiyam, Garjila, and Bobboyi 2017) Poultry manure performed best in plant height could be due to the least value of C/N ratio of poultry manure that enhance faster decomposition and quicker release of nutrients than other organic manure sources

Table 3 Effect of organic manure on the plant height

Treatment	Ph1	Ph2	Ph3	Ph4
T1(Control)	3.28 <sup>b</sup>	6.17 <sup>b</sup>	8.83 <sup>b</sup>	19.7 <sup>b</sup>
T2(FYM)	4.93 <sup>ab</sup>	11.17 <sup>ab</sup>	14.04 <sup>ab</sup>	22.04 <sup>ab</sup>
T3(Poultry)	5.37 <sup>ab</sup>	14.77 <sup>a</sup>	20.42 <sup>a</sup>	24.72 <sup>a</sup>
T4(Goat)	4.75 <sup>ab</sup>	10.99 <sup>ab</sup>	18.46 <sup>a</sup>	23.19 <sup>ab</sup>
T5(FYM+POULTRY)	5.19 <sup>ab</sup>	12.66 <sup>a</sup>	16.53 <sup>ab</sup>	25.3 <sup>a</sup>
T6(FYM+ Goat)	6.62 <sup>a</sup>	14.25 <sup>a</sup>	19.53 <sup>a</sup>	23.51 <sup>ab</sup>
T7(Poultry + Goat)	5.49 <sup>ab</sup>	13.69 <sup>a</sup>	20 <sup>a</sup>	22.45 <sup>ab</sup>
LSD	2.31	5.48	8.13	4.37
CV	25.5	25.78	27.12	10.69
SE <sub>m</sub>	1.29	3.08	4.56	2.45
Mean	5.09	11.96	16.84	22.99

Means followed by the common letter within each column are not significantly different at 5% level of significance by Duncan Test.

### Effect of Organic manure on the leaf number of Radish

There was no significant effect of any manure on leaf number at 30 DAS. All manure except FYM were significant in increasing the leaf number at 40 DAS as compared to control. At 50 DAS leaf number was higher in poultry and combination of FYM and goat as compared to control, but its effect was similar to other manures. At 60 DAS, poultry and its combination with goat, FYM and combination of FYM and goat were superior in increasing leaf number as compared to control.

This observation was supported by Kumar,( 2014) who recorded highest number of leaves from poultry manure and minimum number of leaves was recorded in control at 60 DAS. (Sandeep Kumar *et al.*, 2014), The number of leaves and plant height of radish increases with an increase in nitrogen level, which was mostly found in mustard oil cake and poultry manure.(Shahi *et al.*, 2022)

Table 4 Effect of organic manure on the leaf numbers

Treatment	Lno1	Lno2	Lno3	Lno4
T1(Control)	3.57 <sup>a</sup>	5.49 <sup>c</sup>	10.43 <sup>c</sup>	13.39 <sup>c</sup>
T2(FYM)	4 <sup>a</sup>	7.42 <sup>bc</sup>	12.3 <sup>abc</sup>	14.86 <sup>bc</sup>
T3(Poultry)	4 <sup>a</sup>	9.15 <sup>ab</sup>	15.37 <sup>a</sup>	18.98 <sup>a</sup>
T4(Goat)	4 <sup>a</sup>	9.18 <sup>ab</sup>	12.22 <sup>abc</sup>	15.66 <sup>bc</sup>
T5(FYM+ Poultry)	3.94 <sup>a</sup>	8.71 <sup>ab</sup>	11.16 <sup>bc</sup>	17.54 <sup>ab</sup>
T6(FYM+ Goat)	4.12 <sup>a</sup>	9.84 <sup>ab</sup>	14.73 <sup>ab</sup>	17.48 <sup>ab</sup>
T7(Poultry +Goat)	4 <sup>a</sup>	11.14 <sup>a</sup>	13.7 <sup>abc</sup>	16.67 <sup>ab</sup>
LSD	0.62	2.49	3.42	2.82
CV	8.8	16.09	14.99	9.7
SE <sub>m</sub>	0.34	1.4	1.92	1.58
Mean	3.95	8.71	12.84	16.37

Means followed by the common letter within each column are not significantly different at 5% level of significance by Duncan Test.

### Effect of different sources of organic manures on root weight, root length & root diameter of radish

#### Root Weight

The highest fruit weight (204.33 gm) was recorded from Poultry Manure application, surpassing all other treatments in terms of root weight per plant. Control, on the other hand, was found to be on par with FYM and had the lowest weight. However, when compared to FYM coupled with poultry manure and poultry manure combined with goat manure, goat manure displayed statistically equivalent root weight. FYM mixed with goat dung provided a better response than control and FYM alone but was inferior to all other treatments.

This observation was supported by (Ijoyah and Sophie 2009) who reported that the application of poultry manure increased crop yield. Also, Khatri *et al.* (2019) assigned the highest biomass in PM than FYM to the low C/N ratio of PM contributing to faster decomposition and quick release of nutrients.

(Shiyam, Garjila, and Bobboyi 2017) reported that among the different organic manure, PM helps to improve soil condition, increase water holding capacity of soil and provide more macro as well as micronutrients than FYM.

The lower values radish yield and soil physio-chemical properties obtained from control treatment as compared to other treatments could be because of initial low soil nutrients status that often characterizes continuous cultivation of land without fertilizer application. This finding was supported by (Hasan *et al.*, 2012)

### Root length

FYM combined with goat manure outperformed the control but was insignificant in comparison to all other treatments. The shortest root length was 14.37 cm (control) compared to 20.90 cm (FYM+GM). While all other treatments resulted in lengths ranging from 18 to 19 cm, the difference was less than the LSD value.

### Root diameter

The largest root diameter (31.43 mm) was produced by FYM coupled with poultry manure, which was statistically different from FYM alone but negligible when compared to poultry manure alone. Control was shown to be equivalent to FYM alone, whereas goat manure was found to be equivalent to chicken manure, FYM + goat dung, and poultry manure plus goat manure. The thickest root (FYM+PM) was 31.43 mm, while the thinnest was 17.32 mm (control).

Subedi *et al.* (2018) observed longer and wider roots of radish in the treatments containing PM either alone or in combination with chemical fertilizer due to higher organic manure and nutrient content (NPK and micronutrients). Also, Stephenson *et al.*, (1990) and Oladotun (2002) reported that poultry manure contained macro and micro nutrients such as N, P, K, Ca, Mg, Cu, Bo and Fe that help to increase plant biomass rapidly.

Table 5 Effect of organic manure on root weight, root length & root diameter of radish

Treatment	Root Weight(gm)	Root Length(cm)	Root Diameter(mm)
Control	31.66667 <sup>d</sup>	14.37778 <sup>b</sup>	17.3238 c
FYM	41.00000d	18.38889 <sup>ab</sup>	18.28622 <sup>c</sup>

PM	204.33333 <sup>a</sup>	19.44444 <sup>ab</sup>	26.84533 <sup>ab</sup>
GM	122.33333 <sup>b</sup>	18.41111 <sup>ab</sup>	24.91944 <sup>b</sup>
FYM+PM	111.00000 <sup>b</sup>	19.25000 <sup>ab</sup>	31.43048 <sup>a</sup>
FYM+GM	81.66667 <sup>c</sup>	20.90000 <sup>a</sup>	26.35944 <sup>ab</sup>
PM+GM	123.66667 <sup>b</sup>	18.21000 <sup>ab</sup>	24.35000 <sup>b</sup>
LSD	27.78876	5.650929	5.650929
CV%	15.27855	17.23907	13.11705
Mean	102.2381	18.42603	24.21639
SeM	15.6	3.2	3.2

Means followed by the common letter within each column are not significantly different at 5% level of significance by Duncan Test.

### **Effect of organic manure on Root yield, Harvest index, Leaf yield and Biological yield.**

#### **Harvest index**

Among the different treatments, the significantly highest harvest index of (0.7746691) was recorded in treatment T3 (Poultry manure) which was followed by T5>T7>T4>T6>T1>T2 in descending order. A minimum harvest index (0.5908403) was recorded under the treatment T2 (FYM).

Increased nutrient availability, improved soil structure and enhanced microbial activities due to addition of poultry manure could be the reason for increase in harvest index of radish. Similar finding was observed on the research conducted by (Khatri *et al.* 2019) at the horticulture farm of the Agriculture and Forestry University (AFU), Rampur, Chitwan, Nepal

#### **Leaf yield**

Among the different treatments, the significantly highest leaf yield of (59.33<sup>a</sup>) was recorded in treatment T3 (Poultry Manure) which was followed by T4>T7>T5>T6>T2>T1 in descending order. A minimum leaf yield (20.0) was recorded under the treatment of Control.

Increased nutrient availability, improved soil structure and enhanced microbial activities due to addition of poultry manure could be the reason for the increase in leaf yield of radish. Similar finding was observed on the research conducted by (Khatri *et al.*, 2019) at the horticulture farm of the Agriculture and Forestry University (AFU), Rampur, Chitwan, Nepal.

## Biological yield

Among the different treatments, the significantly highest yield of (263.67) was observed in T3 (Poultry Manure) which was followed by T4>T7>T5>T6>T2>T1 in descending order. A minimum biological yield of 51.6 was recorded under the treatment of Control.

Table 6 Effect of organic manure on root yield, harvest index, leaf yield and biological yield. of radish

Treatment	Harvest Index	Leaf Yield	Biological Yield
control	0.6098 <sup>d</sup>	20.0 <sup>e</sup>	51.67 <sup>d</sup>
FYM	0.590 <sup>d</sup>	28.33 <sup>d</sup>	69.33 <sup>d</sup>
poultry	0.774 <sup>a</sup>	59.33 <sup>a</sup>	263.67 <sup>a</sup>
Goat	0.719 <sup>bc</sup>	47.67 <sup>b</sup>	170.00 <sup>b</sup>
FYM + Poultry	0.753 <sup>ab</sup>	36.33 <sup>c</sup>	147.33 <sup>bc</sup>
FYM + Goat	0.705 <sup>c</sup>	34.00 <sup>cd</sup>	115.67 <sup>c</sup>
Poultry + Goat	0.738 <sup>abc</sup>	43.67 <sup>b</sup>	167.33 <sup>b</sup>
c.v.%	2.87	10.7159	13.688
LSD value	0.035	7.3349	34.265
mean	0.69	38.476	140.71
SEm	0.23	12.82	46.90

## Conclusion:

Tokinashi radishes show a good response to the use of various organic fertilizers, demonstrating improvements in both growth characteristics and yield performance. Among all the treatments tested, the most significant results were observed with the use of poultry manure. Growth parameters, including plant height and the number of leaves per plant, increased significantly, indicating the favorable nutrient composition and efficient absorption provided by poultry manure. Similarly, other important yield parameters, such as root weight, root length, root diameter, total root yield, harvest index, leaf yield, and biological yield, also showed superior performance under this treatment.

Research has shown that poultry manure not only provides essential nutrients but also improves soil health, creating optimal growing conditions for radish cultivation in the Lamjung District. This study highlights the potential of poultry manure as a sustainable and cost-effective organic fertilizer for radish production. It encourages its adoption among local farmers, thereby enhancing crop productivity. Future research could explore the long-term effects of poultry manure application and its impact on soil properties, crop rotations, and other vegetable crops in similar agro-climatic conditions.

## 6. REFERENCES:

1. Brintha I and Seran T H (2009). Effect of paired row planting of radish (*Raphanus sativus* L.) intercropped with vegetable amaranthus (*Amaranthus tricolor* L.) on yield components of radish in sandy regosol. *J. Agric. Sci*, 4: 19-28.
2. Chapagain, Tika Rak et al. 1970. "Comparison of Commercial and Local Varieties of Radish at Different Levels of Manures and Fertilizers." *Nepal Journal of Science and Technology* 11(September 2022): 51–56.
3. Hasan, Mohammad & Solaiman, Dr Abul. (2012). Efficacy of organic and organic fertilizer on the growth of *Brassica oleracea* L. (Cabbage). (Internal Journal of plant science)
4. Kiran, Mehwish, Muhammad Saleem Jilani, Kashif Waseem, and Muhammad Sohail. 2016. "EFFECT OF ORGANIC MANURES AND INORGANIC FERTILIZERS ON GROWTH." 29(4):363–72.
5. KITAMURA, S. (1958). Cultivars of radish and their change. In *Radish in Japan*. (Nishiyama, I. Ed.). Japanese replications of each. Society for the Promotion of Science, Tokyo, Japan. 1–19. (In Japanese).
6. Ijoyah, MO, and VL Sophie. 2009. "Effects Of Different Levels Of Decomposed Poultry Manure On Yield Of Cabbage (*Brassica Oleraceae* L.) At Anse Boileau, Seychelles." *Agro-Science* 8(1): 20–23.
7. Khatri, Keshar Bahadur, Roshan Babu Ojha, Keshab Raj Pande, and Babu Ram Khanal. 2019. "The Effects of Different Sources of Organic Manures in Growth and Yield of Radish (*Raphanus Sativus* L.)." *International Journal of Applied Sciences and Biotechnology* 7(1): 39–42.
8. Kumar, Sandeep, Sutanu Maji, Sanjay Kumar, and H.D Singh. 2014. "Efficacy of Organic Manures on Growth and Yield of Radish (*Raphanus Sativus* L.) Cv." *International Journal of Plant Sciences* 9(1): 57–60.
9. Kumar, Sanjay, Subhash Kumar Verma, Manoj Kumar, and Devendra Kumar. 2016. "Effect of Organic and Inorganic Fertilizers on Growth, Yield and Quality of Radish (*Raphanus Sativus* L.) CV. Japanese White ." *Annals of Horticulture* 9(2): 133.
10. Ministry of Finance. 2022. "Economic Survey Analysis 2021/2022." 3: 378. [https://www.mof.gov.np/uploads/document/file/1674635120\\_Economic\\_Survey\\_2022.pdf](https://www.mof.gov.np/uploads/document/file/1674635120_Economic_Survey_2022.pdf).
12. Oladotun, A. O. (2002). Managing manure as a fertilizer; Saskatchewan, Agriculture, Food and Rural Revitalization. 5 pp.
13. Shahi, Priya et al. 2022. "Effect of Different Organic Manures on Growth and Yield of Radish At Kailali District." *Journal of Wastes and Biomass Management* 4(2): 92–95.

14. Stephenson, A. H., McCaskey T. A. and Ruffin B. G. (1990). A survey of broiler litter composition and potential value as a nutrient resource. *Biological wastes* 34: 1-9.
15. Umar, U. M., I. Ibrahim, Iro, and S. M. Obidola. 2019. "Growth and Yield of Radish (*Raphanus Sativus* L.) as Influenced by Different Levels of Kalli Organic Fertilizer on the Jos Plateau." *Asian Journal of Research in Crop Science* (September): 1-8.

UNDER PEER REVIEW