

# Diversity of Avifauna in the campus of Navsari Agricultural University, Navsari, Gujarat, India

## ABSTRACT:

Educational institutions are important green spaces which reside diverse species of organisms. In this study, a checklist of bird species was compiled using data collected over seven years (2014-2020) from the campus of Navsari Agricultural University (NAU), Navsari, Gujarat. The study area comprises various habitats including agriculture lands, horticultural orchards, agroforestry plantations, monoculture plantations, biodiversity park, arboretum, grasslands, barren lands and water bodies. A total of 169 bird species belonging to 19 orders, 63 families, and 126 genera were documented throughout the study period. Among the recorded species two (Asian woolly necked stork *Ciconia episcopus* and Alexandrine Parakeet *Psittaculaeupatria*) are near threatened and three (Greater Spotted Eagle *Clanga clanga*, Sarus Crane *Antigone Antigone* and River Tern *Sterna aurantia*) are vulnerable. Family Muscicapidae was most diverse within study area comprising 12 species. Species feeding on invertebrates were predominant. The substantial species diversity and presence of nesting activity emphasize the importance of university campuses as significant habitat for avian conservation. Present study provides crucial baseline data for further scientific studies on Avifauna of NAU campus.

**Keywords:** *Avian diversity, Urban green space, Agroecosystems, Educational Institutes, Agricultural university*


## INTRODUCTION:

India is one of the most biodiverse countries and has a forest cover of 25.17% of its geographical area (ISFR 2023). The forest cover of the country includes 3.41% of tree cover, which falls outside the forest area. A considerable part of this tree cover is contributed by the educational institutes, public parks, roadside plantations, farm forests etc. Whilst urbanization is posing an increasing threat to biodiversity (Li et al., 2022). These premises act as important habitats amidst oases of concrete forests. These green spaces act as a source of food, water, and shelter for biodiversity. Several studies have highlighted the role of institutional campuses as home of various species (Colding & Barthel, 2017, Liu et al. 2021, Guthula et al. 2022). Educational institutions traditionally established in natural areas away from urban settlements are known to reside diverse flora and fauna. Many educational campuses in India have carried out study on the diversity of birds (Adhurya et al., 2023; Chandrakar & Dhuria, 2019; Jain et al., 2005; Kiran et al., 2022; Kumar et al., 2024; Patel et al., 2021; Rathod & Bhaduri, 2022; Shivhare et al., 2022; K. Singh, 2018; R. Singh et al., 2024)

Navsari Agriculture University lies in South Gujarat is spread over 333.42 hectares. It encompasses different land use types including agriculture lands, horticultural orchards, agroforestry plantations, monoculture plantations, biodiversity park (~140 plant species), arboretum (~162 tree species), grasslands, barren lands and water bodies. The rich plant diversity and various habitats provide ample space for different organisms. 89 species of butterflies are recorded from the campus (Malek et al., 2024). Agroecosystems are identified as important habitats and prominent nesting grounds for birds (Borad et al., 2000, 2001). With increasing urbanisation pressure identifying the role of institution campuses for biodiversity conservation is important. Given this background our study aims to provide a checklist of bird species found in the campus of Navsari Agricultural University (NAU).

## MATERIAL AND METHODS

Navsari Agricultural University (NAU) is situated in district Navsari of Gujarat. NAU was established in 1965 and situated at 20.9248° N latitude, and 72.9079° E longitude at 10 meters above sea level (MSL). The campus is located 12 km away from the east of the Arabian seashore. The region experiences a warm and humid climate, with an average rainfall of approximately 1500 mm (June to September). Winters (November to February) are moderately cold, whereas summers (March to May) are characterized by hot and humid conditions.



**Figure 1: Study Map of the Navsari Agricultural University, Navsari**

This study is a compilation of 7 years of birding surveys. During the study, different trails of the campus were regularly visited in the seasons of winter, summer and monsoon. At least two visits in each month were ensured throughout the study period. The various habitats include agriculture lands, horticultural orchards, agroforestry plantations, monoculture plantations, biodiversity park (~140 plant species), arboretum (~162 tree species), grasslands, barren lands and water bodies. Since the NAU campus region lacks extensive homogenous areas of a single habitat, systematic transects could not be laid. As a result, pre-existing trails were followed for the bird survey. The survey was carried out in the morning and evening mostly 6.45-9.30 hrs and 16.00-18.00 hrs during peak activity time of birds. Apart from a systematic survey, we have included the opportunistic sightings of birds observed on campus. We observed birds with the aid of 12 X 50 binoculars and identified them using field guides of Ali (2002); Rasmussen & Anderton (2012) and Grimmett et al. (2011). Unidentified birds during field visits were identified from the photographs taken using a DSLR camera. The taxonomy of the given checklist is followed using Praveen et al. (2016); Praveen et al. (2020); Praveen and Jaypal (2023) and Praveen and Jaypal (2024). We determined whether a species was resident or migratory in the study area following Ganpule (2020) and feeding guilds following (Wilman et al., 2014). We also evaluated the threat level to species using the Red List prepared by the International Union for Conservation of Nature (IUCN 2024).

## RESULTS AND DISCUSSION

We recorded a total of 169 bird species belonging to 19 orders, 63 families, and 126 genera throughout the study period (Table 1). We recorded the highest number of species in order of Passeriformes (89), followed by Pelecaniformes (13) and Charadriiformes (10). In present study area,

the family Muscicapidae (12) was most prevalent, followed by Ardeidae (9) and Accipitridae (8) (Figure. 2). Of the recorded species, 112 bird species are residents followed by 44 winter migrants, 5 bird species residents as well as winter migrants and 3 bird species monsoon breeding. As per the IUCN category, Asian woolly necked stork *Ciconia episcopus* and Alexandrine Parakeet *Psittacula eupatria* are near threatened, and Greater Spotted Eagle *Clanga clanga*, Sarus Crane *Antigone Antigone* and River Tern *Sterna aurantia* are vulnerable. Among the feeding guild, 48% (81) of documented bird species consume invertebrates, while 17% (29) of species feed on plant parts and seeds. Additionally, 13% (22) of species each feed on terrestrial vertebrates, fish, and carrion/carcasses, and omnivores, and 9% (15) of bird species feed on fruits and nectar. (Figure. 3).

**Table 1: Checklist of the avifauna of Navsari Agricultural University (Patel et al, 2022)**

	Species	IUCN Status (2024)	Status	Feeding guild
<b>Order: Galliformes</b>				
<b>Family: Phasianidae</b>				
1.	Grey Francolin <i>Ortygornis pondicerianus</i> (J.F. Gmelin, 1789)	LC	R	PS
2.	Indian Peafowl <i>Pavo cristatus</i> Linnaeus, 1758	LC	R	PS
<b>Order: Anseriformes</b>				
<b>Family: Anatidae</b>				
3.	Indian Spot-Billed Duck <i>Anas poecilorhyncha</i> J.R. Forster, 1781	LC	W	PS
<b>Order: Podicipediformes</b>				
<b>Family: Podicipedidae</b>				
4.	Little Grebe <i>Tachybaptus ruficollis</i> (Pallas, 1764)	LC	R	I
<b>Order: Ciconiiformes</b>				
<b>Family: Ciconiidae</b>				
5.	Asian woolly necked stork <i>Ciconia episcopus</i> (Boddaert, 1783)	NT	R	VFC
6.	Painted Stork <i>Mycteria leucocephala</i> (Pennant, 1769)	LC	MM	VFC
7.	Asian Openbill <i>Anastomus oscitans</i> (Boddaert, 1783)	LC	MM	I
<b>Order: Pelecaniformes</b>				
<b>Family: Threskiornithidae</b>				
8.	Black-Headed Ibis <i>Threskiornis melanocephalus</i> (Latham, 1790)	LC	R	VFC
9.	Indian Black Ibis/ Red-Naped Ibis <i>Pseudibis papillosa</i> (Temminck, 1824)	LC	R	OM
10.	Glossy Ibis <i>Plegadis falcinellus</i> (Linnaeus, 1766)	LC	R, W	I
11.	Eurasian Spoonbill <i>Platalea leucorodia</i> Linnaeus, 1758	LC	R	I
<b>Family: Ardeidae</b>				
12.	Yellow Bittern <i>Ixobrychus sinensis</i> (J.F. Gmelin, 1789)	LC	MM	I
13.	Black Crowned Night Heron <i>Nycticorax nycticorax</i> (Linnaeus, 1758)	LC	R	VFC
14.	Indian Pond Heron <i>Ardeola grayii</i> (Sykes, 1832)	LC	R	OM
15.	Grey Heron <i>Ardea cinerea</i> Linnaeus, 1758	LC	R	VFC
16.	Purple Heron <i>Ardea purpurea</i> Linnaeus, 1766	LC	R	VFC
17.	Cattle Egret <i>Bubulcus ibis</i> (Linnaeus, 1758)	LC	R	I
18.	Great Egret <i>Ardea alba</i> Linnaeus, 1758	LC	R	VFC
19.	Intermediate Egret <i>Ardea intermedia</i> Wagler, 1829	LC	R	VFC
20.	Little Egret <i>Egretta garzetta</i> (Linnaeus, 1766)	LC	R	I
<b>Order: Suliformes</b>				
<b>Family: Phalacrocoracidae</b>				
21.	Little Cormorant <i>Microcarboniger</i> (Vieillot, 1817)	LC	R	VFC
<b>Order: Falconiformes</b>				
<b>Family: Falconidae</b>				
22.	Common Kestrel <i>Falco tinnunculus</i> Linnaeus, 1758	LC	W	VFC
<b>Order: Accipitriformes</b>				

<b>Family: Accipitridae</b>				
23.	Black Winged Kite <i>Elanus caeruleus</i> (Desfontaines, 1789)	LC	W	VFC
24a.	Black Kite <i>Milvus migrans</i> (Boddaert, 1783)	LC	R	VFC
24b.	Black-Eared Kite <i>Milvus migrans lineatus</i> (Boddaert, 1783)	LC	W	VFC
25	Oriental Honey Buzzard <i>Pernis ptilorhynchus</i> (Temminck, 1821)	LC	W	I
26.	Western Marsh Harrier <i>Circus aeruginosus</i> (Linnaeus, 1758)	LC	W	VFC
27.	Shikra <i>Accipiter badius</i> (J.F. Gmelin, 1788)	LC	R	VFC
28.	Eurasian Sparrowhawk <i>Accipiter nisus</i> (Linnaeus, 1758)	LC	W	VFC
29.	White-Eyed Buzzard <i>Butasturteesa</i> (Franklin, 1831)	LC	W	VFC
30.	Greater Spotted Eagle <i>Clanga clanga</i> (Pallas, 1811)	VU	W	VFC
<b>Order: Gruiformes</b>				
<b>Family: Gruidae</b>				
31.	Common Crane <i>Grus grus</i> (Linnaeus, 1758)	LC	W	PS
32.	Demoiselle Crane <i>Grus virgo</i> (Linnaeus, 1758)	LC	W	PS
33.	Sarus Crane <i>Antigone antigone</i> (Linnaeus, 1758)	VU	R	PS
<b>Family: Rallidae</b>				
34.	Common Moorhen <i>Gallinula chloropus</i> (Linnaeus, 1758)	LC	R	OM
35.	White Breasted Waterhen <i>Amaurornis phoenicurus</i> (Pennant, 1769)	LC	R	I
36.	Common Coot <i>Fulica atra</i> Linnaeus, 1758	LC	R, W	PS
37.	Purple Swampphen <i>Porphyrioporphyrus</i> (Linnaeus, 1758)	LC	R	PS
38.	Ruddy breasted crane <i>Zapornia fusca</i> (Linnaeus, 1766)	LC	R	PS
<b>Order: Charadriiformes</b>				
<b>Family: Turnicidae</b>				
39.	Barred Buttonquail <i>Turnix suscitator</i> (J.F. Gmelin, 1789)	LC	MM	PS
<b>Family: Jacanidae</b>				
40.	Pheasant-tailed Jacana <i>Hydrophasianus chirurgus</i> (Scopoli, 1786)	LC	R	I
41.	Bronze Winged Jacana <i>Metopidius indicus</i> (Latham, 1790)	LC	R	PS
<b>Family: Recurvirostridae</b>				
42.	Black Winged Stilt <i>Himantopus himantopus</i> (Linnaeus, 1758)	LC	W	I
<b>Family: Charadriidae</b>				
43.	Red Wattled Lapwing <i>Vanellus indicus</i> (Boddaert, 1783)	LC	R	I
<b>Family: Scolopacidae</b>				
44.	Wood Sandpiper <i>Tringaglareola</i> Linnaeus, 1758	LC	W	I
45.	Common Sandpiper <i>Actitis hypoleucos</i> (Linnaeus, 1758)	LC	W	OM
46.	Common Snipe <i>Gallinagogallinago</i> (Linnaeus, 1758)	LC	W	I
<b>Family: Laridae</b>				
47.	River Tern <i>Sterna aurantia</i> J.E. Gray, 1831	VU	R	OM
48.	Whiskered Tern <i>Chlidonias hybrid</i> (Pallas, 1811)	LC	W	VFC
<b>Order: Columbiformes</b>				
<b>Family: Columbidae</b>				
49.	Rock Pigeon <i>Columba livia</i> J.F. Gmelin, 1789	LC	R	PS
50.	Eurasian Collared Dove <i>Streptopelia decaocto</i> (Frivaldszky, 1838)	LC	R	PS
51.	Red Collared Dove <i>Streptopelia tranquebarica</i> (Hermann, 1804)	LC	R	PS
52.	Spotted Dove <i>Spilopelia chinensis</i> (Scopoli, 1786)	LC	R	PS
53.	Laughing Dove <i>Spilopelia senegalensis</i> (Linnaeus, 1766)	LC	R	PS
54.	Yellow Legged Green Pigeon/ Yellow Footed Green Region <i>Treron phoenicopterus</i> (Latham, 1790)	LC	R	FN
<b>Order: Psittaciformes</b>				
<b>Family: Psittaculidae</b>				
55.	Alexandrine Parakeet <i>Psittacula eupatria</i> (Linnaeus, 1766)	NT	R	FN

56.	Rose Ringed Parakeet <i>Psittaculakrameri</i> (Scopoli, 1769)	LC	R	FN
57.	Plum headed parakeet <i>Psittaculacyanocephala</i> (Linnaeus, 1766)	LC	R	FN
<b>Order: Cuculiformes</b>				
<b>Family: Cuculidae</b>				
58.	Pied Cuckoo/ Jacobin Cuckoo <i>Clamatorjacobinus</i> (Boddaert, 1783)	LC	MB	I
59.	Common Hawk Cuckoo <i>Hierococcyxvarius</i> (Vahl, 1797)	LC	R	I
60.	Common Cuckoo/ Eursaian Cuckoo <i>Cuculuscanorus</i> Linnaeus, 1758	LC	MB	I
61.	Grey Bellied Cuckoo <i>Cacomantispasserinus</i> (Vahl, 1797)	LC	R	I
62.	Asian Koel <i>Eudynamysscolopaceus</i> (Linnaeus, 1758)	LC	R	FN
63.	Greater Coucal/ Southern Coucal/ Crow Pheasant <i>Centropus sinensis parroti</i> (Stephens, 1815)	LC	R	OM
<b>Order: Strigiformes</b>				
<b>Family: Tytonidae</b>				
64.	Common Barn Owl/ Barn Owl <i>Tyto alba</i> (Scopoli, 1769)	LC	R	VFC
<b>Family: Strigidae</b>				
65.	Spotted Owlet <i>Athene brama</i> (Temminck, 1821)	LC	R	I
<b>Order: Caprimulgiformes</b>				
<b>Family: Apodidae</b>				
66.	Asian Palm Swift <i>Cypsiurusbalasiensis</i> (J.E. Gray, 1829)	LC	R	I
67.	Indian House Swift/Little Swift <i>Apus affinis</i> (J.E. Gray, 1830)	LC	R	I
<b>Order: Bucerotiformes</b>				
<b>Family: Upupidae</b>				
68.	Common Hoopoe <i>Upupa epops</i> Linnaeus, 1758	LC	R, W	I
<b>Family: Bucerotidae</b>				
69.	Indian Grey Hornbill <i>Ocyerosbirostris</i> (Scopoli, 1786)	LC	R	FN
<b>Order: Coraciiformes</b>				
<b>Family: Coraciidae</b>				
70.	Indian Roller <i>Coracias benghalensis</i> (Linnaeus, 1758)	LC	R	OM
71.	European Roller <i>Coracias garrulus</i> Linnaeus, 1758	LC	PM	I
<b>Family: Alcedinidae</b>				
72.	White Throated Kingfisher <i>Halcyon smyrnensis</i> (Linnaeus, 1758)	LC	R	I
73.	Pied Kingfisher <i>Ceryle rudis</i> (Linnaeus, 1758)	LC	R	VFC
74.	Common Kingfisher <i>Alcedo atthis</i> (Linnaeus, 1758)	LC	R	VFC
<b>Family: Meropidae</b>				
75.	Green Bee Eater <i>Meropsorientalis</i> Latham, 1801	LC	R	I
76.	Blue Tailed Bee Eater <i>Meropsphilippinus</i> Linnaeus, 1767	LC	R	I
<b>Order: Piciformes</b>				
<b>Family: Megalaimidae</b>				
77.	Coppersmith Barbet <i>Psilopogonhaemacephalus</i> (Statius Muller, 1776)	LC	R	FN
<b>Family: Picidae</b>				
78.	Eurasian Wryneck <i>Jynx torquilla</i> Linnaeus, 1758	LC	W	I
79.	Lesser Golden-Backed Woodpecker <i>Dinopiumbenghalense</i> (Linnaeus, 1758)	LC	R	I
80.	Yellow crowned woodpecker/ Yellow-fronted Pied Woodpecker <i>Dendrocoposmahrattensis</i> (Latham, 1801)	LC	R	I
<b>Order: Passeriformes</b>				
<b>Family: Acrocephalidae</b>				
81.	Booted Warbler <i>Iduna caligata</i> (M.H.C. Lichtenstein, 1823)	LC	W	I
82.	Clamorous Reed Warbler <i>Acrocephalus stentoreus</i> (Hemprich& Ehrenberg, 1833)	LC	R	I
83.	Sykes's Warbler <i>Iduna rama</i> (Sykes, 1832)	LC	W	I

<b>Family: Aegithinidae</b>				
84.	Common Iora <i>Aegithina tiphia</i> (Linnaeus, 1758)	LC	R	I
<b>Family: Alaudidae</b>				
85.	Ashy Crowned Sparrow Lark <i>Eremopterix griseus</i> (Scopoli, 1786)	LC	R	PS
86.	Crested Lark <i>Galerida cristata</i> (Linnaeus, 1758)	LC	R	PS
87.	Rufous Tailed Lark <i>Ammomanes phoenicurus</i> (Franklin, 1831)	LC	R	OM
88.	Sykes's Lark <i>Galerida deva</i> (Sykes, 1832)	LC	R	PS
<b>Family: Campephagidae</b>				
89.	Small minivet <i>Pericrocotus cinnamomeus</i> (Linnaeus, 1766)	LC	R	I
<b>Family: Chloropseidae</b>				
90.	Golden Fronted Leafbird <i>Chloropsis aurifrons</i> (Temminck, 1829)	LC	R	FN
91.	Jerdon's Leafbird <i>Chloropsis jerdoni</i> (Blyth, 1844)	LC	R	FN
<b>Family: Cisticolidae</b>				
92.	Ashy Prinia <i>Prinia socialis</i> Sykes, 1832	LC	R	I
93.	Common Tailorbird <i>Orthotomus sutorius</i> (Pennant, 1769)	LC	R	I
94.	Grey Breasted Prinia <i>Prinia hodgsonii</i> Blyth, 1844	LC	R	I
95.	Jungle Prinia <i>Prinia sylvatica</i> Jerdon, 1840	LC	R	I
96.	Plain Prinia <i>Prinia inornata</i> Sykes, 1832	LC	R	I
97.	Zitting Cisticola <i>Cisticola juncidis</i> (Rafinesque, 1810)	LC	R	I
<b>Family: Corvidae</b>				
98.	House Crow <i>Corvus splendens</i> Vieillot, 1817	LC	R	VFC
99.	Jungle Crow / Large -Billed Crow <i>Corvus macrorhynchos</i> Wagler, 1827	LC	R	FN
<b>Family: Dicaeidae</b>				
100.	Pale Billed Flowerpecker <i>Dicaeum erythrorhynchos</i> (Latham, 1790)	LC	R	FN
101.	Thick Billed Flowerpecker <i>Dicaeum agile</i> (Tickell, 1833)	LC	R	FN
<b>Family: Dicrurus</b>				
102.	Ashy Drongo <i>Dicrurus leucophaeus</i> Vieillot, 1817	LC	W	I
103.	Black Drongo <i>Dicrurus macrocercus</i> Vieillot, 1817	LC	R	OM
104.	White Bellied Drongo <i>Dicrurus caerulescens</i> (Linnaeus, 1758)	LC	R, W	I
<b>Family: Emberizidae</b>				
105.	Black Headed Bunting <i>Emberiza melanocephala</i> (Scopoli, 1769)	LC	W	PS
106.	Crested Bunting <i>Emberiza lathami</i> (J.E. Gray, 1831)	LC	R	PS
107.	Red Headed Bunting <i>Emberiza bruniceps</i> (von Brandt, 1841)	LC	W	PS
<b>Family: Estrildidae</b>				
108.	Indian Silverbill <i>Euodice malabarica</i> (Linnaeus, 1758)	LC	R	PS
109.	Red Munia <i>Amandava amandava</i> (Linnaeus, 1758)	LC	R	PS
110.	Scaly Breasted Munia <i>Lonchurapunctulata</i> (Linnaeus, 1758)	LC	R	PS
111.	Tricoloured Munia <i>Lonchura malacca</i> (Linnaeus, 1766)	LC	R	PS
<b>Family: Fringillidae</b>				
112.	Common Rosefinch <i>Carpodacus erythrinus</i> (Pallas, 1770)	LC	W	PS
<b>Family: Hirundinidae</b>				
113.	Red-rumped Swallow <i>Cecropis daurica</i> (Laxmann, 1769)	LC	R	I
114.	Wire Tailed Swallow <i>Hirundo smithii</i> Leach, 1818	LC	R	I
<b>Family: Laniidae</b>				
115.	Bay Backed Shrike <i>Lanius vittatus</i> Valenciennes, 1826	LC	R	I
116.	Isabelline Shrike <i>Lanius isabellinus</i> Hemprich & Ehrenberg, 1833	LC	W	I
117.	Long Tailed Shrike <i>Lanius schach</i> Linnaeus, 1758	LC	R	I
<b>Family: Leiothrichidae</b>				

118.	Common Babbler <i>Argya caudata</i> (Dumont, 1823)	LC	R	OM
119.	Jungle Babbler <i>Turdoides striata</i> (Dumont, 1823)	LC	R	OM
<b>Family: Monarchidae</b>				
120.	Indian Paradise Flycatcher <i>Terpsiphone paradisi</i> (Linnaeus, 1758)	LC	R	I
121.	Black-naped Monarch <i>Hypothymis azurea</i> (Boddaert, 1783)	LC	R	I
<b>Family: Motacillidae</b>				
122.	Citrine Wagtail <i>Motacillactreola</i> Pallas, 1776	LC	W	I
123.	Long Billed Pipit <i>Anthus similis</i> (Jerdon, 1840)	LC	W, R?	I
124.	Paddy field Pipit <i>Anthus rufulus</i> Vieillot, 1818	LC	R	I
125.	Tawny Pipit <i>Anthus campestris</i> (Linnaeus, 1758)	LC	W	I
126.	Tree Pipit <i>Anthus trivialis</i> (Linnaeus, 1758)	LC	W	I
127.	Western Yellow Wagtail <i>Motacilla flava</i> Linnaeus, 1758	LC	W	I
128.	White Wagtail <i>Motacilla alba</i> Linnaeus, 1758	LC	W	I
<b>Family: Muscicapidae</b>				
129.	Asian Brown Flycatcher <i>Muscicapa dauurica</i> Pallas, 1811	LC	R, W	I
130.	Black Redstart <i>Phoenicurus ochruros</i> (S.G. Gmelin, 1774)	LC	W	I
131.	Bluethroat <i>Luscinia svecica</i> (Linnaeus, 1758)	LC	W	I
132.	Brown Rock Chat <i>Oenanthe fusca</i> (Blyth, 1851)	LC	R	I
133.	Indian Robin <i>Saxicoloides fulicatus</i> (Linnaeus, 1766)	LC	R	I
134.	Isabelline Wheatear <i>Oenanthe isabellina</i> (Temminck, 1829)	LC	W	I
135.	Oriental Magpie Robin <i>Copsychus saularis</i> (Linnaeus, 1758)	LC	R	I
136.	Pied Bush chat <i>Saxicola caprata</i> (Linnaeus, 1766)	LC	R	I
137.	Red Breasted Flycatcher <i>Ficedula parva</i> (Bechstein, 1792)	LC	W	I
138.	Siberian Stonechat <i>Saxicola torquatus</i> (Pallas, 1773)	LC	W	I
139.	Taiga Flycatcher <i>Ficedula albicilla</i> (Pallas, 1811)	LC	W	I
140.	Tickell's Blue Flycatcher <i>Cyornis tickelliae</i> Blyth, 1843	LC	R	I
<b>Family: Nectariniidae</b>				
141.	Purple Rumped Sunbird <i>Leptocomazeylonica</i> (Linnaeus, 1766)	LC	R	FN
142.	Purple Sunbird <i>Cinnyris asiaticus</i> (Linnaeus, 1766)	LC	R	FN
<b>Family: Oriolidae</b>				
143.	Indian Golden Oriole <i>Oriolus kundoo</i> Sykes, 1832	LC	R	OM
<b>Family: Paridae</b>				
144.	Great Tit / Cinereous Tit <i>Parus cinereus</i> Vieillot, 1818	LC	W	I
<b>Family: Passeridae</b>				
145.	House Sparrow <i>Passer domesticus</i> (Linnaeus, 1758)	LC	R	PS
146.	Yellow-Throated Sparrow <i>Gymnoris xanthocollis</i> (E. Burton, 1838)	LC	R	OM
<b>Family: Phylloscopidae</b>				
147.	Common Chiffchaff <i>Phylloscopus collybita</i> (Vieillot, 1817)	LC	W	I
148.	Greenish Leaf Warbler <i>Phylloscopus trochiloides</i> (Sundevall, 1837)	LC	W	I
149.	Sulphur Bellied Warbler <i>Phylloscopus griseolus</i> Blyth, 1847	LC	W	I
<b>Family: Pittidae</b>				
150.	Indian pitta <i>Pittabrachyura</i> (Linnaeus, 1766)	LC	MB	I
<b>Family: Ploceidae</b>				
151.	Baya Weaver <i>Ploceus philippinus</i> (Linnaeus, 1766)	LC	R	PS
<b>Family: Pycnonotidae</b>				
152.	Red Vented Bulbul <i>Pycnonotus cafer</i> (Linnaeus, 1766)	LC	R	OM
153.	Red Whiskered Bulbul <i>Pycnonotus jocosus</i> (Linnaeus, 1758)	LC	R	OM
154.	White Browed Bulbul <i>Pycnonotus luteolus</i> (Lesson, 1841)	LC	R	FN
<b>Family: Rhipiduridae</b>				

155.	White Browed Fantail <i>Rhipidura aureola</i> Lesson, 1831	LC	R	I
156.	White Spotted Fantail <i>Rhipidura albogularis</i> (Vieillot, 1818)	LC	R	I
<b>Family: Stenostiridae</b>				
157.	Grey headed canary flycatcher <i>Culicicapaceylonensis</i> (Swainson, 1820)	LC	W	I
<b>Family: Sturnidae</b>				
158.	Bank Myna <i>Acridotheres ginginianus</i> (Latham, 1790)	LC	R	OM
159.	Brahminy Starling <i>Sturniapagodarum</i> (J.F. Gmelin, 1789)	LC	R	OM
160.	Chestnut-Tailed Starling <i>Sturniamalabarica</i> (J.F. Gmelin, 1789)	LC	R	OM
161.	Common Myna <i>Acridotheres tristis</i> (Linnaeus, 1766)	LC	R	OM
162.	Jungle Myna <i>Acridotheres fuscus</i> (Wagler, 1827)	LC	R	OM
163.	Malabar Starling <i>Sturniamalabarica</i> (Jerdon, 1845)	LC	W	OM
164.	Rosy Starling <i>Pastor roseus</i> (Linnaeus, 1758)	LC	W	I
<b>Family: Sylviidae</b>				
165.	Lesser Whitethroat <i>Sylvia curruca</i> (Linnaeus, 1758)	LC	W	I
166.	Yellow Eyed Babbler <i>Chrysommasinense</i> (J.F. Gmelin, 1789)	LC	R	I
<b>Family: Timaliidae</b>				
167.	Tawny Bellied Babbler <i>Dumetiahyperythra</i> (Franklin, 1831)	LC	R	I
<b>Family: Vangidae</b>				
168.	Common Wood shrike <i>Tephrodornispondicerianus</i> (J.F. Gmelin, 1789)	LC	R	I
<b>Family: Zosteropidae</b>				
169.	Oriental White Eye <i>Zosterops palpebrosus</i> (Temminck, 1824)	LC	W	OM

(IUCN Red List categories: EN: Endangered, LC: Least concern, NT: Near Threatened, VU: Vulnerable, DD: Data Deficient; **Residential Status:** R: Resident/ Resident Breeding, MB: Monsoon Breeding, MM: Monsoon Migrant, W: Winter Migrant, PM: Passage Migrant, V: Vagrant and **Feeding guilds:** FN: Fruitnect, I: Invertebrate, OM: Omnivore, PS: PlantSeed and VFC: VertFishScav)





Figure 2: Family and order-wise representation of Avifauna of Navsari Agricultural University, Navsari.



**Figure 3: Feeding guild representation of avifauna of Navsari Agricultural University, Navsari** (*FruitNect* - feeding on fruits and nectar, *Invertebrate* - feeding on aquatic and terrestrial invertebrates, *PlantSeed* - feeding on plant parts and seeds, *VertFishScav* - feeding on vertebrates, fishes and scavenging on carrion).

This study highlights the role of institutional campuses as important places that retain significant biodiversity. Navsari is an important region under the Central Asian flyway with rich avian diversity (Patel et al 2022). The cover of agricultural fields, orchards and plantations which give food and spatial resources attracts diverse species of birds. Some noteworthy migratory species in campus are Indian Pitta *Pittabrachyura*, Pied Cuckoo *Clamatorjacobinus*, Eurasian Cuckoo *Cuculuscanorus*, Yellow Bittern *Ixobrychus sinensis*, Asian Openbill *Anastomusoscitans*, Painted Stork *Mycteria leucocephala*, Taiga Flycatcher *Ficedula albicilla*, Malabar Starling *Sturniablythii*, Isabelline Wheatear *Oenanthe isabellina*, Grey-headed Canary-Flycatcher *Culicicapaceylonensis* and Eurasian Sparrowhawk *Accipiter nisus* among others. Many of these species are not easily encountered in surrounding areas. Sanllorente et al. (2023) have observed more species richness and varying species composition between campus areas and surrounding urban areas. Further study in NAU and its surroundings would help to identify such a pattern which could determine the important role of campus for retaining more functional and phylogenetic diversity of birds.

Well-wooded areas like biodiversity park, arboretum and uncultivated areas on campus are hotspots for bird diversity. Forest specialist birds like Indian Paradise-Flycatcher *Terpsiphone paradisi*, Indian Grey Hornbill *Ocyerosbirostris* and Common Woodshrike *Tephrodornispondicerianus* were observed in these areas. Several species are found to breed on the campus some peculiar to them are Indian Paradise-Flycatcher, Indian Grey Hornbill, Yellow Bittern, Pheasant-tailed Jacana *Hydrophasianuschirurgus*, Black Drongo *Dicrurus macrocercus*, Common Woodshrike and Spotted Owllet *Athene brama*. A Comprehensive study on the biodiversity of ~300 international campuses shows an average of 66 bird species present on the premises of one particular campus (Liu et al., 2021). The study by Guthula et al. (2022) on 335 campuses in India shows an average of 88 species per campus. Our study documented 170 bird species which is considerably more than campus averages. Low averages in given studies might be attributed to the wide geographic area considered in the study. The reason for more richness in NAU could be the longer study period. The study carried out by Adhurya et al. (2023) in West Bengal recorded 106 species in a small area of a 12-hectare campus during the same 7-year period. The other reason could be the vicinity of NAU near the sea which provides year-round moderate climatic conditions for various species to thrive. Moreover, as the universities introduce and raise many diverse species of plants and actively conserve them, (In NAU, biodiversity park and Arboretum) it provides foraging and nesting places for bird species.

Change in land use patterns due to urbanisation severely impacts the birds in the tropics (Jetz et al., 2007). Urbanisation around the campus has been accelerating for a few years, which could pose a serious threat in the future. In addition, land use change on campus for developmental purposes can pose a risk to various habitats of birds. Therefore, mindful developmental works should be planned for the campus area. As the pressure on overall biodiversity rises, identifying secure refugia for different species in urban environments and taking proactive steps to conserve them could significantly aid in global conservation initiatives (Knapp et al., 2020). The present checklist on avifaunal diversity of NAU campus contributes to such conservation efforts.

## CONCLUSION

The NAU campus harbours 169 species of birds and many other organisms. The rich diversity in the campus is attributed to presence of diverse habitats. The presence of rich biodiversity influences positively to students as it can be a valuable natural resource to observe and learn. The present findings on avian diversity can be used as a baseline for further research on studying avian species abundance and composition. Long-term scientific study on avian species in the study area can be continued, with a focus on habitat use, seasonal abundance, and nesting and breeding ecology. This would highlight the significance of the institutional campus of NAU in supporting and conserving biodiversity.

## Ethical Approval

Animal Ethic committee approval has been collected and preserved by the author(s)

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## REFERENCES

- Adhurya, S., Gayen, D., Chakrabarty, M., & Singha Roy, U. (2023). A study of avian diversity in Durgapur Government College Campus, West Bengal, India. *The Holistic Approach to Environment*, 13(2), 48–62.
- Ali, S., 2002. The book of Indian birds (13th Edn.). Bombay Natural History Society, Oxford University Press, New Delhi.
- Borad, C. K., Mukherjee, A., & Parasharya, B. M. (2000). Conservation of the avian biodiversity in paddy (*Oryza sativa*) crop agroecosystem. *The Indian Journal of Agricultural Sciences*, 70(4), Article 4.
- Borad, C. K., Mukherjee, A., & Parasharya, B. M. (2001). Damage potential of Indian sarus crane in paddy crop agroecosystem in Kheda district Gujarat, India. *Agriculture, Ecosystems & Environment*, 86(2), 211–215.
- Chandrakar, A. K., & Dhuria, S. (2019). A Checklist of bird diversity In Guru Ghasidas University campus, Bilaspur, Chhattisgarh, India. *Eco Chronicle*, 14, 53–59.
- Colding, J., & Barthel, S. (2017). The Role of University Campuses in Reconnecting Humans to the Biosphere. *Sustainability*, 9(12), Article 12.
- Ganpule, P., 2020. A checklist of the birds of Gujarat. Bird Conservation Society, Gujarat, Ahmedabad.
- Grimmett R, Inskipp C and Inskipp T. 2011. Birds of the Indian Subcontinent. Oxford University Press, Delhi
- Guthula, V. B., Shrotriya, S., Nigam, P., Goyal, S. P., Mohan, D., & Habib, B. (2022). Biodiversity significance of small habitat patches: More than half of Indian bird species are in academic campuses. *Landscape and Urban Planning*, 228, 104552.
- India State of Forests Report 2023. Forest Survey of India. Dehradun, Uttarakhand, India.
- IUCN. 2024. The IUCN Red List of Threatened Species. Version 2024-2. <https://www.iucnredlist.org>. Accessed on 2<sup>nd</sup> January 2024.
- Jain, N. K., Patel, S. N., & Patel, M. V. (2005). Birds of Gujarat University Campus, Ahmedabad. *Zoos' Print Journal*, 20(12), 2111–2113.

Jetz, W., Wilcove, D. S., & Dobson, A. P. (2007). Projected Impacts of Climate and Land-Use Change on the Global Diversity of Birds. *PLOS Biology*, 5(6), e157.

Kiran, Singh, D., Kour, A., Dahiya, P., Delu, V., & Kumar, R. (2022). Community composition and status of avian diversity at Campus and Agricultural landscapes of Chaudhary Charan Singh Haryana Agricultural University, Hisar (Haryana). *Journal of Applied and Natural Science*, 14(4), 1130–1140.

Knapp, S., Aronson, M.F., Carpenter, E., Herrera-Montes, A., Jung, K., Kotze, D.J., La Sorte, F.A., Lepczyk, C.A., MacGregor-Fors, I., MacIvor, J.S. and Moretti, M., (2021). A research agenda for urban biodiversity in the global extinction crisis. *BioScience*, 71(3), pp.268-279.

Kumar, P., Parmar, B., & Kumar, P. (2024). Checklist and comparison of the bird diversity from the Himachal Pradesh Agricultural University, India. *Journal of Threatened Taxa*, 16(4), Article 4.

Li, G., Fang, C., Li, Y., Wang, Z., Sun, S., He, S., Qi, W., Bao, C., Ma, H., Fan, Y., Feng, Y., & Liu, X. (2022). Global impacts of future urban expansion on terrestrial vertebrate diversity. *Nature Communications*, 13(1), 1628.

Liu, J., Zhao, Y., Si, X., Feng, G., Slik, F., & Zhang, J. (2021). University campuses as valuable resources for urban biodiversity research and conservation. *Urban Forestry & Urban Greening*, 64, 127255.

Malek, S., Patel, M., & Desai, B. (2024). Butterfly Diversity of Navsari Agricultural University (NAU), Navsari, Gujarat, India. *Journal of Advances in Biology & Biotechnology*, 27(9), 44–53.

Patel M, Kapdi P, Tandel N, Desai J, PatelZ (2022). Diversity and conservation status of avifauna in western Navsari, South Gujarat- 10 years of observations. *Species*.2022;23(72):369-389

Patel, H., Jjar, G., Bhatt, D., & Patel, K. (2021). *An Annotated Checklist of Avifauna from Hemchandracharya North Gujarat University Campus, Patan, Gujarat, India*. 3, 121–131.

Praveen J., Jayapal, R., & Pittie, A., 2016. A checklist of the birds of India. *Indian BIRDS* 11 (5&6): 113–172A.

Praveen J., Jayapal, R., & Pittie, A., 2020. Taxonomic updates to the checklists of birds of India, and the South Asian region—2020. *Indian BIRDS* 16 (1): 12–19.

Praveen J., & Jayapal, R., 2023. Taxonomic updates to the checklists of birds of India and the South Asian region—2023. *Indian BIRDS* 18 (5): 131–134.

Praveen, J., & Jayapal, R., 2024. Taxonomic updates to the checklists of birds of India and the South Asian region—2024. *Indian BIRDS* 19 (5): 155–158.

Rasmussen PC and Anderton JC. 2012. *Birds of South Asia: The Ripley Guide*, Vol. I and II, Published by Lynx Editions, Barcelona, Pp.414–415.

Rathod, U., & Bhaduri, R. (2022). Avifaunal diversity in Indian Institute of Technology Guwahati Campus, Assam, India. *Journal of Threatened Taxa*, 14.

Sanllorente, O., Ríos-Guisado, R., Izquierdo, L., Molina, J. L., Mourocq, E., & Ibáñez-Álamo, J. D. (2023). The importance of university campuses for the avian diversity of cities. *Urban Forestry & Urban Greening*, 86, 128038.

Shivhare, P., Mishra, N., & Singh, D. (2022). Studies on avian diversity of Shri Krishna University campus, Chhatarpur, Madhya Pradesh, India. *FLORA AND FAUNA*, 28(2).

Singh, K. (2018). Studies on avian diversity of banda university of agriculture and technology campus, Banda, Uttar Pradesh, India. *International International Journal of Avian & Wildlife Biology*, 3(2).

Singh, R., Tiwari, A. K., Kumar, S., & Singh, G. S. (2024). Avifaunal Diversity in Managed Urban Ecosystem: A Case Study of Banaras Hindu University, Varanasi. *Ecological Questions*, 35(3), 1–25.

Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., & Jetz, W. (2014). EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals: *Ecological Archives* E095-178. *Ecology*, 95(7), 2027–2027.