

# Effect of foliar application of water soluble fertilizer and humic acid on physico-chemical properties and fertility status of soil after harvest of rose

Commented [Mb1]: Rose cultivation-more relevant format

## Abstract

A field experiment on "Effect of foliar application of water soluble fertilizer and humic acid on yield and quality of rose" was carried out during Rabi season at PG Research unit, Horticulture Section, College of Agriculture, Nagpur during 2022-2023. The experiment was laid out in Factorial Randomized Block Design. The treatments comprised of four levels of WSF (19:19:19) viz., Control, 200 g, 300 g and 400 g and three levels of humic acid viz., Control, 500 ppm and 750 ppm giving twelve treatment combinations replicated thrice. In respect of available NPK after harvesting was found superior in treatment combination (F<sub>4</sub>H<sub>3</sub>) i.e. application of 400 g 19:19:19 with 750 ppm humic acid.

Commented [Mb2]: rose (*Rosa* spp.).

Commented [Mb3]: Define abbreviation on their first mention

Commented [Mb4]: Improve phrasing

**Keywords:** Rose, water soluble fertilizer, humic acid, soil fertility

## Introduction

Rose is native of USA and belongs to the family Rosaceae. It is a woody perennial flowering plant of genus *Rosa*. There are over three hundred species and tens thousands cultivars of rose. They form a group of plants that can be erect shrubs, climbing or trailing with stems that are often armed with sharp prickles. Rose flowers vary in size and shape and are usually large and showy, in colour's ranging from white through yellows and reds. Rose is commercially grown mainly for its cut flower production and grown mainly for Rose hips are high in vitamin C, are edible raw and occasionally made into jam, jelly, marmalade, rose syrup and soup, or are brewed for tea (Angier and Bradford, 1974). The antiseptic nature of rose petals make them a wonderful treatment for wounds, bruises, rashes and incisions, their anti-inflammatory properties make them a wonderful treatment for sore throats or ulcers. The extract of the rose petals is used as eye drops or eye wash in burning sensation of the eyes (Chahar, 2016). So, for its cultivation balanced fertilisation are required. Water

Commented [Mb5]: Fix grammatical issues

Commented [Mb6]: Typo

Commented [Mb7]: Improper linking. Avoid informal phrasing and improve grammar.

soluble fertilizer is easy to uptake by plant. As compare to soluble fertilizers, water soluble fertilizers have more nutrient use efficiency. Foliar application provides ample scope for utilization of nutrients more efficiently and for correcting the deficiencies rapidly. It also helps in the reduction in loss of nutrients. The great difficulty in supplying the macro nutrients through foliage is the non-availability of suitable water soluble fertilizers, which are a better source of nutrients for foliar application (Vibhute, 1998). Humic acid is a natural polymeric composition which is produced as a result of decaying organic matter in soil, peat and lignin and can be used in order to increase crop production (Sabzevari *et al.* 2008).

**Commented [Mb8]:** Both the sentences convey same meaning. Instead you may write as "water-soluble fertilizers are easily absorbed by plants and demonstrate higher nutrient use efficiency compared to conventional fertilizers."

## Materials and Methods

The experiment was conducted at PG Research Farm, Horticulture Section, College of Agriculture, Nagpur during the year 2022-23. The experiment was superimposed on ongoing experiment at Section of Horticulture, College of Agriculture, Nagpur. (21.15°N and 79.09°E, 310.50 m above MSL). Nagpur is characterized with hot and dry summer from March to May, warm and humid monsoon from June to October and fairly cold winter from November to February. The area shows wide fluctuation of temperature. The soil of experimental site was medium black in colour with good drainage. The soil properties before start of experiment is neutral in reaction (pH 7.22), medium in salt concentration (0.24 dSm<sup>-1</sup>), moderately low in organic carbon (4.70 g kg<sup>-1</sup>), low in calcium carbonate (2.76%), low in available N (230.2 kg ha<sup>-1</sup>) and available P (13.07 kg ha<sup>-1</sup>), moderately high in available K (248.3 kg ha<sup>-1</sup>) and low in available sulphur (11.68 kg ha<sup>-1</sup>).

**Commented [Mb9]:** Correct formatting of coordinates.

The experiment was laid out to study the effect of water soluble fertilizer and humic acid on fertility status of soil after harvest of rose. The research was carried out on variety First red.

Twelve treatment combinations with four levels of 19:19:19 viz. 0, 200, 300 and 400 g and three levels of humic acid 0, 500 and 750 ppm were tested in factorial randomized block design with three replications. The different combinations of water soluble fertilizer and humic acid as: T<sub>1</sub> - 0 g 19:19:19 + 0 ppm HA (F<sub>1</sub>H<sub>1</sub>), T<sub>2</sub> - 0 g 19:19:19 + 500 ppm HA (F<sub>1</sub>H<sub>2</sub>), T<sub>3</sub> - 0 g 19:19:19

**Commented [Mb10]:** List out the combinations in tabular form to improve structure of the manuscript

**Commented [Mb11]:** Define abbreviation on their first mention

+750 ppm HA (F<sub>1</sub>H<sub>3</sub>), T<sub>4</sub>- 200 g 19:19:19 + 0 ppm HA (F<sub>2</sub>H<sub>1</sub>), T<sub>5</sub>- 200 g 19:19:19 + 500 ppm HA (F<sub>2</sub>H<sub>2</sub>), T<sub>6</sub>- 200 g 19:19:19 + 750 ppm HA (F<sub>2</sub>H<sub>3</sub>), T<sub>7</sub>- 300 g 19:19:19 + 0 ppm HA (F<sub>3</sub>H<sub>1</sub>), T<sub>8</sub>- 300 g 19:19:19 + 500 ppm HA (F<sub>3</sub>H<sub>2</sub>), T<sub>9</sub>- 300 g 19:19:19 + 750 ppm HA (F<sub>3</sub>H<sub>3</sub>), T<sub>10</sub>- 400 g 19:19:19 + 0 ppm HA (F<sub>4</sub>H<sub>1</sub>), T<sub>11</sub>- 400 g 19:19:19 + 500 ppm HA (F<sub>4</sub>H<sub>2</sub>), T<sub>12</sub>- 400 g 19:19:19 + 750 ppm HA (F<sub>4</sub>H<sub>3</sub>).

The raised beds of 25 m length and 0.9 m breadth and 40 cm height were prepared. Distance between the two beds was 50 cm. Three to four months old budded plants were planted in above raised beds during the year 2018-19. The common recommended dose of 300:200:200 ppm NPK plant<sup>-1</sup> was applied at three split doses 1<sup>st</sup> split was given in the month of October, 2<sup>nd</sup> split was given in December and 3<sup>rd</sup> split was given in the month of January to all the plots (Singh and Peter, 2014). Package of practices including irrigation were adopted as per recommendation.

Sterilized growing media (soil + sand + rice husk + FYM + cocopeat) treated with *Trichoderma viridae* was used for planting rose plants. Spraying of water soluble fertilizer and humic acid will be done after pruning of rose plants. As regards the treatment of water-soluble fertilizer 19:19:19 was calculated accordingly 200 g, 300 g, 400 g of WSF has applied @ 5 g L<sup>-1</sup>, 7.5 g L<sup>-1</sup> and 10 g L<sup>-1</sup> in 40 splits in two days interval through foliar application. As regards humic acid application dose of humic acid was also calculated as per treatments and applied as foliar application in two equal splits at 15 days and 30 days after pruning.

Observations on various growth, flowering and yield parameters five plants will be recorded randomly in each treatment plot. A composite soil sample from the experimental site was collected before the crop was transplanted to know the nutrient status of the soil before the application of fertilizer.

After the harvesting of crop, the surface soil samples were collected (15-20 cm depth) from each treatment plot using screw auger. Soil samples collected were mixed and spread on small cotton cloth bags for air drying. Then the samples were ground and sieved through 2 mm sieve. The chemical properties viz., pH was determined in 1:2.5 soil water suspension with the help

Commented [Mb12]: Rose plants

Commented [Mb13]: Write in brackets

of glass electrode using pH meter (Jackson, 1973), Electrical conductivity (EC) of the soil was determined in 1:2.5 soil water suspension using conductivity bridge (Jackson, 1973), Organic carbon was estimated by Walkley and Black's (1934) Wet Oxidation method and Calcium carbonate was estimated by Rapid Titration method (Piper, 1966). Nitrogen content was determined as alkaline permagnate method described by Subbiah and Asija (1956), While phosphorus was estimated by using Olsen's method reagent (Olsen and Sommer 1982), potassium was extracted by 1N ammonium acetate of pH 7.0 and determined by using flame photometer as described by Jackson (1973) and sulphur determined by turbidity method given by Chesnin and Yien (1951). Data were statistically analysed in FRBD (Gomez and Gomez, 1984)

Commented [Mb14]: Spelling correction

Commented [Mb15]: Discuss the significance levels for the interaction

Commented [Mb16]: Define abbreviation on their first mention

## Results and discussion

### Effect of water soluble fertilizer and humic acid on physico-chemical properties of soil

In the present investigations, a residual effect of various nutritional treatments on the physico-chemical properties was also studied. Effect of water soluble fertilizer and humic acid individually and combine application have no significant effect on pH, EC, organic carbon and calcium carbonate of soil (Table 1 and 2).

Commented [Mb17]: remove

**Table 1: Effect of water soluble fertilizer (19:19:19) and humic acid on physico-chemical properties of soil**

Treatments	pH	EC (dSm <sup>-1</sup> )	Organic carbon (g kg <sup>-1</sup> )	Calcium Carbonate (%)
<b>Fertilizer (F)</b>				
F <sub>1</sub> Control	7.24	0.25	4.87	2.78
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	7.26	0.25	4.90	2.81
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	7.24	0.25	5.10	2.84
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	7.22	0.24	4.87	2.88
SE (m) ±	0.01	0.002	0.27	0.031
CD at 5%				
<b>Humic acid (H)</b>				
H <sub>1</sub> (control)	7.24	0.24	4.75	2.82

H <sub>2</sub> (500 ppm)	7.25	0.25	5.03	2.83
H <sub>3</sub> (750 ppm)	7.23	0.25	5.03	2.84
SE (m) ±	0.003	0.002	0.28	0.026
CD at 5%				
<b>Interaction (F x H)</b>				
SE (m) ±	0.02	0.007	0.38	0.043
CD at 5%				

**Table 2: Interaction effect of water soluble fertilizer (19:19:19) and humic acid on physico-chemical properties of soil**

Treatments Fertilizer	pH			
	Humic acid			Mean
H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)		
F <sub>1</sub> Control	7.22	7.23	7.28	7.24
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	7.29	7.24	7.24	7.26
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	7.24	7.27	7.21	7.24
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	7.20	7.25	7.21	7.22
Mean	7.24	7.25	7.23	
	<b>Factor A (F)</b>	<b>Factor B (H)</b>	<b>Interaction (F X H)</b>	
SE (m) ±	0.01	0.003	0.02	
CD at 5%				

Treatments	EC (dS m <sup>-1</sup> )
Fertilizer	Humic acid

	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
F <sub>1</sub> Control	0.25	0.24	0.25	0.25
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	0.25	0.25	0.25	0.25
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	0.25	0.27	0.25	0.25
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	0.23	0.25	0.25	0.24
Mean	0.24	0.25	0.25	
	Factor A (F)	Factor B (H)	Interaction (F X H)	
SE (m) ±	0.002	0.002	0.007	
CD at 5%				

Treatments	OC (g kg <sup>-1</sup> )			
Fertilizer	Humic acid			
	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
F <sub>1</sub> Control	4.60	4.90	5.10	4.87
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	4.80	4.90	5.00	4.90
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	5.00	5.30	5.00	5.10
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	4.60	5.00	5.00	4.87
Mean	4.75	5.03	5.03	
	Factor A (F)	Factor B (H)	Interaction (F X H)	
SE (m) ±	0.27	0.28	0.38	
CD at 5%				

Treatments	Calcium carbonate (%)
Fertilizer	Humic acid

	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
F <sub>1</sub> Control	2.78	2.78	2.80	2.78
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	2.80	2.82	2.83	2.81
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	2.84	2.85	2.85	2.84
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	2.87	2.89	2.90	2.88
Mean	2.82	2.83	2.84	
	Factor A (F)	Factor B (H)	Interaction (F X H)	
SE (m) ±	0.031	0.026	0.043	
CD at 5%				

#### Effect of water soluble fertilizer and humic acid on residual fertility status of soil

The data revealed that the application of individual levels of water soluble fertilizer and humic acid significantly increased the availability of nutrients in soil. Significantly, maximum available N in soil recorded in treatment F<sub>4</sub> with application of 400 g 19:19:19 (245.7 kg ha<sup>-1</sup>) which is found to be at par with 300 g 19:19:19 (F<sub>3</sub>). Similar finding was reported by Singh *et al.* (2015) in carnation. Whereas, maximum nitrogen (243.9 kg ha<sup>-1</sup>) was found in treatment 750 ppm humic acid (H<sub>3</sub>) which was at par with the treatment 500 ppm humic acid (H<sub>2</sub>). Similar finding was reported by Patil (2020). stated that, humic acid application (soil + foliar) was significantly increased available nitrogen in soil. Data shows that available nitrogen in soil was found highest (254.8 kg ha<sup>-1</sup>) in treatment 400 g 19:19:19 WSF and 500 ppm humic acid (F<sub>4</sub>H<sub>2</sub>) which was found to be at par with treatment F<sub>1</sub>H<sub>3</sub>, F<sub>3</sub>H<sub>1</sub> and F<sub>2</sub>H<sub>3</sub>. These results are also in the line with the findings of Chen and Aviad (1990), who reported significant differences in nitrogen uptake

**Commented [Mb18]:** consider providing more explanation on how the differences in values were statistically significant.

**Commented [Mb19]:** Format it properly

**Commented [Mb20]:** Write table number in which data is represented

**Table 3: Effect of water soluble fertilizer (19:19:19) and humic acid on N, P, K and S status of soil**

Treatments	Available N (kg ha <sup>-1</sup> )	Available P (kg ha <sup>-1</sup> )	Available K (kg ha <sup>-1</sup> )	Available S (kg ha <sup>-1</sup> )
<b>Fertilizer (F)</b>				
F <sub>1</sub> Control	234.4	13.41	258.2	12.23
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	239.1	13.62	258.7	12.41
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	243.9	13.74	261.3	12.48
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	245.7	13.82	262.0	12.57
SE (m) ±	1.83	0.05	0.64	0.04
CD at 5%	5.30	0.16	1.93	
<b>Humic acid (H)</b>				
H <sub>1</sub> (control)	235.6	13.47	256.5	12.32
H <sub>2</sub> (500 ppm)	242.7	13.72	261.8	12.44
H <sub>3</sub> (750 ppm)	243.9	13.74	261.7	12.51
SE (m) ±	1.33	0.04	0.86	0.03
CD at 5%	3.85	0.12	2.60	
<b>Interaction (F x H)</b>				
SE (m) ±	5.71	0.14	2.34	0.13
CD at 5%	11.56	0.41	6.78	

**Table 4: Interaction effect of water soluble fertilizer (19:19:19) and humic acid on N, P, K and S status of soil**

Treatments	Available nitrogen (kg ha <sup>-1</sup> )			
	Humic acid			
Fertilizer	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
<b>F<sub>1</sub> Control</b>	220.6	235.6	246.9	234.4
<b>F<sub>2</sub> @ 5 g L<sup>-1</sup> (200 g)</b>	234.3	238.5	244.4	239.1
<b>F<sub>3</sub> @ 7.5 g L<sup>-1</sup> (300 g)</b>	245.5	242.2	244.1	243.9
<b>F<sub>4</sub> @ 10 g L<sup>-1</sup> (400 g)</b>	242.1	254.8	240.4	245.7
<b>Mean</b>	235.6	242.7	243.9	
	<b>Factor A (F)</b>	<b>Factor B (H)</b>	<b>Interaction (F X H)</b>	
SE (m) ±	1.83	1.33	5.71	
CD at 5%	5.30	3.85	11.56	

Treatments	Available phosphorus (kg ha <sup>-1</sup> )			
Fertilizer	Humic acid			
	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
<b>F<sub>1</sub> Control</b>	13.11	13.24	13.90	13.41
<b>F<sub>2</sub> @ 5 g L<sup>-1</sup> (200 g)</b>	13.45	13.86	13.56	13.62
<b>F<sub>3</sub> @ 7.5 g L<sup>-1</sup> (300 g)</b>	13.50	13.89	13.83	13.74
<b>F<sub>4</sub> @ 10 g L<sup>-1</sup> (400 g)</b>	13.82	13.97	13.68	13.82
<b>Mean</b>	13.47	13.72	13.74	
	<b>Factor A (F)</b>	<b>Factor B (H)</b>	<b>Interaction (F X H)</b>	
SE (m) ±	0.05	0.04	0.14	
CD at 5%	0.16	0.12	0.41	

Treatments	Available potassium (kg ha <sup>-1</sup> )			
Fertilizer	Humic acid			
	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
<b>F<sub>1</sub> Control</b>	251.3	257.8	265.4	258.2
<b>F<sub>2</sub> @ 5 g L<sup>-1</sup> (200 g)</b>	254.2	262.2	259.6	258.7
<b>F<sub>3</sub> @ 7.5 g L<sup>-1</sup> (300 g)</b>	257.1	263.6	263.3	261.3
<b>F<sub>4</sub> @ 10 g L<sup>-1</sup> (400 g)</b>	263.6	263.7	258.7	262.0
<b>Mean</b>	256.5	261.8	261.7	
	<b>Factor A (F)</b>	<b>Factor B (H)</b>	<b>Interaction (F X H)</b>	
SE (m) ±	0.64	0.86	2.34	
CD at 5%	1.93	2.60	6.78	

Treatments	Available sulphur (kg ha <sup>-1</sup> )			
	Humic acid			
Fertilizer	H <sub>1</sub> (control)	H <sub>2</sub> (500 ppm)	H <sub>3</sub> (750 ppm)	Mean
F <sub>1</sub> Control	11.97	12.26	12.47	12.23
F <sub>2</sub> @ 5 g L <sup>-1</sup> (200 g)	12.34	12.50	12.40	12.41
F <sub>3</sub> @ 7.5 g L <sup>-1</sup> (300 g)	12.45	12.49	12.51	12.48
F <sub>4</sub> @ 10 g L <sup>-1</sup> (400 g)	12.53	12.49	12.68	12.57
Mean	12.32	12.44	12.51	
	Factor A (F)	Factor B (H)	Interaction (F X H)	
SE (m) ±	0.04	0.03	0.13	
CD at 5%				

The maximum available P in soil recorded in treatment F<sub>4</sub> with application of 400 g 19:19:19 (13.82 kg ha<sup>-1</sup>) which was found to be at par with (F<sub>3</sub>) 300 g 19:19:19 (13.74 kg ha<sup>-1</sup>). Similar finding was reported by Singh *et al.* (2015) in carnation. Significantly, maximum phosphorus (13.74 kg ha<sup>-1</sup>) was found in treatment 750 ppm humic acid (H<sub>3</sub>) which was at par with the treatment (H<sub>2</sub>). Similar finding was reported by Patil (2020) stated that, humic acid application (soil + foliar) was significantly increased availability of phosphorus in soil. Data shows that available phosphorus in soil was found highest (13.97 kg ha<sup>-1</sup>) in treatment 400 g 19:19:19 WSF and 500 ppm humic acid (F<sub>4</sub>H<sub>2</sub>) which was found to be at par with treatment F<sub>3</sub>H<sub>2</sub>, F<sub>2</sub>H<sub>2</sub>, F<sub>3</sub>H<sub>3</sub>, F<sub>4</sub>H<sub>1</sub> and F<sub>4</sub>H<sub>3</sub>.

The maximum available K in soil recorded in treatment (F<sub>4</sub>) with application of 400 g 19:19:19 (262.0 kg ha<sup>-1</sup>) which was found to be at par with (F<sub>3</sub>) 300 g 19:19:19 (261.32 kg ha<sup>-1</sup>). Similar results were also obtained by Singh *et al.* (2015) who also reported increase in available potassium in soil with

Commented [Mb21]: typo

Commented [Mb22]: Discuss the reason for the significant difference

Commented [Mb23]: Write table number in which data is represented

increasing doses of potassium application. Significantly, maximum potassium (261.8 kg ha<sup>-1</sup>) was found in treatment 500 ppm humic acid (H<sub>2</sub>) which was at par with the treatment (H<sub>3</sub>). Similar findings was reported by Patil (2020) reported that, humic acid application (soil + foliar) was significantly increased available potassium in soil. Data shows that available potassium in soil was found highest (263.7 kg ha<sup>-1</sup>) in treatment 400 g 19:19:19 WSF and 500 ppm humic acid (F<sub>4</sub>H<sub>2</sub>) which was found to be at par with treatment F<sub>3</sub>H<sub>2</sub>, F<sub>3</sub>H<sub>3</sub>, and F<sub>2</sub>H<sub>2</sub>. Homogenous results showing the positive co-relationship between the doses of HA and the potassium contents of the leaves were observed by Nikbakht et al. (2008).

The available sulphur was found non-significant with individual application of water soluble fertilizers and humic acid spray as well as in combination.

#### Conclusion:

The combine application of water soluble fertilizer @ 300 g and humic acid @ 500 ppm increased the availability of nutrients in soil and improves fertility status of soil.

#### References:

1. Angier and Bradford, 1974. Field guide to edible wild plants. Harrisburg, PA: Stackpole Books. pp.186. ISBN 0-8117-0616-8. OCLC 799792.
2. Chahar, 2016. Rose: Ornamental as well as medicinal plant. *J. Res. Agric. Animal Sci.* Vol. 4 Issue 1. pp 8-10. ISSN (online): 2321-9459. www.questjournals.org.
3. Chen, Y. and T. Aviad, 1990. Effects of Humic substances on plant growth. Humic substances in soil and crop science: selected readings. SSSA and ASA, Madison, Wisconsin, USA, pp. 161-186.
4. Chesnin, L. and C. H. Yien, 1951. Turbidimetric determination of available Sulphate. *Soil Sci. Soc. America*, 15: 149-151.
5. Gomez, K. A. and A. A. Gomez, 1984. *Statistical Procedure for Agricultural Research*, 2<sup>nd</sup> Ed. New York: John Wiley & Sons.
6. Jackson, M. L. 1973. *Soil Chemical Analysis*, Prentice Hall, India, Pvt. Ltd., New Delhi (India), PP. 498.
7. Nikbakht, A. M., M. Kafi, Y. P. Babalar, Xia, AnchengLuo and N. Etemadi, 2008. Effect of humic acid on plant growth nutrient uptake and postharvest life of gerbera. *J. of Plant Nutrition*, 31: 2155-2167.

**Commented [Mb24]:** Explain why significant difference in the treatments with proper explanation.

**Commented [Mb25]:** Add space after fullstop

**Commented [Mb26]:** Write table number in which data is represented

**Commented [Mb27]:** Write in italics

**Commented [Mb28]:** The conclusion should emphasize the practical implications for nutrient management, and how these findings could be applied in agricultural settings.

**Commented [Mb29]:** Irrelevant to the values mentioned in results and discussions. Check once.

**Commented [Mb30]:** Citation formatting should align with the journal's guidelines.

8. Olsen, S. R. and L. E. Sommer, 1982. Phosphorus methods of soil analysis; chemical and microbiological properties, Part 2. 2<sup>nd</sup> ed., Agron. 403-430.
9. Patil, H. 2020. Influence of humic acid on growth, yield and quality of chrysanthemum (*Dendranthema grandiflora* T.) cv. Marigold (Unpublished master's thesis). University of Agricultural Sciences, Bangalore.
10. Piper, C. S. 1966. Soil and Plant Analysis. Hans Publishers, Bombay pp. 368.
11. Sabzevari, S., H. R. Khazaie and M. Kafi, 2008. Study on the effects of humic acid on germination of four wheat cultivars (*Triticum aestivum* L.). *Iranian J. Field crops res.*, 8(3): 473-480.
12. Singh, D. K. and K. V. Peter, 2014. Protected cultivation of Hort. Crops 88 1<sup>st</sup> New Delhi: 334.
13. Singh, A., B. P. Sharma, B. S. Dilta, B. S. Nomita laishram, Y. C. Gupta and S. K. Bharadwaj, 2015. Effects of fertilization on quality flower production and foliar nutrient content of carnation (*Dianthus caryophyllus* L.) cv. Master. *Bangladesh J. Bot.* 44(1): 133-137.
14. Subbiah, B. V. and G. L. Asija, 1956. A rapid procedure for determination of available nitrogen in Soil *Current Sci.*, 25: 256-260.
15. Vibhute, C. P. 1998. A process for manufacturing complex solid and liquid completely water soluble fertilizers. *Fer. News*, 43(8): 63.
16. Walkley, A. J. and A. I. Black, 1934. Estimation of organic carbon by chromic acid Titration method. *Soil Sci.*, 25: 255-259.



