

Effect of recommended and graded doses of Biochar with and without ZMB biofertilizer and Zinc on the Growth attributes of Rice (*Oryza sativa* L.)

Abstract

A field experiment was conducted during the Kharif seasons of 2022 and 2023 at the Students' Instructional Farm, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj Ayodhya, to assess the impact of graded doses of biochar, varying fertility levels, and the addition of biofertilizers and zinc on rice cultivation under partially reclaimed sodic soils. Nine treatments were applied, including combinations of recommended doses of fertilizers (RDF), biochar (5 t ha⁻¹ and 2.5 t ha⁻¹), biofertilizer (ZMB), and zinc sulfate (ZnSO₄) in a randomized block design with three replications. The treatments were: T₁ (Control), T₂ (100% RDF), T₃ (50% RDF), T₄ (50% RDF + 5 t ha⁻¹ biochar), T₅ (50% RDF + 5 t ha⁻¹ biochar + ZMB), T₆ (50% RDF + 5 t ha⁻¹ biochar + ZMB + ZnSO₄), T₇ (100% RDF + 2.5 t ha⁻¹ biochar), T₈ (100% RDF + 2.5 t ha⁻¹ biochar + ZMB), and T₉ (100% RDF + 2.5 t ha⁻¹ biochar + ZMB + ZnSO₄). The rice variety NDR 2065 was used as the test crop. Growth parameters such as plant height, dry matter accumulation, root length, root volume, number of tillers, and panicle length were evaluated. The results showed that the treatment T₉ (100% RDF + 2.5 t ha⁻¹ biochar + ZMB + ZnSO₄) produced the highest growth and quality parameters, indicating a significant improvement in rice performance. The study suggests combining biochar, biofertilizers, and zinc under reduced fertilizer levels can enhance rice growth and productivity in sodic soils, offering a sustainable approach to soil fertility management.

Keywords – *Kharif, Plant Height, ZnSO₄, ZMB.*

Introduction

Rice (*Oryza sativa* L.) is the most important cereal crop grown under aquatic conditions and mostly under submergence or variable ponding conditions. It belongs to the family *Poaceae (Gramineae)*. It is the most important staple food of about more than 60% of the total world population. The

production of rice in the world is about 509.26 million metric tons with a productivity of 4.60 metric tons ha⁻¹, an area of 165.21 million ha (**Anonymous, 2021- 2022**). In India, rice is cultivated in 450.57 Lakh ha with an annual production of 122.27 million tons and average productivity of 2713 kg ha⁻¹ (**Anonymous, 2022**). Uttar Pradesh is the largest rice-growing state after West Bengal. In Uttar Pradesh, the area under rice cultivation is about 59.70 lakh ha. and the production was about 159.68 Lakh MT with a productivity of 26.75 q ha⁻¹. (**Anonymous, 2022**). Indo-Gangetic and other river-side states like West Bengal, Uttar Pradesh, Bihar, Punjab, Haryana, Odisha, Chhattisgarh, Andhra Pradesh, Telangana, Tamil Nadu, Kerala, Assam, and others are major rice-producing states in India. Punjab has experienced excellent prosperity in rice cultivation during the last 60 years since the Green Revolution began in India in the 1960s. Generally, Rice crop requires hot and humid weather, plentiful water supply, and abundant sunshine. Ideal temperature ranges from 20^o C to 40^o C and ideal rainfall ranges from 100 to 200. Traditionally.

Biochar (BC) is the carbon-rich product obtained when biomass, such as wood, manure, or leaves, is heated in a closed container with little or no available air. In more technical terms, BC is produced by the so-called thermal decomposition of organic material under a limited supply of oxygen (O₂) and at relatively low temperatures. The addition of BC to agricultural soils has been projected as a means to improve soil fertility and mitigate climate change. It is indicated that amending BC into soil improves the structure and properties of soil, such as the water-holding capacity, organic matter content, aeration condition, pH value, cationic exchange capacity (CEC), and the formation of aggregates of soil. The leaching losses of nitrogen and phosphorous in soil and the releases of greenhouse gases (N₂O and CH₄) from soil could be decreased in the presence of BC. In addition, BC has a porous structure, charged surface, and surface functional groups (such as carboxyl, hydroxyl, phenolic hydroxyl, and carbonyl groups). These properties are the important factors that influence the migration, transformation, and bioavailability of contaminants in soil. The application of Biochar improves soil fertility through two mechanisms: adding nutrients to the soil (such as K, to a limited extent P, and many micronutrients) or retaining nutrients from other sources, including nutrients from the soil itself. Furthermore, the application of Biochar to soils has been identified as a low-cost technology that can stabilize organic carbon, reduce greenhouse gas emissions, improve soil physical and chemical properties, and boost crop yield productivity and farm incomes. Biochar application can lead to a reduction in inorganic fertilizer use by farmers. The sole Biochar application, in most cases, did not provide any substantial amount of nutrients.

Material and Methods

The experiments were conducted at the Student's Instructional Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya situated on Ayodhya-Raebareli Road about 42 km from Ayodhya. The experimental site falls under the subtropical climatic zone of Indo-Gangetic plains located at 26.54° N latitude, 81.82° E longitudes, and an altitude of 113 meters above mean sea level. The district Ayodhya falls under a sub-humid climate, receiving a mean annual rainfall of about 1200 mm. About 85% of the total rainfall is received from mid-June to the end of September. However, occasional showers are also common during the winter. The winter months are cold, and an occasional frost occurs during this period. The NDR 2065 variety of rice were taken for the study. The treatments used in the study were: T₁ (Control), T₂ (100% RDF), T₃ (50% RDF), T₄ (50% RDF + 5 t Biochar), T₅ (50% RDF + 5 t Biochar + ZMB), T₆ (50% RDF + 5 t Biochar + ZMB + Zn), T₇ (100% RDF + 2.5 t Biochar), T₈ (100% RDF + 2.5 t Biochar + ZMB), and T₉ (100% RDF + 2.5 t Biochar + ZMB + Zn). Observations taken for study are plant height, dry matter accumulation, root length, root volume, no. of tillers, length of panicle, were recorded during the investigation. The experiment was done using a randomized block design and data was analyzed using OPState Software.

Result and Discussion

The data regarding plant height is presented in Table 1 and Figure 1 at different growth stages, 30 DAT, 60 DAT, 90 DAT, and at harvest stage. Treatment incorporating biochar (T₄-T₉) consistently exhibited higher plant heights compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had a plant height of 30.9 cm and 32.2 cm respectively, while treatment T₂ (100% RDF 100:60:40, N P₂O₅, K₂O) had a plant height of 34.5 cm and 35.9 cm for both years and T₃ (50 % RDF) had a plant height 33.5 cm and 34.5 cm for both the years respectively. However, the maximum plant height was recorded with the application of T₉ (38.9 cm and 40.1 cm) for both years. This trend was evident throughout all growth stages in both years of experimentation, highlighting the positive impact of biochar on plant height. In T₇ (100 % RDF + 2.5 t/ha-1 Biochar) and T₉ (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) at 60 DAT in 2022 had a plant height of 57.5 cm and 61.8 cm respectively, and in 2023 T₇ and T₉ had a plant height of 59.5 and 63.8 cm respectively. while T₄ (50 % RDF + 5 t/ha-1 Biochar) and T₆ (50 % RDF + 5 t/ha-1 Biochar + ZMB + Zn) in 2022 had a plant height of 55.9 cm and 56.1 cm. while in 2023 T₄ and T₆ had a plant height of 57.8 cm and 59.0 cm. Including biofertilizers in treatments (T₅, T₆, T₈, and T₉) further augmented plant height. These treatments consistently exhibited higher plant heights compared to treatments without biofertilizer at all growth stages and in both years. For instance, at 90 DAT in 2022, treatment T₉ (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) had a plant height of 110.5 cm,


while T4 and T7 without biofertilizer had plant heights ranging from 101.1 cm to 104.1 cm and 105.9 cm to 107.1 cm respectively for both the years. The integration of biofertilizers further enhances the plant height response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights for optimizing biochar and fertility management practices to enhance plant height and potentially crop productivity in sustainable agriculture. This result corroborated with findings of **Khan *et al.* (2021)**, **Gul *et al.* (2015)**, **Khedwal *et al.* (2018)**, **Sharanabasappa and Basavanneppa (2019)**, **Ponmozhi *et al.* (2019)**, **Singh *et al.* (2021)** and **Akhtar *et al.* (2021)**.

UNDER PEER REVIEW

Table 1: Effect of Biochar and fertility levels on plant height (cm) at different intervals of rice

Symbol	Treatments	Plant height (cm)							
		30 DAT		60 DAT		90 DAT		At harvest	
		2022	2023	2022	2023	2022	2023	2022	2023
T₁	Control	30.9	32.2	40.6	42.9	70.8	72.9	74.2	76.4
T₂	100 %RDF	34.5	35.9	54.9	57.8	95.5	97.4	98.4	100.5
T₃	50%RDF	33.5	34.3	51.0	52.9	94.5	95.8	97.8	99.9
T₄	50%RDF+5t/ha ⁻¹ Biochar	34.8	36.0	55.9	57.8	101.1	104.1	104.1	105.3
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	35.2	36.7	55.9	57.8	101.1	103.1	104.8	106.0
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	35.5	37.0	56.1	59.0	102.5	105.6	105.5	106.7
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	36.1	37.7	57.5	59.5	105.9	107.1	108.6	110.9
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	38.2	39.8	60.9	62.1	109.9	111.2	113.3	115.7
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	38.9	40.1	61.8	63.8	110.5	112.7	114.6	116.8
	SEm±	1.14	1.18	1.78	1.84	3.20	3.26	3.31	3.37
	CD (P= 0.05)	3.33	3.47	5.20	5.39	9.37	9.54	9.68	9.85

Fig 1 : Effect of Biochar and fertility levels on plant height (cm) at different intervals of rice



Control
 100 %RDF
 50%RDF
 50%RDF+5t/ha-1Biochar
 50 %RDF + 5 t/ha-1Biochar+ZMB
 50 %RDF + 5 t/ha-1Biochar+ZMB + Zn
 100 %RDF+2.5t/ha-1 Biochar
 100 %RDF +2.5t/ha-1Biochar +ZMB
 100 %RDF +2.5t/ha-1Biochar +ZMB + Zn

UNDEFIN

The data regarding the no. of tillers is presented in Table 2 and Figure.2 at different growth stages, 30 DAT, 60 DAT, 90 DAT, and at harvest stage. The data demonstrate the positive influence of biochar and fertility levels on the tillers (m^2) throughout the crop's growth stages in both years. Treatment incorporating biochar (T₄-T₉) consistently exhibited higher number of tillers compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had the no. of tillers (m^2) of 115 (m^2) and 117 (m^2) respectively, while treatment T₂ (100% RDF 100:60:40, N P₂O₅, K₂O) had the no. of tillers (m^2) of 136 (m^2) and 138 (m^2) for both years and T₃ (50 % RDF) had the no. of tillers (m^2) 132 (m^2) and 135 (m^2) for both the years respectively. However, the maximum no of tillers (m^2) was recorded with the application of T₉ with values 152 (m^2) and 155 (m^2) respectively for both the years. This trend was evident throughout all growth stages in both year of experimentation, highlighting the positive impact of biochar on no of tillers (m^2). In T₇ (100 % RDF + 2.5 t ha⁻¹ Biochar) and T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) at 60 DAT in 2022 had the no. of tillers (m^2) with values of 295 (m^2) and 328 (m^2) respectively, and in 2023 T₇ and T₉ had the no. of tillers (m^2) with values of 304 (m^2) and 338 (m^2) respectively. While T₄ (50 % RDF + 5 t ha⁻¹ Biochar) and T₆ (50 % RDF + 5 t ha⁻¹ Biochar + ZMB + Zn) in 2022 had the no. of tillers (m^2) with values of 265 (m^2) and 277 (m^2). But in 2023 T₄ and T₆ had the no. of tillers (m^2) with values of 274 (m^2) and 288 (m^2). The inclusion of biofertilizer in treatments (T₅, T₆, T₈, and T₉) further augmented the no. of tillers. These treatments consistently exhibited higher the no of tillers compared to treatments without biofertilizer at all growth stages and in both years. For instance, at 90 DAS in 2022, treatment T₉ (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) had the no. of tillers with the values of 308 (m^2), while treatments without biofertilizer had number of tillers ranging from 246 (m^2) to 274 (m^2) respectively for both years. The same trends were also found at harvest stage of the crop during both the year of experimentation. Overall, the results demonstrate that the application of biochar and the inclusion of biofertilizer positively influence the no. of tillers at different growth stages of the crop. The integration of biofertilizer further enhances the no. of tillers response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights for optimizing biochar and fertility management practices to enhance the no. of tillers and potentially crop productivity in sustainable agriculture. This result corroborated with finding of **Gowthami et al. (2022)**, **Oo (2022)**, **Oladele et al. (2019)**, **Khan et al. (2021)**.

Table 2: Effect of Biochar and fertility levels on no. of tillers (m²) at different intervals of rice

Symbol	Treatments	Number of tillers (m ²)							
		30 DAT		60 DAT		90 DAT		At harvest	
		2022	2023	2022	2023	2022	2023	2022	2023
T ₁	Control	115	117	183	192	170	177	167	173
T ₂	100 %RDF	136	138	260	268	242	247	237	242
T ₃	50%RDF	132	135	239	248	222	228	218	224
T ₄	50%RDF+5t/ha ⁻¹ Biochar	140	143	265	274	246	252	242	248
T ₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	141	144	269	278	250	256	245	251
T ₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	144	146	277	288	258	265	252	260
T ₇	100 %RDF+2.5t/ha ⁻¹ Biochar	145	148	295	304	274	280	271	275
T ₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	149	153	325	334	302	307	296	302
T ₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	152	155	328	338	308	312	300	310
	SEm±	4.50	4.59	8.72	9.01	8.10	8.29	7.95	8.14
	CD (P= 0.05)	13.16	13.44	25.51	26.35	23.71	24.25	23.27	23.80




Fig 2 : Effect of Biochar and fertility levels on no. of tillers (m²) at different intervals of rice


The data regarding plant height presented in Table 3 and Figure.3 at different growth stages, 30 DAT, 60 DAT, 90 DAT, and at harvest stage. The data demonstrate the positive influence of biochar and fertility levels on plant height throughout the crop's growth stages in both years. The application of biochar and biofertilizer an increased in plant height was observed. Treatment incorporating biochar (T₄-T₉) consistently exhibited higher plant heights compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had a root length of 7.0 cm and 7.3 cm respectively, while treatment T₂ (100% RDF100:60:40, N P₂O₅, K₂O) had a root length of 10.5 cm and 10.7 cm for both years and T₃ (50 % RDF) had a root length 10.1 cm and 10.5 cm for both the years respectively. However, the maximum root length was recorded with the application of T₉ (12.4 cm and 12.9 cm) for both the years. This trend was evident throughout all growth stages in both year of experimentation, highlighting the positive impact of biochar on root length. In T₇ (100 % RDF + 2.5 t ha⁻¹ Biochar) and T₉ (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) at 60 DAT in 2022 had a root length of 33.7 cm and 35.7 cm respectively, and in 2023 T₇ and T₉ had a root length of 34.8 and 36.8 respectively. while T₄ (50 % RDF + 5 t ha⁻¹ Biochar) and T₆ (50 % RDF + 5 t/ha-1 Biochar + ZMB + Zn) in 2022 had a root length of 28.7 cm and 32.8 cm. while in 2023 T₄ and T₆ had a root length of 30.4 cm and 34.2 cm. The inclusion of Biofertilizer in treatments (T₅, T₆, T₈, and T₉) further augmented root length. These treatments consistently exhibited higher root length compared to treatments without biofertilizer at all growth stages and in both years. For instance, at 90 DAS in 2022 and 2023, treatment T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) had a root length of 33.7 cm and 34.5 respectively, while treatments without biofertilizer had root length ranging from 27.9 cm to 28.1 cm and 31.9 cm to 32.7 cm respectively for both the years. The same trends were also found at harvest stage of the crop during both the year of experimentation. Overall, the results demonstrate that the application of biochar and the inclusion of biofertilizer positively influence root length at different growth stages of the crop. Moreover, the dose-response relationship indicates that higher doses of biochar lead to greater root length. The integration of biofertilizer further enhances the root length response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights for optimizing biochar and fertility management practices to enhance root length and potentially crop productivity in sustainable agriculture. This result corroborated with finding of **Thavanesanand Seran (2018)**, **Kamara et al. (2015)**, **Sial et al. (2019)**, **Oladele et al. (2019)**.

Table 3 Effect of Biochar and fertility levels on root length (cm) at different intervals of rice

Symbol	Treatments	Root length (cm)							
		30 DAT		60 DAT		90 DAT		At harvest	
		2022	2023	2022	2023	2022	2023	2022	2023
T₁	Control	7.0	7.3	19.5	20.9	16.7	18.4	14.3	15.7
T₂	100 %RDF	10.5	10.7	27.5	28.9	26.8	27.4	17.9	18.9
T₃	50%RDF	10.1	10.5	25.8	26.8	24.7	25.3	17.2	18.2
T₄	50%RDF+5t/ha ⁻¹ Biochar	10.7	10.9	28.7	30.4	27.9	28.1	18.4	19.8
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	11.0	11.3	30.6	32.7	28.3	30.8	20.3	21.3
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	11.2	11.6	32.8	34.2	30.7	31.7	21.9	22.4
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	11.5	12.2	33.7	34.8	31.9	32.7	23.6	24.5
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	12.0	12.5	34.8	35.6	32.8	33.5	24.6	25.4
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	12.4	12.9	35.7	36.8	33.7	34.5	25.4	26.9
	SEm±	0.35	0.36	0.96	1.01	0.90	0.95	0.65	0.69
	CD (P= 0.05)	1.01	1.05	2.82	2.96	2.64	2.77	1.91	2.01

Fig 3 : Effect of Biochar and fertility levels on root length (cm) at different intervals of rice

UNDER PEER REVIEW



- Control
- 100 %RDF
- 50%RDF
- 50%RDF+5t/ha-1Biochar
- 50 %RDF + 5 t/ha-1Biochar+ZMB
- 50 %RDF + 5 t/ha-1Biochar+ZMB + Zn
- 100 %RDF+2.5t/ha-1 Biochar
- 100 %RDF +2.5t/ha-1Biochar +ZMB
- 100 %RDF +2.5t/ha-1Biochar +ZMB + Zn


UNDEF

The data regarding root volume presented in Table.4 and Figure .4 at different growth stages, 30 DAT, 60 DAT, 90 DAT, and at harvest stage. The data demonstrate the positive influence of biochar and fertility levels on root volume throughout the crop's growth stages in both years. The application of biochar and biofertilizer an increased in root volume was observed. Treatment incorporating biochar (T₄-T₉) consistently exhibited higher root volume compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had a root volume of 18.8 cm and 20.8 cm respectively, while T₃ (50 % RDF) had a root volume of 21.0 cm and 23.4 cm for both the years respectively. However, the maximum root volume was recorded with the application of T₉ (23.6 cm in 2022 and T₈ (25.7 cm) in 2023. This trend was evident throughout all growth stages in both year of experimentation, highlighting the positive impact of biochar on root volume. In T₇ (100 % RDF + 2.5 t/ha-1 Biochar) and T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) at 60 DAT in 2022 had a root volume of 39.0 cm and 40.8 cm respectively, but in 2023 T₇ and T₉ had a root volume of 41.3 cm and 42.9 cm respectively. while T₄ (50 % RDF + 5 t ha⁻¹ Biochar) and T₆ (50 % RDF + 5 t ha⁻¹ Biochar + ZMB + Zn) in 2022 had a root volume of 36.8 cm and 38.5 cm. while in 2023 T₄ and T₆ had a root volume of 38.4 cm and 40.2 cm. The inclusion of biofertilizer in treatments (T₅, T₆, T₈, and T₉) further augmented root volume. These treatments consistently exhibited higher root volume compared to treatments without biofertilizer at all growth stages and in both years. For instance, at 90 DAT in 2022 and 2023, treatment T₉ (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) had a root volume of 50.7 cm and 52.4 respectively, while treatments without biofertilizer had root volume ranging from 44.8 cm to 47.2 cm and 46.3 cm to 49.7 cm respectively for both the years. The same trends were also found at the harvest stage of the crop during both years of experimentation. Overall, the results demonstrate that the application of biochar and the inclusion of biofertilizer positively influence root volume at different growth stages of the crop. Moreover, the dose-response relationship indicates that higher doses of biochar lead to greater root volume. The integration of biofertilizer further enhances the root volume response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights for optimizing biochar and fertility management practices to enhance root volume and potentially crop productivity in sustainable agriculture. This result corroborated with finding of **Thavanesan and Seran (2018)**, **Kamara et al, (2015)**, **Sial et al, (2019)**, **Oladele et al, (2019)**, **Liu et al (2021)**, **Wang et al (2018)**.

Table 4 Effect of Biochar and fertility levels on root volume at different intervals of rice

Symbol	Treatments	Root Volume							
		30 DAT		60 DAT		90 DAT		At harvest	
		2022	2023	2022	2023	2022	2023	2022	2023
T₁	Control	18.8	20.8	32.7	34.6	40.6	42.3	42.8	43.9
T₂	100 %RDF	21.4	22.9	36.0	38.2	44.0	46.8	45.2	46.2
T₃	50%RDF	21.0	23.4	35.4	37.0	43.6	45.7	44.7	45.9
T₄	50%RDF+5t/ha ⁻¹ Biochar	21.9	23.9	36.8	38.4	44.8	46.3	45.8	46.7
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	22.1	24.8	37.8	39.6	45.3	47.2	46.3	47.3
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	22.5	24.6	38.5	40.2	46.0	48.1	47.5	48.5
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	22.7	24.3	39.0	41.3	47.2	49.7	49.2	50.4
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	23.2	25.7	39.7	42.1	48.6	50.8	50.3	51.8
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	23.6	25.1	40.8	42.9	50.7	52.4	52.8	53.9
	SEm±	0.74	0.81	1.26	1.33	1.54	1.61	1.60	1.63
	CD (P= 0.05)	2.14	2.34	3.67	3.86	4.49	4.68	4.64	4.75

Fig 4 : Effect of Biochar and fertility levels on root volume at different intervals of rice



- Control
- 100 %RDF
- 50%RDF
- 50%RDF+5t/ha-1Biochar
- 50 %RDF + 5 t/ha-1Biochar+ZMB
- 50 %RDF + 5 t/ha-1Biochar+ZMB + Zn
- 100 %RDF+2.5t/ha-1 Biochar
- 100 %RDF +2.5t/ha-1Biochar +ZMB
- 100 %RDF +2.5t/ha-1Biochar +ZMB + Zn


UNDER

The data regarding Leaf area index presented in Table 5 and Figure 5 at different growth stages, 30 DAT, 60 DAT and 90 DAT. The data demonstrate the positive influence of biochar and fertility levels on Leaf area index throughout the crop's growth stages in both years. The application of biochar and biofertilizer an increased in Leaf area index was observed. Treatment incorporating biochar (T₄-T₉) consistently exhibited higher Leaf area index compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had a Leaf area index of 2.12 and 2.31 respectively, while treatment T₂ (100% RDF 100:60:40, N P₂O₅, K₂O) had a Leaf area index of 2.39 and 2.61 for both years and T₃ (50 % RDF) had a Leaf area index 2.37 and 2.58 for both the years respectively. However, the maximum Leaf area index was recorded with the application of T₉ (2.60 and 2.84) for both the years respectively. This trend was evident throughout all growth stages in both year of experimentation, highlighting the positive impact of biochar on Leaf area index. In T₇ (100 % RDF + 2.5 t ha⁻¹ Biochar) and T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) at 60 DAT in 2022 had a Leaf area index of 6.37 and 7.02 respectively, but in 2023 T₇ and T₉ had a Leaf area index of 6.81 and 7.52 respectively. While T₄ (50 % RDF + 5 t ha⁻¹ Biochar) and T₆ (50 % RDF + 5 t ha⁻¹ Biochar + ZMB + Zn) in 2022 had a Leaf area index of 5.96 and 6.32. But in 2023 T₄ and T₆ had a Leaf area index of 6.38 and 6.76. The inclusion of biofertilizer in treatments (T₅, T₆, T₈, and T₉) further augmented Leaf area index. These treatments consistently exhibited higher Leaf area index compared to treatments without biofertilizer at all growth stages and in both years. For instance, at 90 DAT in 2022 and 2023, treatment T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) had a Leaf area index of 6.94 and 7.33 respectively, while treatments without biofertilizer (T₄ to T₇) had Leaf area index ranging from (5.85 to 6.25) and (6.20 to 6.63) respectively for both the years. Overall, the results demonstrate that the application of biochar and the inclusion of biofertilizer positively influence Leaf area index at different growth stages of the crop. Moreover, the dose-response relationship indicates that higher doses of biochar lead to greater Leaf area index. The integration of biofertilizer further enhances the Leaf area index response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights for optimizing biochar and fertility management practices to enhance Leaf area index and potentially crop productivity in sustainable agriculture. This result corroborated with finding of **Khan *et al.* (2021)**, **Gul *et al.* (2015)**, **Khedwal *et al.* (2018)** **Sharanabasappa and Basavanneppa (2019)** and **Singh *et al.* (2021)**.

Table 5 Effect of Biochar and fertility levels on leaf area index at different intervals of rice

Symbol	Treatments	Leaf area index (LAI)					
		30 DAT		60 DAT		90 DAT	
		2022	2023	2022	2023	2022	2023
T₁	Control	2.12	2.31	4.46	4.77	4.40	4.66
T₂	100 %RDF	2.39	2.61	5.37	5.75	5.24	5.55
T₃	50%RDF	2.37	2.58	5.33	5.70	5.11	5.42
T₄	50%RDF+5t/ha ⁻¹ Biochar	2.42	2.64	5.96	6.38	5.85	6.20
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	2.44	2.66	6.25	6.69	6.19	6.56
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	2.46	2.68	6.32	6.76	6.23	6.60
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	2.48	2.70	6.37	6.81	6.25	6.63
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	2.58	2.81	6.98	7.47	6.88	7.29
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	2.60	2.84	7.02	7.52	6.94	7.33
	SEm±	0.08	0.09	0.19	0.21	0.19	0.20
	CD (P= 0.05)	0.23	0.25	0.57	0.61	0.56	0.59

Fig 5 : Effect of Biochar and fertility levels on leaf area index at different intervals of rice




The data regarding Leaf area index presented in Table 6 and Figure.6 at different growthstages, 30 DAT, 60 DAT, 90 DAT and harvest stage. The application of biochar and biofertilizer an increased in Leaf area index was observed. Treatment incorporating biochar (T₄-T₉) consistently exhibited higherLeaf area index compared to the control treatment (T₁). In control at 30 DAT in 2022 and 2023 had the dry matter accumulation of 125.3 (g m⁻²) and 129.1 (g m⁻²), while treatment T₂ (100% RDF100:60:40, N P₂O₅, K₂O) had the dry matteraccumulation of 140.5 (g m⁻²) and 144.7(g m⁻²). However, the maximum dry matteraccumulation was recorded with the application of T₉. This trend was evident throughout all growth stages in both year of experimentation, highlighting the positive impact of biochar on dry matter accumulation. In T₇ (100 % RDF + 2.5 t ha⁻¹ Biochar) and T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) at 60 DAT in 2022 had the dry matter accumulation of 531.9 (g m⁻²) and 570.4 (g m⁻²)respectively, But in 2023 T₇ and T₉ had the dry matter accumulation of 542.5(g m⁻²) and 492.7(g m⁻²) respectively, While T₄ (50 % RDF + 5 t ha⁻¹ Biochar) and T₆ (50 % RDF + 5 t ha⁻¹ Biochar + ZMB + Zn) in 2022 had the dry matter accumulation of 522.1(g m⁻²) and 528.8 (g m⁻²) respectively. But in 2023 T₄ and T₆ had the dry matter accumulation of 532.5(g m⁻²) and 539.4 (g m⁻²) respectively. The inclusion of biofertilizer in treatments (T₅, T₆, T₈, and T₉) further augmented dry matter accumulation. These treatments consistently exhibited higher dry matter accumulation compared to treatments without biofertilizer at allgrowth stages and in both years. For instance, at 90 DAT in 2022 and 2023, treatment T₉ (100 % RDF + 2.5 t ha⁻¹ Biochar + ZMB + Zn) had a dry matter accumulation of 772.4 (g m⁻²) and 772.4 (g m⁻²) respectively, while treatments without biofertilizer (T₄ to T₇) haddry matter accumulation ranging from 698.5 (g m⁻²) to 711.6 (g m⁻²) and 712.5(g m⁻²) to 725.8 (g m⁻²) respectively for both the years. The same trends were also found at the harvest stage of the crop during both years of experimentation. Overall, the results demonstrate that the application of biochar and the inclusion of biofertilizer positively influence dry matter accumulation at different growth stages of the crop. The integration of biofertilizer further enhances the dry matter accumulation response, showcasing the potential of combined nutrient management strategies for improved crop growth. Overall, these findings contribute valuable insights foroptimizing biochar and fertility management practices to enhance dry matter accumulation and potentially crop productivity in sustainable agriculture. These results corroborated withfinding of **Cao *et al.* (2024)**, **Islam *et al.* (2018)**, **Khan *et al.* (2022)**, **Gul *et al.* (2015)**, **Roa *et al.* (2016)**, **Koca *et al.* (2016)**, **Khedwal *et al.* (2018)**, **Ponmozhi *et al.* (2019)**, **Kumar *et al.* (2020)** and **Singh *et al.* (2021)**.

Table.6 Effect of Biochar and fertility levels on dry matter accumulation (g m⁻²) at different intervals of rice

Symbol	Treatments	Dry matter accumulation (g m ⁻²)							
		30 DAT		60 DAT		90 DAT		At harvest	
		2022	2023	2022	2023	2022	2023	2022	2023
T₁	Control	125.3	129.1	328.7	335.3	468.6	478.0	690.2	693.7
T₂	100 %RDF	140.5	144.7	453.4	462.5	642.0	654.8	936.8	942.3
T₃	50%RDF	137.1	141.2	426.5	435.0	632.6	645.3	925.3	930.5
T₄	50%RDF+5t/ha ⁻¹ Biochar	152.1	156.7	522.1	532.5	698.5	712.5	1017.7	1025.2
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	152.5	157.1	524.1	534.6	705.7	719.8	1021.0	1028.5
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	154.1	158.7	528.8	539.4	708.6	722.8	1025.2	1032.8
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	156.4	161.1	531.9	542.5	711.6	725.8	1032.3	1040.1
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	164.7	169.6	568.7	580.1	762.4	777.6	1111.3	1121.0
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	167.3	172.8	570.4	592.7	772.4	785.4	1118.5	1128.4
	SEm±	4.84	4.99	16.08	16.40	21.96	22.40	31.93	32.16
	CD (P= 0.05)	14.16	14.59	47.03	47.97	64.23	65.52	93.41	94.08

Fig 6 : Effect of Biochar and fertility levels on dry matter accumulation (g m⁻²) at different intervals of rice

UNDER PEER REVIEW



Control
 100 %RDF
 50%RDF
 50%RDF+5t/ha-1Biochar
 50 %RDF + 5 t/ha-1Biochar+ZMB
 50 %RDF + 5 t/ha-1Biochar+ZMB + Zn
 100 %RDF+2.5t/ha-1 Biochar
 100 %RDF +2.5t/ha-1Biochar +ZMB
 100 %RDF +2.5t/ha-1Biochar +ZMB + Zn

UNDER

The table 7 presents the crop growth rate (CGR) in grams per square meter per day ($\text{gm}^{-2} \text{ day}^{-1}$) at different growth stages, 30-60 DAT, 60-90 DAT, and 90 DAT to at harvest, for the years 2022 and 2023. The study includes nine treatments, each representing different combinations of biochar, fertility levels, and biofertilizer, along with the corresponding standard error of the mean ($\text{SEm} \pm$) and critical difference at the 5% level of significance (CD). The data demonstrate the impact of biochar and fertility levels on the crop growth rate throughout the different growth stages in both years. In the control treatment (T_1), without any biochar or biofertilizer, the crop growth rate ranged from 6.78 to 7.39 $\text{gm}^{-2} \text{ day}^{-1}$ in 2022 and from 6.87 to 7.19 $\text{gm}^{-2} \text{ day}^{-1}$ in 2023. However, treatments with biochar and/or biofertilizer consistently exhibited higher crop growth rates compared to the control. Treatments with biochar (T_4 - T_9) consistently displayed higher crop growth rates compared to the control treatment (T_1). For instance, at 30-60 DAT in 2022, the control treatment (T_1) had a crop growth rate of 6.78 $\text{gm}^{-2} \text{ day}^{-1}$, and treatment T_2 (100% RDF) had a crop growth rate of 10.43 $\text{gm}^{-2} \text{ day}^{-1}$. While treatment T_9 (100% RDF 100:60:40, N P_2O_5 , K_2O with 2.5 t ha^{-1} biochar and ZMB biofertilizer +Zn) had a crop growth rate of 13.44 $\text{gm}^{-2} \text{ day}^{-1}$. This trend was observed across all growth stages and in both years, indicating the positive impact of biochar on the crop growth rate. T_7 (100 % RDF + 2.5 t ha^{-1} Biochar) and T_9 (100 % RDF + 2.5 t/ha-1 Biochar + ZMB + Zn) at 60-90 DAT in 2022 had a crop growth rate of 5.99(g m^{-2}) and 6.73(g m^{-2}) respectively, But in 2023 T_7 and T_9 had a crop growth rate of 6.11 (g m^{-2}) and 6.42 (g m^{-2}) respectively, While T_4 (50 % RDF + 5 t ha^{-1} Biochar) and T_6 (50 % RDF + 5 t ha^{-1} Biochar + ZMB + Zn) in 2022 had a crop growth rate of 5.88 (g m^{-2}) and 5.99 (g m^{-2}) respectively. But in 2023 T_4 and T_6 had a crop growth rate of 6.00(g m^{-2}) and 6.11 (g m^{-2}) respectively, the same trends were also found at 90- at the harvest stage of the crop during both the year of experimentation. The inclusion of biofertilizers in some treatments (T_5 , T_6 , T_8 , and T_9) further enhanced the crop growth rate. These treatments consistently exhibited higher crop growth rates compared to treatments without biofertilizer at all growth stages and in both years. The results indicate the significant influence of biochar and the integration of biofertilizers on the crop growth rate at different growth stages. The consistent increase in crop growth rate with the application of biochar suggests its positive effect on enhancing plant growth and productivity. Moreover, the observed dose-response relationship highlights the importance of optimizing biochar doses to achieve maximum crop growth rates. Additionally, the inclusion of biofertilizers in some treatments further enhances

the crop growth rate, indicating its role in promoting nutrient uptake and overall plant growth. The combination of biochar and biofertilizer shows promising synergistic effects on the crop's growth, which can lead to improved crop performance and yield potential.


Overall, the findings from this study provide valuable insights into the benefits of using biochar and biofertilizer as integrated soil management practices to enhance crop growth rates and potentially optimize agricultural productivity. Further investigation is warranted to better understand the underlying mechanisms driving these responses and to explore the long-term implications of these practices on overall crop performance and sustainable agriculture. Finding of the investigation may be confirmed by the earlier research workers **Prasad *et al.* (2017) and Khedwal *et al.* (2018).**

UNDER PEER REVIEW

Table 7 Effect of Biochar and fertility levels on crop growth rate (g/m²/day) at different intervals of rice

Symbol	Treatments	Crop growth rate (g/m ² /day)					
		30-60 DAT		60-90 DAT		90- at harvest	
		2022	2023	2022	2023	2022	2023
T₁	Control	6.78	6.87	4.66	4.76	7.39	7.19
T₂	100 %RDF	10.43	10.59	6.29	6.41	9.83	9.58
T₃	50%RDF	9.65	9.79	6.87	7.01	9.76	9.51
T₄	50%RDF+5t/ha ⁻¹ Biochar	12.33	12.53	5.88	6.00	10.64	10.42
T₅	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB	12.39	12.58	6.05	6.17	10.51	10.29
T₆	50 %RDF + 5 t/ha ⁻¹ Biochar+ZMB + Zn	12.49	12.69	5.99	6.11	10.55	10.33
T₇	100 %RDF+2.5t/ha ⁻¹ Biochar	12.52	12.71	5.99	6.11	10.69	10.48
T₈	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB	13.47	13.68	6.46	6.58	11.63	11.45
T₉	100 %RDF +2.5t/ha ⁻¹ Biochar +ZMB + Zn	13.44	14.00	6.73	6.42	11.54	11.43
	SEm±	0.38	0.40	0.20	0.21	0.34	0.33
	CD (P= 0.05)	1.10	1.21	0.58	0.62	1.01	0.97

Fig 7 : Effect of Biochar and fertility levels on crop growth rate (g/m²/day) at different intervals of rice



Control
 100 %RDF
 50%RDF
 50%RDF+5t/ha-1Biochar
 50 %RDF + 5 t/ha-1Biochar+ZMB
 50 %RDF + 5 t/ha-1Biochar+ZMB + Zn
 100 %RDF+2.5t/ha-1 Biochar
 100 %RDF +2.5t/ha-1Biochar +ZMB
 100 %RDF+2.5t/ha-1Biochar +ZMB + Zn

UNDER

Conclusion

In conclusion, the study demonstrated the significant influence of biochar and varying fertility levels on rice growth and yield parameters across 2022 and 2023. Biochar consistently enhanced plant height, root length, tiller density, leaf area index, dry matter accumulation, crop growth rate, compared to the control. Treatments integrating biochar with 100% recommended fertilizer dose and biofertilizer, particularly T₉ (100% RDF + 2.5 t/ha Biochar + ZMB + Zn), consistently outperformed other treatments across all parameters measured. This treatment promoted superior growth and enhanced soil health indicators such as microbial populations and organic carbon content. Therefore, T₉ emerges as the optimal treatment choice for maximizing rice productivity and sustainability, offering a balanced approach toward achieving higher benefits in rice cultivation practices.

References

- Akhtar, S., Ullah Md, J., Ahamed Md, K. U., Hamid, A., Islam Md, S., & Islam, M. R. (2021). Sowing date effects on phenology and yield of white maize genotypes during Kharif season under subtropical environment. *Int J Nat Soc Sci*, 8, 81-90.
- Anonymous (2022). Agricultural Statistics at a Glance. Annual data of Economics & Statistics Division, Department of Agriculture & Farmers Welfare, Ministry of Agriculture & Farmers Welfare, Government of India. P. 38-39. DOI: [https://agricoop.nic.in/Documents/CWWGDATA/Agricultural Statistics at a Glance 2022 0. pdf](https://agricoop.nic.in/Documents/CWWGDATA/Agricultural%20Statistics%20at%20a%20Glance%202022%200.pdf)
- Anonymous, 2021-22. Annual Report, Department of Agriculture and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Krishi Bhawan, New Delhi; p:4-5.
- Anonymous, April 2022. Department of Agriculture, National Conference on Kharif Crops, Government of Uttar Pradesh, p: 8-9.
- Cao, X., Liu, J., Wu, Y., Wei, Y., Zhang, L., Li, J., Lv, H. and Idimesheva, O., 2024. Effects of biochar application on rice physiological growth and water use efficiency under dry direct-seeded mode. *Irrigation and Drainage*, 73(2), pp.526-537.
- Gowthami, V., Venkateswarlu, B., Prasad, P.V.N., Rao, C.S. and Kumari, S.R., 2022. Growth and Yield of Rice as Influenced by Rice Husk Biochar and Nitrogen Levels during Kharif Season. *International Journal of Environment and Climate Change*, 12(10), pp.284-289.
- Gul, S., Khan, M. H., Khanday, B. A., & Nabi, S. (2015). Effect of sowing methods and NPK levels on growth and yield of rainfed maize (*Zea mays* L.). *Scientifica*,
- Islam, S.J.M., Mannan, M.A., Khaliq, Q.A. and Rahman, M.M., 2018. Growth and yield response of maize to rice husk biochar. *Australian Journal of Crop Science*, 12(12), pp.1813-1819.
- Kamara, A., Kamara, H.S. and Kamara, M.S., 2015. Effect of rice straw biochar on soil quality and the early growth and biomass yield of two rice varieties. *Agricultural Sciences*, 6(08), p.798.
- Khan, M.A., Basir, A., Fahad, S., Adnan, M., Saleem, M.H., Iqbal, A., Amanullah, Al-

- Huqail, A.A., Alosaimi, A.A., Saud, S. and Liu, K., 2022. Biochar optimizes wheat quality, yield, and nitrogen acquisition in low fertile calcareous soil treated with organic and mineral nitrogen fertilizers. *Frontiers in Plant Science*, 13, p.879788.
- Khan, Z., Nauman Khan, M., Luo, T., Zhang, K., Zhu, K., Rana, M.S., Hu, L. and Jiang, Y., 2021. Compensation of high nitrogen toxicity and nitrogen deficiency with biochar amendment through enhancement of soil fertility and nitrogen use efficiency promoted rice growth and yield. *Gcb Bioenergy*, 13(11), pp.1765-1784.
- Khedwal, R. S., Yadav, D. B., & Hooda, V. S. (2018). Crop Residue management in no-till maize: Influence the growth, yield and economics of kharif maize (*Zea mays* L.). *Forage Res*, **44**, 90-95.
- Koca, Y. O., & Erekul, O. (2016). Changes of dry matter, biomass and relative growth rate with different phenological stages of corn. *Agriculture and Agricultural Science Procedia*, 10, 67-75.
- Kumar, A., Mishra, S. R., Pandey, A. K., Kushwaha, R. K., Singh, S. K., Mishra, A. N., & Singh, A. K. (2020). Effect of heat use efficiency on leaf area index, plant height and yield attributes on kharif maize (*Zea mays* L.) cultivars. *Journal of Pharmacognosy and Phytochemistry*, **9**(5), 2732-2735.
- Liu, B., Li, H., Li, H., Zhang, A. and Rengel, Z., 2021. Long-term biochar application promotes rice productivity by regulating root dynamic development and reducing nitrogen leaching. *Gcb Bioenergy*, 13(1), pp.257-268.
- Oladele, S.O., Adeyemo, A.J. and Awodun, M.A., 2019. Influence of rice husk biochar and inorganic fertilizer on soil nutrients availability and rain-fed rice yield in two contrasting soils. *Geoderma*, 336, pp.1-11.
- Oo, S., 2022. Yield and Yield Contributing Characters of Rice (*Oryza sativa* L.) as Affected by Biochar and Nitrogen Fertilizer Application.
- Ponmozhi, C. I., Kumar, R., Baba, Y. A., & Rao, G. M. (2019). Effect of integrated nutrient management on growth and yield of maize (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences*, **8**(11), 2675-2681.
- Prasad, G. (2017). Effects of Sowing Dates on Yield Attributes and Yield in Maize (*Zea mays* L.) Hybrids Sown In Kharif Season in Haryana. *Chemistry Science Rev Lett*, **6**(24), 2393-2396.

- Rao, C. R., Prasad, P. V. N., & Venkateswarlu, B. (2016). Assessment of different herbicides on yield and economics of kharif maize (*Zea mays* L.). *Intern. J. Agric. Sci. Res*, **6**, 409-414.
- Sharanabasappa, H. C., & Basavanneppa, M. A. (2019). Influence of plant population and fertilizer levels on growth, yield and quality parameters of quality protein maize (*Zea mays* L.) in irrigated ecosystem. *Int. J. Chem. Stud*, **7**, 1425-1429.
- Sial, T.A., Lan, Z., Wang, L., Zhao, Y., Zhang, J., Kumbhar, F., Memon, M., Lashari, M.S. and Shah, A.N., 2019. Effects of different biochars on wheat growth parameters, yield and soil fertility status in a silty clay loam soil. *Molecules*, **24**(9), p.1798.
- Singh, J., Partap, R., & Singh, A. (2021). Effect of nitrogen and zinc on growth and yield of maize (*Zea mays* L.). *International Journal of Bio-resource and Stress Management*, **12**(Jun, 3), 179-185.
- Thavanesan, S. and Seran, T.H., 2018. Effect of rice straw and husk biochar on vegetative growth and yield attributes of *Oryza sativa* L. *International Journal of Crop Science and Technology*, **4**(2), pp.49-56.
- Wang, C., Liu, J., Shen, J., Chen, D., Li, Y., Jiang, B. and Wu, J., 2018. Effects of biochar amendment on net greenhouse gas emissions and soil fertility in a double rice cropping system: A 4-year field experiment. *Agriculture, ecosystems & environment*, **262**, pp.83-96.