

Short Research Article

Effect of neem (*Azadirachta indica*) and castor (*Ricinus communis*) leaf extracts on root-knot nematodes of the genus *Meloidogyne* in tomato (*Solanum lycopersicum*) culture

ABSTRACT

Aims: During production, tomatoes are attacked by root-knot nematodes of *Meloidogyne* genus, which are probably the most damaging. Given the harmful effect of certain nematicides used to control this nematode, the use of plant extracts could be an alternative for sustainable tomato production. The aim of this study was to evaluate the efficacy of neem and castor leaf extracts on root-knot nematodes in the field.

Place and Duration of Study: Côte d'Ivoire, from April to August 2023.

Methodology: A plot previously infested with root-knot nematodes was set up using a randomized complete block design. Neem and castor extracts were applied at a dose of 100 g per tomato plant, and furasol at 5 g, planted once a month until harvest. The density of root-knot nematodes was estimated in the soil each month prior to extract application and in tomato roots at harvest. Parameters such as severity and prevalence of galls, nematode density in soil and roots, plant height and fruit yield were measured during the study.

Results: The prevalence and severity of root galls on tomato plants were statistically identical on plants treated with furasol and those treated with castor oil. The results showed that neem and castor extracts had similar effects to furasol on tomato plants in terms of plant height and fruit yield. These plant extracts induced a low rate of multiplication of these nematodes in the same way as furasol, and led to a reduction in the number of nematodes in the soil and roots of treated tomato plants.

Conclusion: The results obtained in this study are encouraging, and the long-term use of these plant extracts could be a sustainable control method against root-knot nematodes.

Keywords: plant extracts, *Meloidogyne*, tomato, severity, prevalence

1. INTRODUCTION

Market garden produce plays an important role in the general diet of human beings. Among these products, tomatoes are an important source of minerals, essential amino acids, sugars, dietary fiber and vitamins, as they contain vitamin A in the form of carotene, vitamin B1 (thiamine), B2 (riboflavin), niacin and vitamin C (Sesso *et al.*, 2002). World production in 2022 amounted to over 186 million tonnes, compared with 44578.83 tonnes for Côte d'Ivoire (Faostat, 2022). Although tomatoes are nutritionally and economically important, their production is limited by biotic and abiotic factors (Oluwatayo *et al.*, 2019). During cultivation and after harvest, tomatoes can be affected by over 200 pathogens, including fungi, bacteria, viruses and viroids, as well as parasitic nematodes (Panno *et al.*, 2021). In sub-Saharan Africa, particularly Côte

Commented [1]: Rewrite in active voice

Commented [2]: Insert prefix "the"

Commented [3]: Rewrite as "Study duration"

Commented [4]: Rewrite in active voice

Commented [5]: Rewrite in active voice

Commented [6]: delete

Commented [7]: Delete coma

Commented [8]: Rewrite in active voice

Commented [9]: Replace with "before"

Commented [10]: Rewrite the sentence in active voice

Commented [11]: delete

Commented [12]: Replace the words with "regarding"

Commented [13]: Replace the words with "reduced"

Commented [14]: Replace with "a vital"

Commented [15]: Rewrite the sentence in active voice

Commented [16]: Add coma

Commented [17]: Replace the words with "and"

d'Ivoire, tomatoes are attacked by several pests, including nematodes (Doga *et al.*, 2022). These are microscopic worms, mostly living in the soil, which parasitize plants. They severely infect almost all crops worldwide, causing an estimated annual global crop loss of around 80 billion USD (Jones *et al.*, 2013). Historically, the main method of controlling nematode infestations was chemical control, such as the fumigant methyl bromide (Desaeger *et al.*, 2020). These chemicals are known to have a negative impact on the crop environment, such as a decrease in the distribution of beneficial microorganisms in the soil, an increase in chemical-resistant pests, environmental pollution due to residual toxicity and toxicity to users (Kiewnick and Sikora, 2006). Recently, alternative products for nematode control have come into use (Jagdale and Grewal, 2002). These include botanical extracts with a nematicidal effect, which are one of the potential alternatives for sustainable control of root-knot nematodes (Gahukar, 2012). This natural pest control strategy can improve both soil quality and human health by reducing harmful residues or contamination (Hulot and Hiller, 2021). Authors such as Alam (1989) and Mojumder (1995) have confirmed the efficacy of soil-incorporated neem (*Azadirachta indica*) and castor (*Ricinus communis*) cakes against *Meloidogyne* spp, as well as other plant-parasitic nematode species. The aim of the present study was to evaluate the efficacy of neem and castor leaf extracts on root-knot nematodes in the field.

Commented [18]: delete

Commented [19]: Replace with 'parasiting'

Commented [20]: Insert "and"

2. MATERIAL AND METHODS

2.1. Material composition

The material used in this study consisted of neem (*Azadirachta indica*) and castor (*Ricinus communis*) leaves, as well as the chemical nematicide (furasol) for the treatments and tomato seeds of the Cobra 26 variety.

2.2. Choice of plot

The experimental plot was chosen on the basis of its cropping history. In fact, this plot was previously used to grow tomatoes, which were severely infested by root-knot nematodes. Before setting up the trial, soil samples were taken and analyzed in the laboratory, in order to assess the proportion of root-knot nematodes in 100 g of soil.

Commented [21]: Rewrite the sentence in active voice

Commented [22]: Replace with "based on"

Commented [23]: Rewrite the sentence in active voice

Commented [24]: delete

2.3. Preparation of neem and castor extracts

The extracts used in this study were castor and neem leaf powders. To prepare these extracts, fresh leaves from each plant were collected. They were dried in the shade for a period of 2 weeks, then ground to powder using a blender. The powders obtained were used at a dose of 100 g/plant for the tests.

Commented [25]: Rewrite the sentences in active voice

Commented [26]: delete

Commented [27]: Rewrite the sentence in active voice

2.4. Setting up the experimental plot

The experiment was carried out on a 282 m² (23.5 m x 12 m) plot from April to August 2023, following a complete randomized block design with four blocks separated by 1.5 m each. Each block was made up of four elementary plots measuring 2 m x 2 m. Thirty seedlings were planted in each elementary plot. The individual pots were spaced 0.5 m apart on the row and between rows. Each elementary plot thus comprised 30 tomato plants.

Commented [28]: Rewrite the sentence in active voice

Commented [29]: Rewrite the sentence in active voice

A nursery was set up by sowing seeds in honeycomb plates containing soil amended with 50 g of each plant extract, and 2 g for the synthetic nematicide. In the case of control plants, the soil was left untreated.

Commented [30]: Delete coma

At the same time, the plant extracts were applied to the planting plots at a depth of 10 cm, depending on the treatment. The plots were watered daily to promote decomposition of the plant extracts. Twenty-one days later, the vigorous seedlings were transplanted into the pits. The treatments carried out were as follows :

Commented [31]: Rewrite the sentence in active voice

Commented [32]: Add prefix "the"

Commented [33]: Rewrite the sentence in active voice

- T⁻ : tomato plants not treated with plant extracts or synthetic nematicides;
- T⁺ : tomato plants treated with synthetic nematicide (furasol) ;
- T_{Fn} : tomato plants treated with neem leaf powder ;
- T_{Fc} : tomato plants treated with castor leaf powder.

The experiment was carried out under natural conditions, without the application of chemical fertilizers, and was repeated twice.

Commented [34]: Replace with "applying"

Commented [35]: Rewrite in active voice

After transplanting the tomato seedlings, 100 g of plant extracts were applied monthly to the plants until harvest, to assess their efficacy against root-knot nematodes. For furasol, 5 g were applied.

Commented [36]: Delete coma

2.5. Determination of nematode density as a function of treatments carried out

The density of root-knot nematodes was estimated in soil and tomato roots. In the case of soil, this density was determined each month before the application of plant extracts, in order to determine the evolution of nematode density in 100 g of sample. For this purpose, 10 plants per elementary plot were randomly selected and soil samples were taken from their roots. The density of nematodes in the roots was determined at harvest on the same tomato plants in 100 g of root samples.

Commented [37]: Rewrite in active voice

Commented [38]: delete

Commented [39]: Replace with "assess"

Commented [40]: Rewrite the sentence in active voice

Commented [41]: Rewrite the sentence in active voice

2.6. Determination of tomato agronomic parameters according to treatments

The following parameters were determined at harvest on 10 randomly selected plants per elementary plot :

Commented [42]: Rewrite in active voice

- plant height, measured using a tape measure ;
- fruit yield, determined according to treatments using the following formula :

Commented [43]: Capitalised P

Commented [44]: CAPITALISED f

$$\text{Yield (t/ha)} = \frac{\text{Total weight of harvested fruit (t)}}{\text{Surface area (ha)}}$$

- degree of fruit firmness, determined on a scale from 1 to 5 (1 = completely soft fruit, 2 = soft fruit, 3 = moderately firm fruit, 4 = firm fruit, 5 = very firm fruit) from Noupé *et al.* (2019).

Commented [45]: Capitalised d

Commented [46]: Replace with "substantial"

In addition to these agronomic parameters, the number of nematode-induced galls on tomato plant roots was determined by counting, and the nematode multiplication rate was determined using the following formula :

Commented [47]: Rewrite in active voice

$$Tm (\%) = \frac{Nnf}{Nni} \times 100$$

Tm : multiplication rate ;
Nnf : final number of nematodes ;
Nni : initial number of nematodes.

2.7. Evaluation of the severity and prevalence of galls on the root system of tomato plants according to treatments

In order to assess the effect of the different treatments on root-knot nematode damage on treated tomato plants, severity and prevalence were assessed on their root systems at harvest.

2.7.1. Assessment of gall prevalence

Gall prevalence was assessed using an X pattern, i.e. by selecting 60 plants along the two diagonals of each plot. On each diagonal, 30 plants of roughly equal distance were selected. The number of plants showing galls was then recorded, and the prevalence of this symptom was calculated using the following formula :

Commented [48]: Rewrite in active voice

Commented [49]: Rewrite in active voice

$$PM(\%) = \frac{Pt}{N} \times 100$$

PM: Average prevalence of galls, Pt = number of plants with galls, N = total number of plants selected

2.7.2. Assessment of gall severity

To assess the severity of galls, the Zeck scale (1971) was used. This scale corresponds to a rating of root systems from 0 to 10, based on observation of the state of the root system, and reflects the reality of symptomatological situations observed on tomatoes. It is summarized as follows :

0 = root system completely free of galls; 1 = a few rare and small galls detected during close observation; 2 = a few rare and small galls easily detected; 3 = numerous small galls; 4 = numerous small galls, a few large galls ; 5 = 25% of root system gall-affected and non-functional (no rootlets); 6 = 50% of root system gall-affected; 7 = 75% of root system gall-affected; 8 = no root unaffected, plant still green; 9 = root system decaying; 10 = plant dead. The severity of the galls was calculated using the formula of McKinney (1923) :

$$S(\%) = \frac{\sum(n_i \times n_i)}{N_t \times n_{ie}} \times 100$$

S(%) : severity of galls ;

n_i : severity score assigned to galls on the plant ;

n_t : number of plants to which the score n_i was assigned ;

N_t : total number of plants used ;

n_{ie} : highest severity score recorded in this study.

2.8. Statistical analysis

The data collected in this study were analyzed using SPSS version 26 software. Tests of variance homogeneity and data distribution were performed to determine the choice of parametric or non-parametric test. The severity and prevalence of galls, the number of nematodes in soil and roots, nematode multiplication rates, plant height, fruit yield and fruit firmness were analyzed using the non-parametric Kruskal-Wallis test (comparison of several means). In the event of a significant difference, at the 5% threshold, the Mann-Whitney U test was used to compare modalities in pairs to form homogeneous groups.

3. RESULTS

3.1. Description of plants obtained according to treatments

The tomato plants obtained showed different aspects. Plants treated with neem, castor oil and furasol generally showed green leaves with fairly vigorous stems (Fig. 1 A) and firm fruit (Fig. 1 B). The roots of these plants showed small galls that were barely detectable at first glance. Untreated plants, on the other hand, showed a rather different appearance for the most part. They showed yellow discoloration on the leaves, dwarfing on some plants (Fig. 1 C) and galls on the roots (Fig. 1 D).

3.2. Agronomic parameters obtained according to treatments

These parameters changed with each treatment.

Plant height varied from 48.60 ± 17.50 cm to 78.40 ± 11.27 cm. Plant height was highest in plants treated with castor extract, at 78.40 ± 11.276. However, no difference was observed in plants treated with neem, castor oil and furasol. On the other hand, statistical tests revealed a significant difference between these 3 treatments and the control (Table 1).

In terms of fruit firmness, the plants treated with furasol, neem and castor were firmer. With these three products, firmness was greater than 4. Statistical tests showed a significant difference between the firmness of fruit obtained with these three extracts and that of fruit from untreated plants (Table 1).

Tomato plant yields ranged from 10.32 ± 2.89 t/ha to 17.80 ± 2.54 t/ha (Fig 2). The lowest yield was observed in untreated plants. In contrast, the highest yield was observed in plants treated with castor extract, followed by those treated with neem and finally furasol. However, the yields obtained from plants treated with these three products were statistically identical. Nevertheless, the yields obtained with these three products were statistically different from those obtained with untreated plants (Fig. 2).

Commented [50]: Delete the words

Commented [51]: Capitalised t

Commented [52]: Replace with "uttery"

Commented [53]: Rewrite in active voice

Commented [54]: Rewrite in active voice

Commented [55]: Add coma

Commented [56]: Rewrite in active voice

Commented [57]: Add coma

Commented [58]: Replace with "pretty"

Commented [59]: Replace the words with "leaf discoloration"

Commented [60]: Add coma

Commented [61]: Rewrite in active voice

Commented [62]: Add coma

Commented [63]: Rewrite as "three"

Commented [64]: Replace the words with "Regarding"

Commented [65]: Add coma

Commented [66]: delete

Commented [67]: Rewrite in active voice

Commented [68]: Rewrite in active voice




Fig. 1 . Aspect of tomato plants observed according to extract treatments

A: leaves of a tomato plant treated with neem; B: tomato fruit of a plant treated with castor oil; C: untreated plant showing symptoms of leaf yellowing; D: galls on the roots of an untreated plant.

Table 1 : tomato plant height and fruit firmness as a function of treatments

Traitements	Plant height (cm)	Fruit firmness
T ⁻ (Indicator)	48,60 ± 17,506 ^a	3,02 ± 0,48 ^a
T ⁺ (Furasol)	72,40 ± 18,38 ^b	4,20 ± 0,632 ^b
Neem	73,00 ± 17,506 ^b	4,80 ± 0,422 ^b

Ricin	$78,40 \pm 11,276^b$	$4,70 \pm 0,483^b$
--------------	----------------------	--------------------

In each column, means marked with the same letter are statistically similar according to the Mann-Whitney U test at the 5% threshold.
p = probability value.




Fig. 2 . Average yield of tomato fruit according to treatments carried out
Means marked with the same letter are statistically similar according to the Mann-Whitney U test at the 5% threshold.

3.3. Prevalence and severity of symptoms observed

The prevalence of nematode-induced galls ranged from $30.33 \pm 2.94\%$ to $71.23 \pm 3.62\%$ according to the different treatments. The prevalence of nematode-induced galls was higher in untreated tomato plants (controls) than in those treated with aqueous extracts of neem, castor oil and furasol. In the case of treated plants, the prevalence of galls was statistically lower in those treated with castor oil and furasol than in those treated with neem (Table 2).

The severity of galls varied from $25.89 \pm 1.33\%$ to $60.51 \pm 2.58\%$, depending on the different treatments. The highest severity was observed in untreated tomato plants. Statistical analysis showed that there was a significant difference between the severity of these plants and those treated with neem, castor oil and furasol. On the other hand, tomato plants treated with castor oil and furasol had statistically lower severities than those treated with neem (Table 2).

3.4. Evolution of nematode density in the soil and their numbers in the roots

At harvest, the number of nematodes in the roots of tomato plants with galls ranged from 4380.67 ± 47.39 to 16280.35 ± 1327.734 . The highest value was observed in untreated plants. Statistical analysis showed a significant difference between the number of nematodes extracted from tomato roots of untreated plants and those treated with neem, castor oil and furasol ($P < 0.05$). In contrast, the number of nematodes extracted from tomato plants treated with neem, castor oil and furasol was statistically identical (Table 3).

The rate of nematode multiplication in roots was higher in untreated plants than in those treated with neem, castor oil and furasol. Statistical analysis confirmed a significant difference in nematode multiplication rates between untreated plants and those treated with neem, castor oil and furasol. For all three treatments, the multiplication rate was statistically identical (Table 3).

The density of nematodes in the soil varied according to the sampling periods and treatments carried out. In fact, for tomato plants that had not undergone any treatment, density increased over time. It rose from

Commented [69]: delete

Commented [70]: Add coma

Commented [71]: Rewrite in active voice

Commented [72]: Delete the words

Commented [73]: Rewrite in active voice

Commented [74]: Add coma

Commented [75]: Add coma

Commented [76]: Add coma

Commented [77]: Add coma

Commented [78]: Delete the words

Commented [79]: Insert the words " density increased over time"

Commented [80]: Delete the words

176 ± 15.05 in the first month, to 673 ± 84.72 in the fourth month. On the other hand, in plants treated with neem, castor oil and furasol, a decrease in density was observed from the first to the fourth month after sowing. For tomato plants treated with castor oil, density fell from 86 ± 10.75 to 70 ± 8.16. For plants treated with neem, density was 91 ± 16.63 in the first month after sowing and 69 ± 7.37 in the fourth month after sowing. For plants treated with furasol, density fell from 99 ± 13.70 to 76 ± 9.66 between the first and fourth months (Fig. 3).

Statistical analysis showed that there was a significant difference in nematode density between untreated tomato plants and those treated with neem, castor oil and furasol. However, no significant difference was observed between tomato plants treated with neem, castor oil and furasol at the different sample collection periods.

Commented [81]: Delete coma

Commented [82]: Add coma

Commented [83]: Rewrite the sentence in active voice

Commented [84]: Delete the words

Commented [85]: Dd coma

Commented [86]: Rewrite in active voice

Table 2 : Mean prevalence and severity of galls caused by nematodes on tomato roots according to treatments carried out

Treatments	Prevalence (%)	Severity (%)
T ⁻ (Indicator)	71,23± 3,62 ^c	60,51 ± 2,58 ^c
T ⁺ (Furasol)	30,33± 2,94 ^a	26,62 ± 1,03 ^a
Neem	35,43± 2,55 ^b	30,5 ± 1,67 ^b
Ricin	30,67 ± 3,91 ^a	25,89 ± 1,33 ^a

In each column, means marked with the same letter are statistically similar according to the Mann-Whitney U test at the 5% threshold.
p = probability value.

Table 3 : Number of nematodes extracted from the roots of tomato plants and their reproduction rate as a function of the treatments applied

Treatments	Number of nematodes in roots	Nematode multiplication rate
T ⁻ (Indicator)	26280,35 ± 1327,73 ^c	2,24 ± 0,52 ^b
T ⁺ (Furasol)	4380,67 ± 47,39 ^a	0,37 ± 0,00 ^a
Neem	5860,53 ± 50,35 ^b	0,49 ± 0,00 ^a
Ricin	5381,33 ± 11,66 ^b	0,45 ± 0,00 ^a

In each column, means marked with the same letter are statistically similar according to the Mann-Whitney U test at the 5% threshold.

p = probability value.




Fig. 3 . Number of *Meloidogyne* J2 in the soil according to treatment and period after sowing

4. DISCUSSION

In the field, neem and castor leaf extracts, together with the chemical nematicide furasol, reduced the number of nematodes in the soil and roots of tomato plants, as well as their rate of multiplication. Indeed, neem contains azadirachtin, which may induce nematostasis, a process that prevents nematodes from invading plants without killing them directly, according to Rehman *et al.* (2009). Such an effect was observed by Gharras *et al.* (2011) in zucchini. These authors showed that treatments with castor oil cake, argan oil cake, argan leaves and neem oil cake reduced nematodes in the soil by more than 25% and the gall index by more than 50%. Affokpon *et al.* (2012) also obtained similar results in their study. In fact, they showed that the use of neem seed derivatives significantly reduced the rate of root-knot nematode multiplication in tomato and greater nightshade crops. Similarly, this rate was reduced by 81% in plots of greater nightshade treated with neem seed derivatives. Castor powder also had a reducing effect on root-knot nematodes of the *Meloidogyne* genus in tomato plots, with a rate of around 51% compared with that of untreated plots.

These powders also improved certain agronomic traits of the plants as well as tomato fruit yield. For Khan *et al.* (2012), the improvement in crop yields on root-knot nematode-infested soil following the application of neem derivatives was due both to nematode control and improved soil fertility levels. For Akhtar and Alam (1993), the triterpene compounds in neem seeds inhibit the nitrification process and increase the quantity of available nitrogen, in addition to their nematicidal effects, which explains the plants' growth and good yield. Gharras *et al.* (2011) showed in their study that argan, neem and castor oil cakes resulted in very significant vegetative growth. Indeed, the addition of these extracts significantly improved plant height and tomato fruit yield. Moreover, an increase in plant height compared with the control was observed with castor oil and neem.

Commented [87]: Replace the words with "They"

Commented [88]: Replace the words with "using"

Commented [89]: Replace the word with "more excellent"

Commented [90]: Rewrite in active voice

Commented [91]: Replace with "reduced the "

Commented [92]: delete

Commented [93]: delete

Commented [94]: delete

Commented [95]: delete

Commented [96]: Replace the words with "adding"

Commented [97]: Rewrite in active voice

5. CONCLUSION

At the end of this study, it was found that neem and castor leaf extracts significantly reduced the evolution of nematodes in the soil and in tomato roots, and their effect was comparable to that of the synthetic nematicide used. These extracts also improved certain agronomic parameters, such as fruit yield and firmness, as well as the size of treated tomato plants. These plants offer a promising option for sustainable control of root-knot nematodes.

Commented [98]: Rewrite in active voice

Commented [99]: Replace with "specific"

COMPETING INTERESTS

Authors have declared that no competing interests exist.

Commented [100]: Add the prefix "The"

REFERENCES

1. Affokpon, A., Dan, C, Houédjissi, M, Hèkpazo B, & Tossou, C (2012). The efficacy of neem seed derivatives against root-knot nematodes in market gardening differs

UNDER PEER REVIEW

depending on the type of derivative. *Benin Agricultural Research Bulletin (BRAB)*, 72: 48-58.

2. Alam. MM. (1989). Control of Root-knot and stunt nematode with horn meal, bone meal and silseed cakes. *Indian Journal of Nematology*, 19: 166 -170.

3. Akhtar, M., & Alam, MM (1993). Control of plantparasitic nematodes by 'Nimin' an ureacoating agent and some plant oils, 100: 337342.

4. Coulibaly. K (2016). Impacts of dry sequences on rice crop calendars in Côte d'Ivoire. Master's thesis in Geography, IGT, Felix Houphouët Boigny University, Abidjan, Côte d'Ivoire, 100 p

5. Desaegeer, J., Wram, C, & Zasada, I (2020). New reduced-risk agricultural nematicides - justification and review *Journal of Nematology*, 1: 52. pp. 1–16

6. Diarra, A., & Tagbo, MP (2022). Climate variability and rainfed rice production in humid areas: the case of the Gagnoa sub-prefecture (Côte d'Ivoire)

7. Doga, D., Ouattara, K, Zirihi, G, & Zeze, A (2022). Nematicidal property of the aqueous extract of *Crotalaria retusa* L. on *Meloidogyne* spp in vitro *Journal of Animal and Plant Sciences*, 54 (3), 9952-9957

8. Faostat. (2022). <https://www.fao.org/faostat/en/#data>

9. Gahukar. RT (2012). Evaluation of plant-derived products against pests and diseases of medicinal plants: a review. *Crop Protection*, 42: 202-209.

10. Hulot, JF., Hiller, N (2021). Exploring the benefits of biocontrol for sustainable agriculture: a literature review on biocontrol in light of the European Green Deal. Institute for European Environmental Policy, 37 p.

11. Jagdale, GB., Grewal, PS (2002). Identification of alternatives for the management of foliar nematodes in floriculture. *Pest Management Science*, 58(5), 451-458.

12. Jones, J, T., Haegeman, A, Danchin, EGJ, Gaur, HS, Helder, J, Jones, MGK, & Perry RN (2013). Top 10 plant-parasitic nematodes in molecular plant pathology, *Molecular Plant Pathology*, 14 (9), 946-961.

13. Khan, MR., Mohiddin, FA, Ejaz, MN, Khan, M (2012). Management of root-knot disease in eggplant through the application of biocontrol fungi and dry neem leaves. *Turkish Journal of Biology*, 36 (2), 161-169.
14. Kiewnick, S., & Sikora, R.A. (2006). Biological control of the root-knot nematode *Meloidogyne incognita* by *Paecilomyces lilacinus* strain 251. *Biological control*, 38 (2), 179-187.
15. Mojumder. V (1995). Effect of neem on nematodes. In "The neem tree: source of unique natural products for integrated pest management, medicine, industry and other purposes", SCHMUTTERER H., VCH Publication, New York, USA, p.129-150.
16. Oluwatayo, JI., Jidere, CI, Nwankiti, A (2019). Nematicidal Effect of Some Botanical Extracts for the Management of *Meloidogyne incognita* and on Growth of Tomato. *Asian Journal of Agricultural and Horticultural Research*, 4(2), 1-8, 2019
17. Panno, S., Davino, S, Caruso, AG, Bertacca, S, Crnogorac, A, Mandić, A, Noris, E, & Matic, S. (2021). A review of the most common and economically important diseases affecting tomato cultivation in the Mediterranean basin. *Agronomy*, 47 p.
18. Rehma, AU., Javed, N, Ahmad, R, Shahid, M (2009). Protective and curative effect of bioproducts against the invasion and development of root-knot nematode in tomato. *Pakistan Journal of Phytopathology*, 21(1), 37-40
19. Sesso, HD., Lius, GJM, Buring JE (2003). Dietary lycopene tomato - Based food products and cardiovascular disease in women. *Journal of Nutrition*,133(7), 2336-2341.