

Optimizing Planting Date and Spacing for Enhanced Yield and Oil Production of *Kaempferia galanga* L.: A Field Experiment.

Abstract

Kaempferia galanga L., a rhizomatous plant species, is highly valued for its medicinal and culinary properties. Studying the effects of planting dates and spacing is crucial for optimizing the cultivation of *K. galanga*, as it can help farmers and cultivators maximize yields, improve resource allocation, and promote sustainable agricultural practices. This study investigated the impact of planting date and spacing on the growth and yield of *K. galanga*. The experiment was conducted over three consecutive years, with three planting dates (April 15, April 30, and May 15) and four spacing treatments (20 cm x 20 cm, 30 cm x 30 cm, 40 cm x 40 cm, and 50 cm x 50 cm). The results showed that planting date and spacing significantly affected canopy diameter, yield per plant, rhizome yield per hectare, and oil yield per hectare. The optimal planting window for maximizing yield and oil production was April 15-30, with a spacing of 20 cm x 20 cm. Wider spacing resulted in higher yield per plant, while closer spacing led to higher rhizome yields per hectare.

Keywords

Kaempferia galanga L., Planting date, Spacing, Rhizome yield, Agronomic optimization

Introduction

Kaempferia galanga L., commonly known as Kencur, Sand Ginger, or lesser Galangal, is a plant species belonging to the Zingiberaceae family. Native to tropical Asia, particularly in India, Southeast Asia, and southern China, this herb has been used for centuries in traditional medicine, cooking, and rituals (Zeitler, 2024). *K. galanga* prefers well-drained soil and partial shade, making it a popular crop in tropical regions (Kumar, 2007). The plant grows up to 30 cm tall, with narrow, lance-shaped leaves and small, white or yellowish flowers. The rhizome is the most valuable part, with a pungent, aromatic flavor and a sweet, slightly bitter taste.

In traditional medicine, the rhizome treats various ailments like digestive issues, fever, and respiratory problems. The plant is also used as a spice and flavoring agent in traditional dishes like

curries and soups. *K. galanga* is also utilized in traditional ceremonies and rituals and is valued for its spiritual and medicinal significance. Recent studies have validated the traditional uses of *K. galanga*, revealing its potential pharmacological properties, including anti-inflammatory, antioxidant, and antimicrobial activities (Munda, *et al.*, 2018). The plant's essential oils and bioactive compounds have also been explored for potential applications in medicine and industry (Kumar, 2020).

The demand for *K. galanga* rhizomes is increasing due to their medicinal properties, making it essential to standardize agronomic practices that ensure higher yield and quality. Planting date and spacing are critical factors that affect crops' growth, yield and quality (Abbas *et al.*, 2019). The optimal planting date and spacing can vary depending on the crop species, climate, and soil conditions. This study aimed to standardize the planting dates and spacing for *Kaempferia galanga* L. to achieve higher rhizome yields.

Materials and Methods

The experiment was conducted at the All India Coordinated Research Project on Medicinal, Aromatic plants, and Betel vine (AICRP on MAP&B), Thrissur, Kerala, India, over three consecutive years (2021, 2022, and 2023). The experimental site is located in a humid tropical climate region, with geographical coordinates of 13° 32'N latitude and 76° 26'E longitude and an elevation of about 40 meters above Mean Sea Level (MSL).

The soil at the experimental site is a lateritic sandy loam of the Oxisol group, comprising 46.8% sand, 21.6% silt, and 31.3% clay. The soil has a moderate organic carbon content of 0.49% and available nutrients, including 250.21 kg/ha nitrogen, 30.96 kg/ha phosphorus, and 240.50 kg/ha potassium. The soil pH is slightly acidic, measuring 5.3.

The experiment was laid out in a split-plot design, with three planting dates (April 15, April 30, and May 15) as the main plot and four spacing treatments (20 cm x 20 cm, 30 cm x 30 cm, 40 cm x 40 cm, and 50 cm x 50 cm) as the subplot. Each plot measured 3 meters x 3 meters. The rhizomes were harvested after 9 months of planting, and data were recorded on canopy diameter, yield per plant, rhizome yield per hectare, and oil yield per hectare.

The data were statistically analyzed using the KAU GRAPES online software, as described by Gopinath *et al.* (2020).

Results And Discussion

Direct effect of planting time and spacing

The canopy diameter of the plants was significantly influenced by the spacing treatment but not by planting dates (Table 1.). The widest spacing of 50 cm x 50 cm resulted in the highest canopy diameter (48.44 cm), followed by the spacing of 40 cm x 40 cm (47.78 cm). In contrast, the closest spacing of 20 cm x 20 cm resulted in the lowest canopy diameter (41.33 cm).

Similarly, the yield per plant was affected by planting date and spacing. The April 30 planting date resulted in the highest yield per plant of 71.92 g, which was on par with the April 15 planting (70.33 g). The widest spacing (50 cm x 50 cm) resulted in the highest yield per plant (84.44 g), followed by 40 cm x 40 cm (75.67 cm). The lowest per-plant yield was obtained in the closest spacing of 20 cm x 20 cm (43.89 cm) (Table 1., Fig.1 and Plate 1.).

The variations in yield due to planting dates can be attributed to the prevailing weather parameters during the plant growth period. Research has shown that optimal weather conditions, such as temperature and rainfall, during the growing season can significantly impact plant growth and development (Tripura and Hegde, 2021). The variation in rhizome yield per plant can be attributed to the differences in planting dates and spacings. Wider spacing provided more plant growth space, resulting in higher rhizome yields. In contrast, narrower spacing resulted in competition among plants for resources, leading to lower rhizome yields.

The narrower spacing of 20 x 20 cm resulted in a higher plant population and rhizome yields per hectare (13289 kg/ha). In contrast, the wider spacing of 50 x 50 cm resulted in a lower plant population, leading to lower rhizome yields per hectare (3555 kg/ha). Appropriate plant density is an important crop management practice that can optimize canopy light distribution and increase canopy photosynthetic capacity (Yao *et al.*, 2016). Several studies have emphasized the significance of optimal plant density in achieving maximum yield potential in various crops (Meng *et al.*, 2022; Jia *et al.*, 2023; Osunleti *et al.*, 2023).

The oil yield per hectare was also affected by planting date and spacing. The April 15 planting date resulted in the highest oil yield per hectare (82.62 kg/ha), indicating that this date allowed for optimal growth and development of the plants. Similarly, the closest spacing (20 cm x 20 cm) resulted in the highest oil yield per hectare.

Interaction effect of planting dates and spacing

The interaction between planting dates and spacing significantly impacted rhizome yield per plant, per hectare, and oil yield (Table 2). Consistent with these findings, Bertila *et al.* (2020) reported that planting time significantly influenced the quality of ginger rhizome, including oil content, dry recovery, and oleoresin content.

The highest rhizome yield per plant (88.33 g/plant) was achieved by planting on April 30 with a spacing of 50 x 50 cm, closely followed by planting on April 15 with the same spacing (85.67 g/plant). In contrast, lower per-plant yields were observed with May planting and closer spacing of 20 cm x 20 cm.

In contrast, the per hectare rhizome yield showed a different trend, with higher yields achieved through closer spacing (20 cm x 20 cm) combined with April planting (April 15 or 30). A similar trend was observed for oil yield per hectare, underscoring the complex interplay between planting dates, spacing, and yield parameters. Notably, yield per plant increased with wider spacing, but yield per hectare was influenced more by plant density. This finding is consistent with the report by Divya and Sindhu (2023), who observed similar trends in *Cucuma aromatica*.

Conclusion

This study conclusively shows that planting date and spacing significantly impact the growth and yield of *K. galanga*. The optimal planting window for maximum yield and oil production is April 15-30, with a spacing of 20 cm x 20 cm. These findings have significant implications for *K. galanga* cultivation and production, informing evidence-based agronomic practices to boost productivity and sustainability. Environmental factors, such as soil type, moisture levels, and pest management, might have influenced the results. Hence, future research should address these limitations by incorporating comprehensive environmental and soil management data. Replicating this study in diverse environments and exploring the effects of various soil types, climate conditions, and management practices on *K. galanga* growth and yield would be highly beneficial.

By acknowledging and addressing these limitations, future studies can build upon the findings presented here, ultimately contributing to a more robust understanding of *Kaempferia galanga* L. cultivation and optimization.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

1. Abbas, G., Younis, H., Naz, S., Fatima, Z., Hussain, S., Ahmed, M., & Ahmad, S. (2019). Effect of planting dates on agronomic crop production. In *Agronomic Crops: Volume 1: Production Technologies* (pp. 131-147).
2. Bertila, S., Ariina, M. S., Maiti, C. S., Gadi, Y., & Daiho, L. (2020). Effect of planting time and harvest on yield and quality of ginger (*Zingiber officinale* Rosc.) cv. Nadia. *IJCS*, 8(5), 1922-1925.
3. Divya, V. U., & Sindhu, P. V. (2023). Spacing and propagule size on yield and quality of *Curcuma aromatica* Salisb. *Journal of Crop and Weed*, 19(1), 266-270.
4. Gopinath, P. P., Prasad, R., Joseph, B., & Adarsh, V. S. (2020). GRAPES: General Rshiny Based Analysis Platform Empowered by Statistics (Version 1.0.0) [Computer software].
5. Jia, T. T., Chen, B., & Ma, M. (2023). Effects of planting density on the growth, taproots yield and quality of *Glycyrrhiza uralensis*. *Legume Research-An International Journal*, 46(1), 62-68.
6. Kumar, A. (2020). Phytochemistry, pharmacological activities and uses of traditional medicinal plant *Kaempferia galanga* L.- An overview. *Journal of Ethnopharmacology*, 253, 112667.
7. Kumar, B. M. (2007). Agroforestry systems and practices of Kerala. In *Agroforestry systems and practices of India* (pp. 459-483). New India Publishing Agency.
8. Meng, C., Wang, Z., Cai, Y., Du, F., Chen, J., & Xiao, C. (2022). Effects of planting density and nitrogen (N) application rate on light energy utilization and yield of maize. *Sustainability*, 14(24), 16707.

9. Munda, S., Saikia, P., & Lal, M. (2018). Chemical composition and biological activity of essential oil of *Kaempferia galanga*: A review. *Journal of Essential Oil Research*, 30(5), 303-308.
10. Osunleti, S. O., Olorunmaiye, P. M., Adeyemi, O. R., Asiribo, O. E., & Lagoke, S. T. O. (2023). Growth and yield of mango ginger (*Curcuma amada* Roxb.) as influenced by plant density, organo-mineral fertilizer, and weeding frequency. *Journal of Plant Nutrition*, 46(7), 1377-1390.
11. Shekhar, G. C., & Hore, J. K. (2021). Growth and yield of ginger (*Zingiber officinale* Rosc.) as influenced by different seed size and spacing.
12. Tripura, U., & Hegde, N. K. (2021). Influence of planting dates and VA-Mycorrhiza association on growth, yield and yield attributes of turmeric. *Plant Cell Biotechnology and Molecular Biology*, 1-6.
13. Yao, H., Zhang, Y., Yi, X., Zhang, B., & Gong, J. (2016). Cotton responds to different plant population densities by adjusting specific leaf area to optimize canopy photosynthetic use efficiency of light and nitrogen. *Field Crops Research*, 188, 10-16.
14. Zeitler, L. (2024). *Kaempferia galanga* L. (Aromatic Ginger): A high potential Zingiberaceae species. In *Advances in Medicinal and Aromatic Plants* (Vol. 1, pp. 1-283). Apple Academic Press.

Table 1. The effect of planting dates and spacing on crop diameter (cm), yield per plant (g), yield (kg/ha) and oil yield (kg/ha) of *Kaempferia galanga* L. (Pooled mean of three years)

Treatments	Canopy diameter (cm)	Rhizome yield/plant (g)	Rhizome yield (kg/ha)	Oil yield (kg/ha) Fresh weight basis
Main plot (Date of planting)				
April 15	45.08	70.33	8262	82.62
April 30	47.17	71.92	7889	78.89
May 15	44.75	64.42	7117	717.0
CD(0.05)	NS	2.67	375	5.01
Subplot (Spacing)				
20 cm x 20 cm	41.33	43.89	13289	13.28
30 cm x 30 cm	45.11	71.56	8829	88.29
40 cm x 40 cm	47.78	75.67	5220	52.20
50 cm x 50 cm	48.44	84.44	3555	35.55
CD (0.05)	1.79	5.13	1122	1122
CV(%)	9.99	7.51	11.45	10.11

Table 2. Interaction effect of treatments on per plant and per hectare rhizome yield and oil yield of *Kaempferia galanga* L. (Pooled mean of three years)

Treatment combination	Rhizome yield/plant (g)	Rhizome yield (kg/ha)	Oil yield (kg/ha)
15 th April 20 x 20 cm	45.00	9000	89.10
15 th April 30 x 30 cm	73.00	6489	64.24
15 th April 40 x 40 cm	77.67	3883	38.45
15 th April 50 x 50 cm	85.67	2741	27.14
30 th April 20 x 20 cm	46.67	9333	92.40
30 th April 30 x 30 cm	74.33	6607	65.41
30 th April 40 x 40 cm	78.33	3917	38.78
30 th April 50 x 50 cm	88.33	2827	27.98
15 th May 20 x 20 cm	40.00	8000	79.20
15 th May 30 x 30 cm	67.33	5985	59.25
15 th May 40 x 40 cm	71.00	3550	35.15
15 th May 50 x 50 cm	79.33	2539	25.13
CD (0.05)	8.89	728.68	7.21
CV (%)	8.84	9.82	7.72

Figure 1. Influence of spacing on rhizome yield of *Kaempferia galanga* L.




Plate 1. Growth of rhizome under varying spacing treatments

