

## Original Research Article

### Alkylphenols and Alkylphenol Ethoxylates in Freshwater Fish: Levels and Human Health Risk Estimates from River Benue, Central Nigeria

#### **ABSTRACT**

Alkylphenols (APs) and Alkylphenols ethoxylates (APEs), are ubiquitous estrogenic chemicals, widely distributed in fish due to their persistence and lipophilicity. This study determined levels of APs and APEs, and the associated human health risk in two fish species, 16 *Labeocoubie* (African carp) and 16 *Hydrocynus vittatus* (African Tiger fish) sampled quarterly from River Benue, central Nigeria between September 2019 and June 2020. Compounds were identified and quantified using Gas Chromatography with Mass Spectrometric detection (GC-MS). Levels of twenty APs ( $\sum_{20}$ APs) in *Labeo cubie* ranged from ND-16.0 ngg<sup>-1</sup> wet wt, with total and maximum values of 94.5 ngg<sup>-1</sup> wet wt, and 16.0 ngg<sup>-1</sup> wet wt respectively. Residues of four APEs ( $\sum_4$ APEs) ranged from ND-3.1 ngg<sup>-1</sup> wet wt, total of 5.0 ngg<sup>-1</sup> wet wt, and maximum of 3.1 ngg<sup>-1</sup> wet wt respectively. In *Hydrocynus vitatus*,  $\sum_{20}$ APs ranged from ND-27.8 ngg<sup>-1</sup> wet wt, total and maximum of 205 and 27.8 ngg<sup>-1</sup> wet wt, respectively. Estimated risk from ingestion of compounds in *Labeo cubie* and *Hydrocynus vitatus* by adult and children indicate no immediate potential harm as the hazard quotient (HQ) value was less than one (HQ < 1). However, children could suffer higher body burdens of these pollutants than adults if they are fed with these fish. This study provides baseline data on levels and estimated human health risk from ingestion of these two species of fish contaminated by APs and APEs to regulatory agents and future researchers.

**Keywords:** Alkylphenols, Alkylphenol Ethoxylates, Fish, River Benue, Central Nigeria

## **1. INTRODUCTION**

Alkylphenols (APs) are phenolic compounds obtained when one or more hydrogen atoms of the phenol are substituted by alkyl group, while ethoxylation of APs result into the nonionic surfactant known as Alkylphenol ethoxylates (APEs) [1]. APs and APEs are ubiquitous in the environment owing to their wide use in agriculture, household products and industry since 1950s [1]. These groups of chemicals have been described as pollutants of emerging concern, and are estrogenic because their human and wildlife exposure can cause adverse effects even at micro or nano concentrations [2].

Due to poor waste disposal practices in most developing countries such as Nigeria, wastes containing chemicals of various kinds are either discharged directly into surrounding water bodies or heaped on illegal waste dump sites in cities [3], constituting potential source for pollutants such as AP and APEs. About 60 % of the total global APs and APEs manufactured per year enter the world aquatic environment [4], where the pollutants degrade into various mono-, di-, tri-, etc. metabolites which could cause varying degrees of health effects to wildlife and humans [5].

Nigeria, the largest consumer of fish in Africa produces about 635,399 tons of fish annually, and imports about 560,000 tons in order to meet her local demand for fish [6]. Fish is widely consumed in Nigeria principally because of its high nutritive value [7]. As such, fish could constitute a major route of Nigerians exposure to pollutants such as APs and APEs.

To the best of our knowledge, this research is pilot, aimed at generating baseline data on the human health risk assessment through ingestion of APs and APEs in fish from river Benue by adult and children.


## **2. MATERIALS AND METHODS**

### **2.1 The Study Area**

River Benue is the second largest river in Nigeria. Its course originates from the fringes of the Cameroon mountains in the Republic of Cameroon, enters Nigeria

through Adamawa highlands, passes through Taraba plains, reaches Abinsi town, then the city of Makurdi before it empties its content into the River Niger at the confluence at Lokoja in Central Nigeria [8]. Abinsi, a fishing community is situated about 127 km east of the Makurdi Metropolitan area, also known for its fishing. These study areas experience rainy season (April to October) and dry season (November to March), and have mostly unconventional waste management practices and weak regulations. Generally, waste of various compositions is mostly dumped illegally in open areas, community-designated dumpsites or are discharged directly into River Benue, constituting potential source for many pollutants including APs and APEs [3, 9].

Four sampling sites (one at Abinsi town, and three at Makurdi) were purposively selected along River Benue as they act as potential flash points for pollutants (Fig. 1). Site 1 (Abinsi: 7° 76.060'N, 8° 72.610'E), receives effluents from town drains, fishing, laundry activities, sand dredging, boat station, riverbank commerce, and serves as domestic source of water. Site 2 (Mikap: 7° 43.649' N, 8° 35.302' E, has upstream from itPure Biotech Company Nigeria Limited, makers of bioethanol and Benue Brewery Nigeria Ltd). The site is prone to effluents from the industries upstream and has vibrant fishing, sand dredging and farming activities. Site 3 (Wurukum Abattoir: 7° 44.076' N, 8° 32.840' E) receives effluence from a major abattoir, city and rice mill drains, runoff from illegal sites for open-air incineration of old vehicle tires. Here too there is a lot of fishing and sand dredging, laundry, irrigation activities and serves as a source of water for rice processing and domestic use. Site 4 (Wadata: 7° 44.789'N, 8° 30.624'E). Illegal waste dump sites, runoff from Wadata open market, laundry, fishing, sand dredging, source of water for domestic and commerce, boat station, and riverbank commerce are the human activities impacting on this site. Direct waste discharge into the river is a common characteristic of all the sites.



**Fig. 1: Map of River Benue showing sampling sites in the study areas**

## 2.2 Sample Collection

Samples were collected according to Chokwe et al.[10] with slight modification. In brief, with the assistance of artisanal fishermen, samples of one bottom feeder, *Labeo cubie*, Rupell 1832, or African Carp, and one predator fish, *Hydrocynus vitatus*, Castelnau 1861, or Tiger fish were caught using nets in order to minimize skin damage; these were subsequently immobilized by the percussive stunning method [11], and wrapped individually in heavy-duty aluminium foil, labelled properly and placed in a chest packed with ice, and transported to the laboratories of Joseph Sarwuan Tarka University, Makurdi and kept frozen pending the time for sample preparation. The fish species were purposively selected as they are the commonly consumed species by the local population. A total of 32 fishes, 16 each of *Labeo cubie* and *Hydrocynus vitatus* were sampled quarterly between September 2019 and June 2020.

## 2.3 Sample Preparation, Extraction, Cleanup and Derivatization

The frozen fish were thawed enough to remove the foil wrapper and rinsed with tap water, then deionized water to remove any adhering debris, internal organs removed within 48 hours, and muscle tissue filleted, then homogenized on electric blender, and the homogenate stored frozen at -20 °C until the time of extraction.

The sample extraction procedure was a modification from Chokwe et al. [10]. Briefly, 5 g of the homogenized fish tissues were weighed and ground with anhydrous sodium

sulfate (1:4 w/w) to uniform paste using mortar and pestle, then extracted with 300 mL of hexane/acetone (4:1v/v) mixture in 500 mL round bottom flask at 55 °C for 12 hours, and extracts concentrated to 20 mL by solvent recovery. The resulting 20 mL extracts were placed in 250 mL separatory funnel, and 10mL of concentrated sulfuric acid added to remove lipids, and the mixture mechanically shaken for 3 minutes, then 7 mL acetone was added and shaken for 5 minutes for enhanced lipids removal and separation of layers. Subsequently, the acid layer was separated first followed by the hexane/acetone layer. The acid layer was washed with 25 mL hexane for further extraction of compounds. The hexane/acetone layer and recovered solvent were combined and washed with 20 mL of 40% (v/v) sulfuric acid to further remove residual lipids, and organic layer collected, and reduced to incipient dryness at 30 °C.

The residue was re-constituted with 2.5 mL methanol, diluted to 250 mL with MilliQ water, acidified to pH 3 with acetic acid, then subjected to solid phase extraction (SPE) process as follows.

The SPE column was plugged at the bottom with glass wool and 0.5 g of C<sub>18</sub> reversed phase silica gel placed on top of the wool, conditioned with 6 mL of 30% methanol in dichloromethane, followed by 6 mL dichloromethane, then solution extracted under gentle suction at 10 mL<sup>-1</sup>. The column was dried under gentle suction for 15 minutes, and compounds eluted with 6 mL of dichloromethane/hexane (4:1v/v) mixture. The eluate was subsequently concentrated to dryness under a gentle stream of nitrogen, then subjected to derivatization.

Reconstitution of APs and APEs residues was carried out using 0.1 mL hexane in a Pyrex tube, 40 µL of 0.1 M triethylamine (TEA) and 4 µL heptafluorobutyric anhydride (HFBA) added, and vortexed for 1 min, and the contents gently heated to 50°C for 30 minutes using GC to ensure derivatization of compounds. Contents were cooled, quenched with 0.3 mL of 5% aqueous solution of potassium carbonate (K<sub>2</sub>CO<sub>3</sub>), the organic phase drawn off. The aqueous phase was washed twice with 0.5 mL of hexane for compound recovery, and the organic phase extracts were concentrated to

100 µL. Chrysene was added as internal standards, volume made up to 200 µL, and 1 µL of the mixture injected into GC-MS.

#### **2.4 Instrumental Analysis**

The derivatized extracts were analyzed using an Agilent 6890 GC equipped with 5975 mass selective detector (MSD) comprising of capillary column: Restek RTx1614, film thickness of 0.10 µm, 15 m x 0.25 mm I.D. and fitted to an Agilent autosampler A673. Using Helium as carrier gas, the GC/MS conditions were: linear velocity of 40 cm/s, 150 °C ion source, maximum injector temperature of 280 °C, and 300°C transfer line temperature. The temperature programme was: a starting temperature of 50 °C, heated to 120 °C by a temperature ramp of 7.5 °C/min, then 275 °C by a temperature ramp of 15 °C/min, and finally heated to 300 °C (held for 2 min) by a temperature ramp of 25 °C/min.

#### **2.5 Quality Assurance and Statistical Analysis**

Fish were caught using nets to minimize skin damage, and fish spines carefully sheared to minimize punctures during the aluminium foil packaging. Samples were transported in chest packed with ice to avoid decomposition. Edible parts (muscle tissue) were filleted within 48 hours to avoid contamination by ruptured internal organs. Monitoring chromatographic conditions to check variations during analysis, then running blanks at interval of five samples, and duplicate sample analysis were the other quality assurance measures adopted. All data were analyzed using Microsoft Windows Excel 2019 version.

#### **2.6 Human Health Risk Estimates**

In this study, the risk assessment tool recommended by USEPA 1989 [12], was adopted to estimate the non-carcinogenic hazard quotient (HQ) due to adult and children exposure through ingestion of fish contaminated by NP, NPE, OP, OPE and 4t-BP as follows:

$$HQ = ADD/RfD \quad (1)$$

$$ADD = C_i \times IR/BW \quad (2)$$

Where, ADD is the average daily intake of a chemical ( $\mu\text{kg}^{-1} \text{BWday}^{-1}$ ) and RfD is the daily intake reference dose ( $\mu\text{kg}^{-1} \text{BWday}^{-1}$ ). The tolerable daily intake (TDI) for NP was  $5000 \text{ ngkg}^{-1} \text{BWday}^{-1}$ , and  $13000 \text{ ngkg}^{-1} \text{BWday}^{-1}$  for NPE as proposed by the Danish Institute of Safety and Toxicology were adopted as their RfD [13]. The United States Environmental Protection Agency (USEPA) sets a RfD for phenol at  $300\ 000 \text{ ngkg}^{-1} \text{BWday}^{-1}$ [14], and this RfD value was employed for estimating the HQ of OP, OPE and 4t-BP during this study, since their RfD have not yet been reported in literature.  $C_i$  is the concentration of APs and/or APEs in fish ( $\mu\text{gkg}^{-1}$  wet wt). Ingestion rate (IR) is the amount of fish ingestion per day ( $\text{kg person}^{-1}\text{day}^{-1}$ ) and was calculated based on per capital consumption of  $0.021 \text{ kg adult}^{-1}\text{day}^{-1}$  for adult Nigerians, while one-third IR of the adult is consumed by children (i.e.  $0.007 \text{ kg child}^{-1}\text{day}^{-1}$ ). BW is the body weight (kg) of adult (70 kg), and children (16 kg) respectively. The above parameters were obtained from [15], [16],[17] and [12]. An  $\text{HQ} > 1$  indicates that there may be concern for potential human health effects, and vice versa.

### 3. RESULTS AND DISCUSSION

#### 3.1 Levels of APs and APEs in *Labeo cubie* from River Benue

Sum of twenty APs ( $\sum_{20}\text{APs}$ ) detected in *Labeo cubie* samples from River Benue as obtained from Table 1, ranged from  $14.6\text{-}28.0 \text{ ngg}^{-1}$  wet wt, with total and maximum values of  $94.5 \text{ ngg}^{-1}$  wet wt, and  $16.0 \text{ ngg}^{-1}$  wet wt respectively. Meanwhile, the range, total and maximum levels of sum of nine short chain APs ( $\sum_9\text{SCAPs}$ ) were ND- $4.16$ ,  $11.4$  and  $4.16 \text{ ngg}^{-1}$  wet wt, respectively. Similarly, sum of eleven long chain APs documented were in range of ND- $12.3 \text{ ngg}^{-1}$  wet wt, total and maximum of  $83.1 \text{ ngg}^{-1}$  wet wt, and  $16.00 \text{ ngg}^{-1}$  wet wt, respectively. Sum of four APEs ( $\sum_4\text{APEs}$ ) were observed to range from ND- $3.12 \text{ ngg}^{-1}$  wet wt, total of  $5.01 \text{ ngg}^{-1}$  wet wt, and maximum of  $3.12 \text{ ngg}^{-1}$  wet wt respectively (Table 1).  $\sum_{11}\text{LCAPs}$  contributed about 88 % abundance of total mean concentration over about 12 % abundance donated by  $\sum_9\text{SCAPs}$ (Table 1). Concentrations of  $\sum_9\text{SCAPs}$  were highest in site 2

followed by site 3, then site 4, but below the limit of detection of the instrument used for this work in site 1, while sum of three ethyl phenol ( $\Sigma_3\text{EP}$ ) was most abundant metabolites(8 %) among the short chain APs detected, but levels of sum of three methyl phenol ( $\Sigma_3\text{MP}$ ) and sum of three propyl phenol ( $\Sigma_3\text{PP}$ ) were less than limit of detection of our analytical instrument (Table 1). Again, site 3 recorded the highest level of sum of eleven long chain APs ( $\Sigma_{11}\text{LCAPs}$ ) followed by site 4, then sites 1 and 2, respectively. 2-octyl phenol (2-OP) was most abundant (63 %) among the LCAPs detected in *Labeo cubie*, followed by 4-octylphenol (4-OP) with 50%, then, 4-nonylphenol (4-NP) (19%), 4-hexylphenol (4-HxP) and 4-heptylphenol (13% each), 4-butylphenol (4-BP) and 2-hexylphenol (2-HxP) (6% each), while 4-tert butylphenol (4t-BP), 2-pentylphenol (2-PnP), 4-pentylphenol (4-PnP), 4-tert octylphenol (4t-OP) recorded the detection frequency of zero percent (0%) respectively (Table 1)

Exception of site 1, concentrations of  $\Sigma_4\text{APEs}$  in tissues of *Labeo cubie* obtained in all other sites were below the limit of detection of our analytical instrument, and NPE1 was the dominant homologue among these group of compounds. Generally, the long chain AP homologues were most frequently detected in the *Labeo cubie* tissues, while the SCAPs homologues were least detected (Table 1). Variation in the sources of these pollutants at the studied sites might account for the differences in concentrations of these pollutants at the studied sites. Although fish are mobile, therefore, the effect of site-specific anthropogenic activities on their tissue accumulation of pollutants might be less effective [18], nevertheless, this factor becomes effective when considering the broader habitat of the fish. The impact of environmental concentration of APs and APEs on the body burden of the chemicals in fish was previously shared by [12].

**Table 1:** Concentrations of APs and APEs ( $\text{ngg}^{-1}$  wet wt) in *Labeo cubie* from River Benue

| Analyte | Site 1 |     |     | Site 2 |     | Site 3 |     | Site 4 |  | FD |
|---------|--------|-----|-----|--------|-----|--------|-----|--------|--|----|
|         | Sum    | Max | Sum | Max    | Sum | Max    | Sum | Max    |  |    |
| <hr/>   |        |     |     |        |     |        |     |        |  |    |


|                     |             |             |             |             |             |             |             |             |    |
|---------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----|
| 2-MP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| 3-MP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| 4-MP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_3$ MP       | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | 0  |
| 2-EP                | ND          | ND          | 2.82        | 2.82        | ND          | ND          | ND          | ND          | 6  |
| 3-EP                | ND          | ND          | ND          | ND          | 4.16        | 4.16        | 1.14        | 1.14        | 13 |
| 4-EP                | ND          | ND          | 3.30        | 3.3         | ND          | ND          | ND          | ND          | 6  |
| $\Sigma_3$ EP       | <b>ND</b>   | <b>ND</b>   | <b>6.12</b> | <b>5.12</b> | <b>4.16</b> | <b>4.16</b> | <b>1.14</b> | <b>1.14</b> | 8  |
| 2-PP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| 3-PP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| 4-PP                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_3$ PP       | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | 0  |
| $\Sigma_9$ SCAPs    | <b>ND</b>   | <b>ND</b>   | <b>6.12</b> | <b>6.12</b> | <b>4.16</b> | <b>4.16</b> | <b>1.14</b> | <b>1.14</b> | 3  |
| 4-BP                | ND          | ND          | 0.73        | 0.73        | ND          | ND          | ND          | ND          | 6  |
| 4t-BP               | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_2$ BP       | <b>ND</b>   | <b>ND</b>   | <b>0.73</b> | <b>0.73</b> | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | 3  |
| 2-PnP               | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| 4-PnP               | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_2$ PnP      | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | <b>ND</b>   | 0  |
| 2-HxP               | 3.84        | 3.84        | ND          | ND          | ND          | ND          | ND          | ND          | 6  |
| 4-HxP               | ND          | ND          | 2.89        | 2.89        | 3.13        | 3.13        | ND          | ND          | 13 |
| $\Sigma_2$ HxP      | <b>3.84</b> | <b>3.84</b> | <b>2.89</b> | <b>2.89</b> | <b>3.13</b> | <b>3.13</b> | <b>ND</b>   | <b>ND</b>   | 6  |
| 4-HP                | ND          | ND          | ND          | ND          | 1.13        | 1.13        | 3.13        | 3.13        | 13 |
| 2-OP                | 9.71        | 3.82        | 2.37        | 2.37        | 9.29        | 4.5         | 7.71        | 3.62        | 63 |
| 4-OP                | ND          | ND          | 2.89        | 2.89        | 15.7        | 6.19        | 16.0        | 5.93        | 50 |
| 4t-OP               | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_3$ OP       | <b>9.71</b> | <b>9.71</b> | <b>5.26</b> | <b>2.89</b> | <b>25.2</b> | <b>15.7</b> | <b>23.7</b> | <b>16.0</b> | 38 |
| 4-NP                | 1.09        | 1.09        | 3.37        | 2.7         | ND          | ND          | ND          | ND          | 19 |
| $\Sigma_{11}$ LCAPs | <b>14.6</b> | <b>9.71</b> | <b>12.3</b> | <b>3.37</b> | <b>29.3</b> | <b>15.7</b> | <b>26.9</b> | <b>16.0</b> | 15 |
| $\Sigma_{20}$ APs   | <b>14.6</b> | <b>9.71</b> | <b>18.4</b> | <b>3.37</b> | <b>33.5</b> | <b>15.7</b> | <b>28.0</b> | <b>16.0</b> | 10 |
| OPE1                | 0.27        | 0.27        | ND          | ND          | 1.61        | 1.61        | ND          | ND          | 13 |
| OPE2                | ND          | ND          | ND          | ND          | ND          | ND          | ND          | ND          | 0  |
| $\Sigma_2$ OPE      | <b>0.27</b> | <b>0.27</b> | <b>ND</b>   | <b>ND</b>   | <b>1.62</b> | <b>1.61</b> | <b>ND</b>   | <b>ND</b>   | 6  |
| NPE1                | 3.12        | 3.12        | ND          | ND          | ND          | ND          | ND          | ND          | 6  |

|               |      |      |    |    |      |      |    |    |   |
|---------------|------|------|----|----|------|------|----|----|---|
| NPE2          | ND   | ND   | ND | ND | ND   | ND   | ND | ND | 0 |
| $\sum_2$ NPE  | 3.12 | 3.12 | ND | ND | ND   | ND   | ND | ND | 3 |
| $\sum_4$ APEs | 3.39 | 3.12 | ND | ND | 1.62 | 1.62 | ND | ND | 5 |

$\sum_3$ MP = 2-MP + 3-MP + 4-MP.  $\sum_3$ PP = 2-PP + 3-PP + 4-PP.  $\sum_9$ SCAPs =  $\sum_3$ MP +  $\sum_3$ EP +  $\sum_3$ PP.  $\sum_2$ BP = 4-BP + 4t-BP.  
 $\sum_2$ PnP = 2-PnP + 4-PnP.  $\sum_2$ HxP = 2-HxP + 4-HxP.  $\sum_3$ OPs = 2-OP + 4-OP + 4t-OP.  $\sum_{11}$ LCAP =  $\sum_2$ BP +  $\sum_2$ PnP + 4-HP +  $\sum_2$ HxP +  $\sum_3$ OPs + 4-NP.  $\sum_{20}$ APs =  $\sum_9$ SCAPs +  $\sum_{11}$ LCAP + 4-HP + 4-NP.  $\sum_2$ OPEs = OPE1 + OPE2.  $\sum_2$ NPEs + NEP1 + NPE2. ND = Not Detected


When compared with similar other studies, levels of sum of the compounds recorded in *Labeo cubie* from River Benue were in range with the reports of [19] and [20], but lower than results reported by [10], and [21], respectively. The sum of mono-, and di-nonylphenol ethoxylate ( $\sum_2$ NPE) obtained from *Labeo cubie* were dominant over that of mono-, and di-octylphenol ethoxylate ( $\sum_2$ OPE), which is consistent with the reports of [10] and [22].

Distribution of compounds in *Labeo cubie* per chain length per site was also considered (Fig. 2). Accumulation of  $\sum_{11}$ LCAPs in *Labeo cubie* was consistently highest in all the sites followed by  $\sum_4$ APEs in Abinsi, then  $\sum_9$ SCAPs in Abattoir and Wadata, respectively, which suggests variation in degradation rates of APEs into APs, and lipophilicity of these pollutants. The impact of degradation on the tissue accumulation of APs and APEs was previously reported in literature [21].



**Fig.2: Accumulation of  $\sum_9$ SCAPs,  $\sum_{11}$ LCAPs and  $\sum_4$ APEs in *Labeo cubie* from River Benue across the sites**

Accumulation of compounds in *Labeo cubie* (Fig. 3) was predominant in the 3<sup>rd</sup> quarter (March) in all the sites. APs and APEs are lipophilic, as such, their bioavailability might be highest in March, the month of the year in Nigeria when fishes are gorged with fats[18]. Fish are mobile, this may account for the observed irregular or diffused pattern of accumulation of the chemicals in *Labeo cubie* across the sites (Fig. 3).



**Fig. 3: Quarterly variation of the concentrations of APs, APEs and metabolites taken together in *Labeo cubie* from River Benue across sites**

### 3.2 Levels of APs and APEs in *Hydrocynus vitatus* from River Benue

The levels of  $\sum_{20}$ APs detected in *Hydrocynus vitatus* (Table 2) ranged from 40.31-68.60 ngg<sup>-1</sup> wet wt, with total and maximum of 205.32 and 68.60 ngg<sup>-1</sup> wet wt respectively. Similarly,  $\sum_9$ SCAPs ranged from 0.61-35.27 ngg<sup>-1</sup> wet wt, total and maximum concentrations of 65.81 ngg<sup>-1</sup> wet wt and 35.27 ngg<sup>-1</sup> wet wt, and  $\sum_3$ MP were most frequently detected (10%) homologue among the short chain APs, while concentrations of  $\sum_3$ EP were below the limit of detection of the instrument used for this study.

**Table 2:** Concentrations of APs and APEs (ngg<sup>-1</sup> wet wt) in *Hydrocynus vitatus* from River Benue


| Analyte     | Site 1    |           | Site 2       |              | Site 3      |             | Site 4       |              | FD |
|-------------|-----------|-----------|--------------|--------------|-------------|-------------|--------------|--------------|----|
|             | Sum       | Max       | Sum          | Max          | Sum         | Max         | Sum          | Max          |    |
| 2-MP        | ND        | ND        | 1.35         | 1.35         | 4.25        | 4.25        | ND           | ND           | 13 |
| 3-MP        | ND        | ND        | 20.52        | 20.52        | 3.80        | 3.80        | ND           | ND           | 13 |
| 4-MP        | ND        | ND        | ND           | ND           | ND          | ND          | 25.04        | 25.04        | 6  |
| $\sum_3$ MP | <b>ND</b> | <b>ND</b> | <b>21.87</b> | <b>20.52</b> | <b>8.06</b> | <b>4.25</b> | <b>25.04</b> | <b>25.04</b> | 10 |

|                    |              |              |              |              |              |              |              |              |    |
|--------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|----|
| 2-EP               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| 3-EP               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| 4-EP               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| $\Sigma_3$ EP      | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | 0  |
| 2-PP               | ND           | ND           | ND           | ND           | ND           | ND           | 4.11         | 4.11         | 6  |
| 3-PP               | ND           | ND           | ND           | ND           | ND           | ND           | 6.12         | 6.12         | 6  |
| 4-PP               | 0.61         | 0.61         | ND           | ND           | ND           | ND           | ND           | ND           | 6  |
| $\Sigma_3$ PP      | <b>0.61</b>  | <b>0.61</b>  | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>10.23</b> | <b>6.12</b>  | 6  |
| $\Sigma_9$ SCAP    | <b>0.61</b>  | <b>0.61</b>  | <b>21.87</b> | <b>20.52</b> | <b>8.06</b>  | <b>4.25</b>  | <b>35.27</b> | <b>25.04</b> | 6  |
| 4-BP               | ND           | ND           | 5.85         | 4.24         | ND           | ND           | 11.15        | 5.61         | 25 |
| 4t-BP              | ND           | ND           | ND           | ND           | ND           | ND           | 5.11         | 5.116        | 6  |
| $\Sigma_2$ BP      | <b>ND</b>    | <b>ND</b>    | <b>5.83</b>  | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>16.26</b> | <b>11.15</b> | 16 |
| 2-PnP              | ND           | ND           | ND           | ND           | 4.39         | 4.39         | ND           | ND           | 6  |
| 4-PnP              | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| $\Sigma_2$ PnP     | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>4.39</b>  | <b>4.39</b>  | <b>ND</b>    | <b>ND</b>    | 3  |
| 2-HxP              | ND           | ND           | 2.11         | 2.11         | ND           | ND           | 0.97         | 0.97         | 13 |
| 4-HxP              | ND           | ND           | 6.80         | 6.80         | ND           | ND           | ND           | ND           | 6  |
| $\Sigma_2$ HxP     | <b>ND</b>    | <b>ND</b>    | <b>8.91</b>  | <b>6.80</b>  | <b>ND</b>    | <b>ND</b>    | <b>0.97</b>  | <b>0.97</b>  | 9  |
| 4-HP               | 3.28         | 3.28         | 4.87         | 4.87         | ND           | ND           | ND           | ND           | 13 |
| 2-OP               | 27.77        | 24.73        | ND           | ND           | 22.94        | 22.01        | 2.54         | 2.54         | 31 |
| 4-OP               | 8.22         | 4.26         | 3.36         | 3.36         | 4.92         | 2.44         | 2.48         | 2.48         | 44 |
| 4t-OP              | 3.90         | 3.90         | ND           | ND           | ND           | ND           | ND           | ND           | 6  |
| $\Sigma_3$ OP      | <b>39.89</b> | <b>27.77</b> | <b>3.36</b>  | <b>3.36</b>  | <b>27.86</b> | <b>22.94</b> | <b>5.02</b>  | <b>2.54</b>  | 29 |
| 4-NP               | ND           | ND           | 7.79         | 3.94         | ND           | ND           | 11.08        | 6.71         | 25 |
| $\Sigma_{11}$ LCAP | <b>43.17</b> | <b>27.77</b> | <b>30.76</b> | <b>12.35</b> | <b>32.25</b> | <b>27.77</b> | <b>33.33</b> | <b>11.15</b> | 16 |
| $\Sigma_{20}$ APs  | <b>43.78</b> | <b>27.77</b> | <b>52.63</b> | <b>21.87</b> | <b>40.31</b> | <b>27.77</b> | <b>68.60</b> | <b>25.04</b> | 12 |
| OPE1               | ND           | ND           | 3.06         | 3.06         | ND           | ND           | ND           | ND           | 6  |
| OPE2               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| $\Sigma_2$ OPE     | <b>ND</b>    | <b>ND</b>    | <b>3.06</b>  | <b>3.06</b>  | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | 3  |
| NPE1               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| NPE2               | ND           | ND           | ND           | ND           | ND           | ND           | ND           | ND           | 0  |
| $\Sigma_2$ NPE     | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | 0  |
| $\Sigma_4$ APE2    | ND           | ND           | <b>3.06</b>  | <b>3.06</b>  | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | <b>ND</b>    | 2  |

Again, 4-OP was the most frequently detected (44 %) among the LCAPs, whereas, 4-pentyl phenol (4-PnP) was least detected (0 %).  $\sum_4$ APEs ranged from ND-3.06 ngg<sup>-1</sup> wet wt, total and maximum of 3.06 ngg<sup>-1</sup> wet wt each (Table 2). Concentration of  $\sum_2$ NPE were below the limit of detection of our instrument, while  $\sum_2$ OPE were most frequently detected (3 %), and most abundant homologue recorded among the APEs in the *Hydrocynus vitatus*. Generally, variation in the detection frequencies of compounds in the tissues of *Hydrocynus vitatus* suggests point and non-point source emissions of these pollutants around the sites studied.  $\sum_9$ SCAPs contributed about 37 % of the sum of mean concentration of  $\sum_{20}$ APs while  $\sum_{11}$ LCAPs homologues represented about 63 %. APs are lipophilic, and this property increases with chain length [23], this might account for the differences in percentage absorption of these chemicals to the fish tissues


In comparison with similar other literatures, [19] and [20] reported levels of APs and APEs in similar range with our study, while [10], and [12] published higher values of these pollutants. Long chain APEs can degrade into mono and di-ethoxylate metabolites which tend to be more persistent, more lipophilic and bioaccumulative step-wise in the trophic [24].

Trend of chemicals accumulated by *Hydrocynus vitatus* at the various sites followed the pattern:  $\sum_{11}$ LCAP >  $\sum_9$ SCAP >  $\sum_4$ APE (Fig.4). Variation in degradation rate and/or fat solubility of these chemicals might be responsible for this trend [21].



**Fig. 4: Accumulation of  $\sum_9$ SCAPs,  $\sum_{11}$ LCAPs and  $\sum_4$ APEs in *Hydrocynus vitatus* from River Benue across the sites**

APs and APEs are lipophilic, and as such, are most bioavailable around March, the month of the year in Nigeria when fishes are rich in lipids [18]. This explains why the highest concentrations of chemicals reported in *Hydrocynus vitatus* in this research were observed in the 3<sup>rd</sup> quarter (March) in all the sites (Fig. 5).



**Fig. 5: Quarterly Trend of concentrations of APs, APEs and metabolites taken together in *Hydrocynus vitatus* from River Benue across sites**

Accumulation of compounds across the sites followed a diffused pattern, which might be the consequence of mobility of fish which makes site-specific influence of these pollutants on fish difficult to determine.

### 3.3 Accumulation of APs, APEs and their Metabolites by *Hydrocynus vitatus* Versus *Labeo cubie* from River Benue


It could be observed from Tables 1 and 2 that LCAPs are the dominant homologues among the pollutants detected during this study, followed by SCAP, then APEs.  $\sum_3$ OP was observed to be most concentrated among the LCAP with the total concentration of 63.87 ng g<sup>-1</sup> wet wt, and detection frequency of 38 % for *Labeo cubie*, while the total concentration of 76.13 ng g<sup>-1</sup> wet wt, and detection frequency of 29 % were recorded for *Hydrocynus vitatus*. 2-OP monopolized dominance with total concentration of 29.08 ng g<sup>-1</sup> wet wt for *Labeo cubie*, and 53.25 ng g<sup>-1</sup> wet wt for *Hydrocynus vitatus*. Mixed ratio of dominance was observed among the SCAP.  $\sum_3$ EP recorded the highest total concentration (11.42 ng g<sup>-1</sup> wet wt) in *Labeo cubie*, and 3-EP topped the total concentration (5.3 ng g<sup>-1</sup> wet wt), and frequency of detection (13 %)

among the ethylphenol metabolites in this species of fish.  $\Sigma_3$ MP had the highest total concentration (54.97 ngg<sup>-1</sup> wet wt) and detection frequency (13 %) among the SCAPs for *Hydrocynus vitatus*, while 4-MP dominated total metabolite concentration (25.04 ngg<sup>-1</sup> wet wt) of this AP homologue in this species of fish.

APEs showed inverse abundance of total metabolite concentration of pollutants among the two fish species: NPE > OPE in *Labeo cubie*, while OPE > NPE in *Hydrocynus vitatus* respectively (Tables 1 and 2), and this suggests variation in sources and/or tissue metabolism of these compounds by the two fish species. Chokwe et al. [10] reported higher abundance ratios of di-NPE and mono-NPE (23.52 ng g<sup>-1</sup> versus 12 ng g<sup>-1</sup>lipids) in fish, the remaining 10.6 ngg<sup>-1</sup> lipids was contributed by sum of mono- and di- OPEs, and this report was consistent with [22] and [25].

Result in Fig.6 indicates that, the predatory *Hydrocynusvittus* consistently accumulated higher levels of pollutants than the bottom feeder *Labeo cubie* across sites.

This trend of accumulation is consistent with the fact that the feeding habit of fish contributes to the bioaccumulation of pollutants in its tissues[12, 10], although, fish physiology, metabolism and environmental levels of pollutants are factors to consider as well. Trend of accumulation of compounds in the two species of fish was statistically significant across sites except for abattoir, and this implies variation in of level of exposure fish to these pollutants.



**Fig.6: Comparison of the accumulation of total APs, APEs and their metabolites taken together in *Labeo cubie* and *Hydrocynus vitatus* from River Benue**

### **3.4 Human Health Risk Assessment of APs and APEs in *Labeo cubie* and *Hydrocynus vitatus* from River Benue**

The HQ for ingestion of APs and APEs in *Labeo cubie* by adults and children (Table 3) was less than one ( $HQ < 1$ ), which suggests that consumption of this species of fish by the two groups posed no significant health risk. Lee *et al.* [21] reported HQ for the ingestion of NP in fish by adults and children from Tamsui river to be less than one as well.

**Table 3:** Hazard Quotient for Ingestion of APs and APEs ( $\text{ngg}^{-1}$  wet wt) in Fish (*Labeo cubie*) from River Benue by Adult and Children

| Analyte | Abinsi   |          | Mikap    |          | Abattoir |          | Wadata |          | Adult    |          | Children |          |
|---------|----------|----------|----------|----------|----------|----------|--------|----------|----------|----------|----------|----------|
|         | Adult    | Children | Adult    | Children | Adult    | Children | Adult  | Children | Sum      | Mean     | Sum      | Mean     |
| 4t-BP   | NA       | NA       | NA       | NA       | NA       | NA       | NA     | NA       | NA       | NA       | NA       | NA       |
| 4t-OP   | NA       | NA       | NA       | NA       | NA       | NA       | NA     | NA       | NA       | NA       | NA       | NA       |
| 4-NP    | 3.76E-06 | 5.36E-06 | 1.17E-05 | 1.67E-05 | NA       | NA       | NA     | NA       | 1.55E-05 | 3.87E-06 | 2.21E-05 | 5.52E-06 |

\* NA = Not applicable as the contaminants were not detected


The risk estimates of exposure to 4t-BP and 4t-OP through ingestion of *Labeo cubie* by the two groups was effectively not applicable (NA), as concentrations of these chemicals in this fish tissues were less than the limit of quantitation of our instrument. Nevertheless, HQ of 4-NP across sites was found to be less than the threshold value of one for the two groups, and risk was higher for adults in Abinsi and Mikap, respectively, while children recorded higher sum and mean HQ than adults (Table 3). The body burden of 4-NP for the two groups is approximately close, which might account for variation of the inverse sum and mean risk observed.

Sum and mean HQ for consumption of APs and APEs in *Hydrocynus vitatus* was consistent for the two groups and decreased in the sequence: 4-NP > 4t-BP > 4t-OP, with children having the highest body burden of the pollutants in Abinsi, Mikap and Wadata (Table 4).  $\Sigma$ HQ recorded in *Hydrocynus vitatus* ranged from ND-1.70E-04 for adult and ND-2.50E-04 for children, while their total and sum of mean as obtained from Table 4 were 7.67E-05 and 8.69E-05, respectively. Children have lower body weight (16 kg), and their bodies are more sensitive to pollutants than adults (body weight = 70 kg), which explains why children were observed to have suffered more body burden through ingestion of APs and APEs in *Hydrocynus vitatus* than adults.

Although the risk values for consumption of *Hydrocynus vitatus* contaminated by APs and APEs by the two groups suggests no immediate potential harm (HQ <1), continuous exposure to these chemicals might eventually become harmful since APs and APEs are persistent and bioaccumulative, hence the need for routine monitoring of the chemicals in the River Benue.

**Table 4:** Hazard Quotient ( $\text{ngg}^{-1}$  wet wt) for Ingestion of *Hydrocynus vitatus* from River Benue by Adult and Children

| Analyte     | Abinsi   |          | Mikap    |          | Abattoir |          | Wadata   |          | Adult    |          | Children |          |
|-------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
|             | Adult    | Children | Adult    | Children | Adult    | Children | Adult    | Children | Sum      | Mean     | Sum      | Mean     |
| 4t-BP       | NA       | NA       | NA       | NA       | NA       | NA       | 1.86E-06 | 2.70E-06 | 1.86E-06 | 4.65E-07 | 2.7E-06  | 6.75E-07 |
| 4t-OP       | 1.41E-06 | 2.05E-06 | NA       | NA       | NA       | NA       | NA       | NA       | 1.41E-06 | 3.53E-07 | 2.05E-06 | 5.13E-07 |
| 4-NP        | NA       | NA       | 1.69E-04 | 2.46E-04 | NA       | NA       | 4.00E-06 | 5.84E-06 | 1.73E-04 | 4.33E-05 | 2.52E-04 | 6.30E-05 |
| $\Sigma$ HQ | 1.41E-06 | 2.05E-06 | 1.70E-04 | 2.50E-04 | NA       | NA       | 5.86E-06 | 8.54E-06 | 1.80E-04 | 4.41E-05 | 2.60E-04 | 6.41E-05 |



**Fig. 7: Hazard Quotient (HQ) for Ingestion of APs and APEs Taken Together in *Labeo cubie* and *Hydrocynus vitatus* Fish from River Benue**

Comparative trend of exposure through ingestion of the two fish species (*Labeo cubie* and *Hydrocynus vitatus*) by adult and children was equally evaluated as shown in Fig.7. Exposure to APs and APEs through ingestion of *Hydrocynus vitatus* by the two groups was consistently higher than *Labeo cubie*, except that the maximum for adults was higher in *Labeo cubie*. It is worth to note that *Hydrocynus vitatus* is predatory, while *Labeo cubie* is bottom feeder, hence, feeding habit might account for the higher tissue levels of the pollutants recorded in *Hydrocynus vitatus* than *Labeo cubie*. Bioaccumulation of APs and APEs in fish was previously reported in literature [12,10, 26]. Nevertheless, overlapping of the maximum levels of the pollutants in adult through ingestion of pollutants in the bottom feeding *Labeo cubie* over the predatory *Hydrocynus vitatus* suggests that factors other than feeding habit might favour accumulation of the pollutants in fish tissues as well. Lee et al. [12] reported that factors such as environmental concentration of chemicals (exposure level), physiology and burrowing behavior of fish might also favour accumulation of chemicals in fish tissues, which demands for further research in the future.

#### 4. CONCLUSION

Baseline data on levels and human health risk estimates on exposure to APs and APEs through consumption of *Labeo cubie* and *Hydrocynus vitatus* from Rive Benue was generated from this study. Results indicated that the two species of fish were minimally contaminated by the studied pollutants, and as such, adults and children were not likely to be affected by consumption of these fishes in their diets, since the HQ was less than one for the two groups. Nevertheless, the high level of local consumption of fish [6], coupled with the lipophilicity of the chemicals is a concern because these chemicals can bioaccumulate and likely biomagnify in food chain, and eventually cause harm to ecology, then humans, hence, the need for routine monitoring of these compounds in the area studied. This study provides baseline data for regulatory authorities. Modification of the lipid extraction and SPE cleanup methods reported by [10] provides an opportunity for researchers working in less equipped laboratories to use our own approach.

**Acknowledgement** The authors acknowledge Mrs. Martha Sewuese Tongu of Tse Pool Layout, Makurdi for her assistance during sample treatment, Dr. Akange Edward of the Department of Fisheries and Aquaculture, Joseph Sarwuan Tarka University, Makurdi for identifying the fish species, and Mr. Emmanuel Ukpoko of the Department of Chemistry, Joseph Sarwuan Tarka University, Makurdi for his assistance during sampling, and technical support in the laboratory.

**Author Contributions** Geoffrey Aernyi Wase in collaboration with Sylvester Mbaavaga Tongu, collected field samples, analyzed and interpreted data; G. A. Wase also drafted the manuscript for this publication. Rufus Sha'Ato, Ishaq Shaibu Eneji and Raymond Ahule Wuana, conceived and designed the experiments, contributed reagents, materials, analysis tools, and reviewed the draft manuscript of the paper.

**Declaration of Competing Interest:** The authors declare no competing funding or other interests that could influence the work reported in this paper.

**Data Availability** Data for this work will be made available on official request.

**Ethical Declaration/consent** Not applicable.

## REFERENCES

1. Anne, P., Nadia, M., Coline, D., Sophie, G., Corina, B., Céline L., Giangiaco, T., Peter, W. And Gre´gorio, C. (2017). Alkylphenol and Alkylphenol Polyethoxylates in Water and Wastewater: A Review of Options for their Elimination. *Arabian Journal of Chemistry*, 10: S3749-S3773.
2. Hong, Y., Feng, C., Jin, X., Xie, H., Liu, N., Bai, Y., Wu, F. and Raimondo, S. (2022). A QSAR–ICE–SSD Model Prediction of the PNECs for Alkylphenol Substances and Application in Ecological Risk Assessment for Rivers of a Megacity. *Environment International*, 167: 107367.
3. Sha’Ato, R., Aboho, S. Y., Oketunde, F. O., Eneji, I. S., Unazi, G. and Agwa, S. (2007). Survey of Solid waste Generation and Composition in a Rapidly Growing Urban Area in Central Nigeria, *Waste Manag.* 27: 352–358.
4. Barse, A. V., Chakrabarti, T., Ghosh, T. K., Pal, A. K. and Jadhao, S. B. (2006). One-tenth dose of LC<sub>50</sub> of 4-Tert-Butylphenol Causes Endocrine Disruption and Metabolic Changes in *Cyprinus carpio*. *Pesticides Biochemistry and Physiology*, 86: 172-179.
5. Xie, W., Zhao, J., Zhang, Q., Ye, C., Zheng, G., Shan, Q., Li, L. and Shao, X. (2020). Occurrence, distribution and bioaccumulation of alkylphenols in the Pearl River networks, South China. *Ecological Indicators*, 110: 105847.
6. Ucherwuhe, S. I., Ashiko, F. T.G and Akange, E. T. (2018). Socio-Economic Analysis of Fishing Practices of the Fisherfolks of Abinsi Settlement, Benue State, Nigeria. *International Journal of Research*, 5: 17.
7. Adewale, A., Adegbola, P. I., Owoade, A. O. and Aborisade, A. B. (2022). Fish as a Bioindicator of Polycyclic Aromatic Hydrocarbon Pollution in Aquatic Ecosystem of Ogun and Eleyele Rivers, Nigeria, and Risk Assessment for Consumer’s Health. *Journal of Hazardous Materials Advances*, 7: 100096.

8. Tongu. S. M., Sha'Ato, R., Wase, G. A., Okonkwo, J. O., Vesuwe, R. N. (2023). Organochlorine Pesticides and Polychlorinated Biphenyls in City Drains in Makurdi, Central Nigeria: Seasonal Variations, Source Apportionment and Risk Assessment. *Heliyon*, 9: e14324.
9. Wuana, R. A., Aide, P. A. and Asegh, I. N. (2012). Seasonal Variation in Bioavailability of Some Toxic Metals – Waste Dumps of Makurdi North Central Nigeria, *Journal of Bio-Diversity and Environmental Science*, 2 (11): 7–17.
10. Chokwe, B., Okonkwo, O. J., Sibali, L. L., Krüger, E., Du Preez, H., Hariram, R. and Ncube, E. J. (2015b). A Simplified Analytical Procedure for Simultaneous Determination of Alkylphenol Ethoxylates and Brominated Flame Retardants in Fish Tissue Samples from Vaal River, South Africa. *American Journal of Analytical Chemistry*, 6: 422–428.
11. Roth, B., linde, E. and Robb. D. H. F. (2007). Percussive Stunning of Atlantic Salmon (*Salmo Salar*) and the Relation Between Force and Stunning. *Aquacultural Engineering*, 36: 192–197,
12. Lee, C-C., Jiang, L.Y., Kuo, Y.L, Chen, C.Y., Hsieh, C.Y., Hung, C.F. and Tien, C.J. (2015). Characteristics of Nonylphenol and Bisphenol A Accumulation and Implication for Ecological and Human Health. *Science of the Total Environment*, 502: 417-425.
13. Nielsen, E., Ostergaard, G., Thorup, I., Ladefoged, O. and Jelnes J. E. (2000). Toxicological Evaluation and Limit Values for Nonylphenol, Nonylphenol Ethoxylates, Tricresyl, Phosphates and Benzoic Acid. Danish Environmental Project No. 512.
14. EFSA: European Food Safety Authority (2013). Scientific Opinion on the Toxicological Evaluation of Phenol, EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids. *European Food and Safety Authority*, 11(4):3189.

15. Ukoha, P. O., Ekere, N. R., Udeogu U. V. and Agbazue, V. E. (2014). Potential Health Risk Assessment of Heavy Metals [Cd, Cu and Fe] Concentrations in Some Imported Frozen Fish Species Consumed in Nigeria. *International Journal of Chemical Sciences*, 12 (2): 366-374.
16. Tesi, G. O. and Iniaghe, P. O. (2020). Polychlorinated Biphenyls in Canned Sardines in Nigeria and Risk Assessment. *Food Additives and Contaminants: Part B*.
17. Chen, C., Zou, W., Chen, S., Zhang, K. and Ma, L. (2020). Ecological and Health risk Assessment of Organochlorine Pesticides in an Urbanized River Network of Shanghai, China. *Environmental Science Europe*, 32:42.
18. USEPA: United States Environmental Protection Agency (2000). Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories. Vol. 1: Fish Sampling and Analysis, 3<sup>rd</sup> Edition, Pp 169-291.
19. Adeyi, A. A. (2020). Distribution and Bioaccumulation of Endocrine Disrupting Chemicals (EDCS) in Water, Sediment and Fish in Lagos Lagoon. *Ife Journal of Science*, 22: 2.
20. Mao, 1-F, Lu, Y-Y. and Chen, M-L. (2006) A Simplified Method for Simultaneous Quantitation of Alkylphenols and Alkylphenol Ethoxylates in Meat and Fish Using high-performance Liquid Chromatography with Fluorescence Detection, *International Journal of Environmental Analytical Chemistry*, 86 (10): 713-722.
21. Lee, C. C., Hsieh, C-Y., Chen, C. S. and Tien, C-J. (2020). Emergent Contaminants in Sediments and Fishes from the Tamsui River (Taiwan): Their Spatial-Temporal Distribution and Risk to Aquatic Ecosystems and Human Health. *Environmental Pollution*, 258: 113733
22. Datta, S., Loyo-Rosales, J. E. and Rice, C. P. (2002). A Simple Method for the Determination of Trace Levels of Alkylphenolic Compounds in Fish Tissues Using Pressurized Fluid Extraction, Solid Phase Cleanup, and High-Performance Liquid

- Chromatography Fluorescence Detection. *Journal of Agriculture and Food Chemistry*, 50: 1350-1354.
23. Sauter, M. and Licha, T. (2002). Use of Short Chained Alkylphenol (SCAP) in Analysis Transport Behaviour of Oil Contaminated Ground Water. *Agricultural Science*, 7(2): 29-37.
24. Verma, C., Hussain, C.M., Quraishi, M. A. and Alfantazi, A. (2023). Green Surfactants for Corrosion Control: Design, Performance and Applications. *Advances in Colloid and Interface Science*, 311:
25. Rice, C. P., Schmitz-Alfonso, I., Loyo-Rosales, J. E., Link, E., Thoma, R., Fay, L., Altfater, D. and Camp, M. J. (2003). Alkylphenol and Alkylphenol Ethoxylates in Carp, Water and Sediment from the Cuyahoga River, Ohio. *Environmental Science and Technology*, 37: 3747-3754.
26. Lalonde, B. and Garron, C. (2021). Nonylphenol, Octylphenol and Nonylphenol Ethoxylates Dissemination in the Canadian Fresh Water Environment. *Archives of Environmental Contamination and Toxicology*, 80: 319–330.