

Advanced Treatment of Type 1 Diabetes with Teplizumab: Mechanism and Clinical Efficacy

ABSTRACT

Teplizumab, a humanized anti-CD3 monoclonal antibody, has emerged as a significant advancement in the treatment of type 1 diabetes (T1D), an autoimmune disease characterized by the destruction of insulin-producing pancreatic beta cells. As the first FDA-approved disease-modifying drug for T1D, teplizumab has demonstrated efficacy in delaying the onset of stage 3 T1D in adults and pediatric patients 8 years and older with stage 2 T1D. Clinical trials have shown that teplizumab can preserve beta cell function, reduce insulin requirements, and improve glycemic control in newly diagnosed T1D patients. The drug's mechanism of action involves modulating the immune response responsible for beta cell destruction by altering the function of T-lymphocytes, inducing regulatory T cells, and promoting the accumulation of exhausted-like CD8 T cells. Teplizumab's efficacy appears to be influenced by factors such as the microbiome and the induction of partially exhausted CD8 T cells, which exhibit reduced secretion of inflammatory cytokines. While teplizumab has shown promise in preserving beta cell function and delaying T1D onset, challenges remain in optimizing treatment protocols and addressing potential side effects, including lymphopenia and skin disorders. Future research directions include exploring combination therapies, identifying predictive biomarkers, and refining patient selection to maximize treatment outcomes. Teplizumab's success in T1D has implications for the broader field of immunotherapy in autoimmune diseases, demonstrating the potential of targeted immunomodulation in altering disease.

Keywords: teplizumab, type 1 diabetes, autoimmune disease, beta cells, FDA-approved, disease-modifying drug, delayed onset, preserved beta cell function, modulating immune response, immunotherapy

1. INTRODUCTION

1.1 Background on Type 1 Diabetes (T1D) and its autoimmune nature

Type 1 Diabetes (T1D) is a chronic autoimmune disease that is characterized by the specific destruction of the insulin-producing pancreatic beta cells, leading to severe insulin deficiency and hyperglycemia^{1,2}. The autoimmune process usually commences in genetically predisposed subjects mostly affected in childhood and adolescence¹. The destruction of beta cells is progressive, and clinical manifestations typically set in once more than 80% of the cells are gone². However, it is important to note that the autoimmune nature of T1D cannot be said to rest on any dysregulation of the immune system exclusively. Recent studies suggest that some apoptosis ensues as the beta cells engage in complex interactions with the immune system³. This fundamentally challenges the view that T1D is merely an immune-mediated disease and places greater importance on understanding the vulnerability and heterogeneity of beta cells in disease progression. T1D pathogenesis arises from a complex interplay among the genetic factors, environmental triggers, and the resulting immune responses^{4,5}. Although genetic susceptibility

majorly affects, with HLA-DQ being one risk factor, many of those diagnosed with T1D have no family history or high-risk HLA haplotypes⁴. Such heterogeneity in disease onset and progression depicts the requirements For a better understanding of T1D pathogenesis that could lead to partly the concept of T1D endotypes and personalization of disease diagnosis and treatment^{4,6}

1.2) Overview of the significance of teplizumab treatment.

Teplizumab is a CD3-directed monoclonal antibody that is a significant milestone in type 1 diabetes (T1D) treatment. It was approved in November 2022 in the USA for delaying the onset of Stage 3 T1D among adult and pediatric patients aged 8 years and older who are suffering from Stage 2 T1D⁷. First promising examples of retaining beta cell function and delaying the diagnosis of T1D in high-risk individuals have emerged⁸.

Interestingly, the effect of teplizumab extends long ahead of the immediate treatment. Teplizumab treatment for a period of 14 days demonstrated extended effects characterized by delaying T1D diagnosis and retaining beta cell function in high-risk individuals⁸. There are evidence that drug can induce specific and substantial changes in T cell function and reactivity, still after 3 months and as lasting as 18 months after drug treatment began⁹. Teplizumab may induce long-lasting immune changes that contribute to its ameliorative effect. As a conclusion, teplizumab represents a huge breakthrough in T1D treatment, offering the first-ever therapy to modify a disease for individuals at risk of developing T1D. This could significantly improve the quality of life for patients by maintaining insulin production and delaying the onset of the disease. However, while it shows much promise, teplizumab is also associated with potential adverse effects with respect to the gastrointestinal, dermatological, and hematological systems⁹. Ongoing research builds this up further to enhance both its efficacy and safety profile¹⁰.

2)The Immune System in T1D

2.1) Description of the immune system's role in T1D

Teplizumab is a humanized monoclonal antibody tethered to CD3 antigen and plays a critical role in modulation of immune response in type 1 diabetes (T1D). In T1D, destruction of insulin-producing β -cells begins with the activation of effector T-lymphocytes that are cytotoxic¹¹. Teplizumab does induce regulatory T-cells (Tregs) responses and inhibits inflammatory and destructive processes taking place in target tissues^{11,12}. Interestingly, teplizumab's mode of action is one that complexes immunological arrayed changes. PD-1+ Foxp3+ Treg cell numbers were observed to rise in the ones off anti-drug antibodies in their first course of therapy¹². Additionally, teplizumab was found to enhance the population of CD8 T-cells resembling exhausted T-cells characterized by high expression of EOMES transcription factors, effector molecules, and multiple inhibitory receptors¹³. In conclusion, teplizumab complexes multiple mechanisms in the immune system of T1D, including induction of Tregs, modulation of CD4 and CD8 populations, and promotion of T-cell exhaustion. These synergized immunomechanisms underlie the ability of the drug in preserving β -cell function and delaying the onset of T1D in high-risk individuals¹⁴⁻¹⁶.

2.2) Mechanisms of T-cell involvement in β -cell destruction.

Phase III clinical trials which are underway for teplizumab have shown promise in delaying the onset of T1D^{17,18}. Teplizumab works on multiple targets and inhibits the autoimmune destruction of insulin-producing β -cells mediated by effector T-lymphocytes¹¹. Mechanism-wise, teplizumab has demonstrated increased Treg activity and promotes immune tolerance¹⁸. The use of teplizumab has led to the transient increasing of PD-1+Foxp3+ Tregs and perhaps a CD4 T cell that was not capable of functioning any longer, as well as a profound tally of exhausted CD8 T cells¹⁹. The phenotypic changes across T cells were correlated with great treatment responses in the nonresponders to therapies, who otherwise developed anti-drug antibodies¹⁹. To sum up, teplizumab protects β -cell function by modulating multiple T-cell subtypes that include enhancing Treg function while inhibiting the activation of effector CD4 and CD8 T cells¹⁹. This coordinated immune modulation may contribute to halting the autoimmune destruction of β -cells in T1D. The underlying mechanisms remain ambiguous, however, and additional work will be needed before the intricate immunological effects of teplizumab in T1D might be known^{19,20}.

3) Overview of Teplizumab

3.1) Overview of Teplizumab

Humanized anti-CD3 monoclonal antibody that acts to prevent and treat type 1 diabetes (T1D)^{7,10}. This includes the first from the FDA to have its use in delaying the onset of stage 3 T1D in patients aged 8 years or older with stage 2 T1D. Teplizumab alters the function of T-lymphocytes that mediate the destruction of insulin-producing β -cells in the pancreas²¹. Clinical trials have demonstrated teplizumab's efficacy in preserving C-peptide response, reducing exogenous insulin use, and improving glycemic control in newly diagnosed T1D patients²²⁻²⁴. Not all patients respond similarly to the drug and the duration of response for some can be up to 7 to 10 months¹⁵. Interestingly, metabolic and immunologic features at baseline may help identify subgroups with robust responses to teplizumab¹⁵. However, teplizumab not only had great promise in T1D management is also associated with a few adverse effects. These side effects can include skin rashes, upper respiratory infections, headaches and nausea¹⁵. Likewise, when prescribed teplizumab, patients carry a greater risk of experiencing gastrointestinal, dermatological, and hematological side effects from treatment, with already the potential risk for serious side effects and/or even active Epstein-Barr Virus infection⁹. Though many such side effects may seem serious, teplizumab is seen as a major advancement in T1D therapy with a new way to slow down disease progression in T1D patients and perhaps improve their long-term health outcomes.

3.2) Pharmacological profile and classification of monoclonal antibodies.

Teplizumab is humanized anti-CD3 monoclonal antibody that modifies the function of T-lymphocytes in the destruction of insulin-producing β -cells in type 1 diabetes (T1D)²¹. It is an immunomodulatory agent that belongs to a class of CD3-directed monoclonal antibodies⁷. Teplizumab works by targeting CD3+ cells, which activate T-lymphocytes of both CD4+ and CD8+ types for the destruction of pancreatic beta-cells in T1D²⁵. Fascinatingly, while teplizumab shows promise in delaying the onset of stage 3 T1D in high-risk individuals, its efficacy may vary depending on the timing of administration and dosage. A Phase 3 trial did not

establish the benefit in T1D patients diagnosed with T1D within the past 12 weeks with the lower cumulative dose²¹. It stresses the need for determining the right dose and time to make monoclonal antibodies reach their therapeutic potential. In conclusion, teplizumab showcases the pharmacological complexity of monoclonal antibodies, highly selective, specific, and enduring in the body²⁶. Its approval has been a landmark achievement in the treatment of T1D^{7,25}. Remarkably, like all monoclonal antibodies, teplizumab may face an uphill task in terms of bioavailability, absorption, and potential immune responses, and hence deserves meticulous investigation to display its application in clinical practices and further scrutiny.

4) Mechanism of Action

4.1) Detailed explanation of teplizumab's interaction with the immune system

Teplizumab is an Fc receptor non-binding anti-CD3 monoclonal antibody that acts through various mechanisms to modulate the immune system and control T1D progression. Teplizumab modulates T cell populations by decreasing circulation CD4+ effector memory T cells while increasing the CD8+ T cell population, specifically CD8+CM in clinical responders²⁷. The expanded CD8+CM T cells show evidence of proliferation and also differentiation from non-CM T cells. These CD8CM T cells showed decreased expression of genes related to immune activation and increased expression of genes related to T cell differentiation and regulation²⁷. Furthermore, it has been shown to induce regulatory CD8+ T cells (CD8+ Treg) capable of inhibiting CD4+ T cell proliferation-somehow in reliance on TNF and CCL4²⁸. They express markers of CD25, CTLA-4, Foxp3, and TNFR2 with the subpopulation capable of co-expressing TNFR2 and CD25 classifiable as a very potent one²⁸. Interestingly, teplizumab causes gut-tropic CCR6+ T cells to migrate to the small intestine, where they produce the regulatory cytokine IL-10²⁹. This effect becomes a key element in the efficacy of the drug since blocking the milieu will render treatment through teplizumab futile²⁹. The long-term effects include induction of the partially exhausted profile in CD8+ T cells expressing EOMES and the effector molecules along with a wide range of inhibitory receptors like TIGIT and KLRG1³⁰. These cells recognize a broad spectrum of environmental and auto-antigens but are hypo-proliferative in response to stimulation-polyclonal. In summary, teplizumab interacts with the immune system by modulating T cell numbers, inducing regulatory T cells, changing T cell migration patterns, and rendering T cells: still preferring a state of-exhausted state among CD8+ T cells. All of these mechanisms contribute to the drug's ability to delay T1D progression in at-risk patients and preserve β -cell function in patients with early-onset T1D.

4.2) Effects on CD3 signaling and T cell modulation

Teplizumab, being an anti-CD3 monoclonal antibody, has some significant effects on CD3 signaling and T cell modulation in type 1 diabetes (T1D). The molecule acts on CD3 critical for T cell activation and signaling³¹. Teplizumab operates through modulation of CD3 on T cells resulting in altered T cell function and phenotype. Studies have shown that teplizumab treatment decreases circulating CD4+ effector memory T cells but does increase CD8+ central memory T cells in clinical responders²⁷. These CD8CM T cells show lowered expression of activation genes and increased expression of genes related to T cell differentiation and regulation. Furthermore, teplizumab expands the KLRG1+TIGIT+CD8+ T cell subpopulation with partial exhaustion, as

evidenced by diminished secretion of the pro-inflammatory cytokines $IFN\gamma$ and $TNF\alpha$ ^{8,32}. Notably, T cells experience a biphasic response after teplizumab treatment: any initial transcriptional signature of cellular activation—a phenomenon expressed in both CD4+ and CD8+ T cells—will revert to lower levels of expression of genes involved in T cell receptor and activation signaling pathways in clinical responders over time³³. This ultimately will promote operational tolerance in T1D in a manner where T cells activate, successively show exhaustion and regulation. Besides, teplizumab treatment wards off autoreactive CD8+ T cells from profusing in groups at the control³³. Therefore, the actions of teplizumab on CD3 signaling and T cell modulation are complex and multilayered. The drug modifies the composition of T cell subsets, alters the expression profiles of genes and induces a state of partial exhaustion and regulation in CD8+ T cells. These immunomodulated effects contribute to the delay in T1D progression and better beta cell function that have been observed in clinical trials^{7,8}. The ability of teplizumab to induce these changes in T cell function and phenotype highlights its potential as a promising therapeutic approach for T1D management.

4.3) Consequences for beta cell preservation and insulin production

The new research has used teplizumab for considerable beta cell function preservation and to delay the onset of type 1 diabetes (T1D). Clinical trials have established that teplizumab can stabilize C-peptide, and that must be preserved to assess the function of beta cells, and the use of exogenous insulin diminished, particularly for patients who had an onset of T1D³⁴. In high-risk subjects, teplizumab would delay the median time to market diagnosis by about 24 months compared to placebo (48.4 vs 24.4 months)³⁵. Interestingly, the effects of teplizumab seemed to persist even beyond the treatment period. Some subjects have shown long-lasting positive results, with teplizumab halting the severe decline of beta cells and perhaps even contributing to their recovery after treatment among high-risk populations^{36,37}. Nevertheless, it has to be borne in mind that, notwithstanding the promise of teplizumab in trial patients, long-term blood glucose modulation and insulin requirement has not yet been established in everyone³⁷. In conclusion, teplizumab represents a significant development in the management of T1D, with the promise of delaying disease onset and preserving beta cell function. Its immune-modulating capacity and potential to enhance beta cell function give teplizumab the prospect of a disease-modifying treatment for T1D³⁸. However, more work is needed towards optimizing treatment regimens, identifying ideal patient populations, and exploring combination therapies to maximize benefits in insulin production and beta cell function preservation³⁹.

5) Clinical Evidence

5.1) Summary of major clinical trials and studies

Teplizumab, an FcR nonbinding anti-CD3 monoclonal antibody, has been shown to have therapeutic effects in several randomized clinical trials on type 1 diabetes (T1D)²⁷. Multiple trials have demonstrated its efficacy in preserving β -cell function and decreasing insulin dependence in T1D patients. A systematic review and meta-analysis of 8 RCTs found that teplizumab was effective in substantially reducing the need for insulin and improving C-peptide levels, implying an intrinsic generation of the insulin. No significant alteration of HbA1c levels was noted⁴⁰. Another meta-analysis of 8 RCTs with 1908 T1D patients showed findings consistent with the previous analysis; decreased insulin consumption, enhanced C-peptide responses and significant

changes in HbA1c with reported minimum side effects⁴¹. Quite interestingly however, other studies reported conflicting results. The phase 3 study of 516 patients showed no significant difference between teplizumab and placebo arms with respect to the primary outcome endpoint, defined as the percentage of patients with insulin use of less than 0.5 U/kg each day and HbA1c less than 6.5% at 1 year. Only 5% of teplizumab patients were free from insulin therapy at 1 year versus none in the placebo group⁴². Thus, while the majority of studies support teplizumab in the preservation of β -cell function and decreased insulin dependence in T1D patients, there are trials that have also suggested conflicting results. Safety profiles appear well established, given that rashes represented the most frequent adverse event⁴². Opening further lines of research will be required to mandate an appropriate dose regimen together with the patient population that will optimally benefit from teplizumab treatment.

5.2) Efficacy outcomes regarding beta cell function and long-term diabetes management.

Teplizumab therapy indeed shows optimistic efficacy results, particularly towards pancreatic beta cell function and its long-term management for the case of type 1 diabetes (T1D). Studies have demonstrated that a single 14-day course of teplizumab treatment could delay T1D diagnosis significantly while improving beta cell function in high-risk individuals. In a randomized controlled trial with extended follow-up, median time to diagnosis was 59.6 months in teplizumab-treated individuals, compared to 27.1 months in placebo-treated individuals⁸. Notably, 50% of those treated with teplizumab remained diabetes-free, in contrast to 22% of placebo participants. Treatment with teplizumab also improved beta cell function, as shown by greater average C-peptide area under the curve (1.94 vs. 1.72 pmol/ml; $P = 0.006$)⁸. Interestingly, the efficacy of teplizumab seems to be correlated with some changes in T cell subsets. C-peptide levels improved with the rise of partially exhausted memory KLRG1+TIGIT+CD8+ T cells, which were shown to produce lower levels of inflammatory cytokines IFN γ and TNF α . It indicates that teplizumab may act directly upon the immune response, thereby allowing the preservation of beta cell function. To conclude, teplizumab therapy is promising for delaying T1D onset, preserving beta cell function, and facilitating long-term management of diabetes. The ongoing positive result of a single course of treatment on both clinical outcomes and immunological parameters gives it renewed promise as a novel therapeutic approach in T1D. Further research is still required to develop better treatment protocols and to assess long-term safety and efficacy outcomes in larger patient populations^{34,43}.

5.3) Comparative analysis with other treatments

Teplizumab is a CD3-directed monoclonal antibody that shows preliminary promise for treatment and prevention of type 1 diabetes (T1D). Several studies have shown its effect in the preservation of β -cell function and in decreasing the insulin requirement in T1D^{23,40,41}. When compared to placebo, teplizumab demonstrated consistently greater outcomes in maintaining C-peptide levels, a marker of endogenous insulin production. Meta-analyses demonstrated that C-peptide levels were significantly elevated in teplizumab-treated groups compared to control groups at various time points up to 24 months after treatment^{23,40}. Furthermore, teplizumab treatment was correlated with decreased use of exogenous insulin with comparable glycemic control^{15,23}. Interestingly, while teplizumab has demonstrated apparent benefits in preserving β -cell function and decreasing insulin dependence, whether this affects long-term glycemic control,

as measured by HbA1c levels, remains undetermined. Some studies describe significant changes in HbA1c levels⁴¹, but others have found no significant differences among teplizumab and placebo groups⁴⁰. Such differences underscore the need for further studies elucidating teplizumab long-term glycemic effects. In summary, teplizumab adds to the latest advances in T1D treatment, which offer a disease-modifying approach that goes beyond traditional insulin therapy. Its ability to postpone the onset of stage 3 T1D in high-risk individuals and preserve β -cell function in patients newly diagnosed with the disease positions it as a promising therapeutic opportunity. However, inherent variability in response to treatment and the need for repeated dosing indicates that further research is needed to develop better treatment protocols for the identification of the most suitable recipient population for this therapy.

6) Challenges and Safety Considerations

6.1) Overview of potential side effects and risks

This includes teplizumab, a monoclonal antibody targeted against CD3 that was approved for delaying the onset of Stage 3 T1D among patients at risk. These results are promising despite the potential tradeoffs. Various adverse effects have been reported with teplizumab treatment⁷. Patients receiving teplizumab had a greater risk of developing gastrointestinal, dermatological, and hematological side effects than the placebo. Specifically, this included a considerable increase in the odds of nausea, rash, and lymphopenia^{9,23}. Moreover, teplizumab-treated patients had a higher risk of active Epstein-Barr virus infection⁹. Notably, while individual adverse effects were enriched for teplizumab, no individual study reported significantly elevated total adverse effects from teplizumab relative to placebo²³. Studies provide contradictory evidence as to whether teplizumab is correlated with higher-grade adverse events and discontinuation events. Overall, teplizumab shows promise in delaying T1D progression but exhibits multiple side effects, the main ones affecting the gastrointestinal, dermatological, and hematological systems. Teplizumab's safety profile requires monitoring, given that this drug is newly introduced and hardly has any long-term data available to predict the safety and consequential benefit of teplizumab treatment. However, post-marketing surveillance and more research is needed to address safety, tolerability, and efficacy-related issues with teplizumab treatment^{9,14}.

6.2) Consideration of patient eligibility and monitoring protocols.

Teplizumab, a CD3-directed monoclonal antibody, has shown promise in delaying the onset of stage 3 type 1 diabetes (T1D) in high-risk individuals. Patient eligibility for teplizumab treatment typically includes adults and children aged 8 years and older with stage 2 T1D^{7,44}. The drug is administered parenterally over a 14-day course^{40,45}, with some studies evaluating the effects of one or two courses. Monitoring protocols for teplizumab treatment involve assessing several key parameters. C-peptide levels are consistently used as a primary outcome measure, indicating endogenous insulin production. Insulin use and HbA1c levels are also monitored to evaluate the drug's efficacy in maintaining glycemic control^{40,46}. Additionally, safety monitoring is crucial, with common adverse events including lymphopenia, rash, and headache, which generally resolve without intervention⁴⁵. Interestingly, a study found that parents with prior experience of diabetic ketoacidosis (DKA) showed a higher inclination to consent to teplizumab treatment for their at-risk children⁴⁷. This highlights the importance of educating patients and caregivers about

the potential benefits and risks of teplizumab. Long-term monitoring of T cell phenotypes and gene expression patterns may also be valuable, as persistent EOMES expression was identified as a predictor for the time to clinical T1D diagnosis⁴⁸.

7)Future Perspectives

7.1)Emerging research directions and potential for combination therapies.

Whereas teplizumab, a humanized anti-CD3 monoclonal antibody, has become an ideal candidate disease-modifying therapy for T1D, it has shown efficacy in delaying T1D onset in at-risk individuals and preserving insulin production in newly diagnosed patients^{43,49}. Clinical trials have demonstrated such teplizumab treatment to slow the decline of C-peptide production, improve glycemic control, and decrease exogenous insulin requirement for up to two years post-treatment^{23,45}. Interestingly, research has identified subgroups of responders with unique metabolic and immunological attributes, to the effect that future applications may focus on personalized treatment approaches^{45,50}. This finding opens a new area of investigation for the identification of predictive biomarkers that could point to the patients most likely to benefit from teplizumab therapy. In one regard, teplizumab shows promise as a monotherapy, but with an increase in various combination therapies to enhance its efficacy to address the multifaceted nature of T1D pathogenesis. Future research directions for teplizumab, optimizing treatment regimens, long-term efficacy and safety, and exploring for potential synergies with other emerging therapies such as mesenchymal stem cells, gene therapy⁴³, and islet transplantation. In addition, studies are needed to evaluate the cost and effectiveness of teplizumab treatment, address challenges of treatment accessibility, and explore for its possible applications in other autoimmune diseases⁵⁰. As T1D therapy keeps on evolving, teplizumab remains a great milestone, with research around it ongoing to optimize its therapeutic potential and improve patient outcomes.

7.2)Teplizumab's role in the broader context of immunotherapy for autoimmune diseases.

Teplizumab is an anti-CD3 monoclonal antibody that could make a significant advance in immunotherapy for autoimmune diseases, especially Type 1 Diabetes (T1D). It shows some promise in delaying T1D onset in individuals at a higher risk of developing this disease and/or slowing disease progression in newly diagnosed patients^{18,50}. As an immunomodulator, teplizumab exerts its role by targeting T-cells as well as enhancing regulatory T-cell activity and consequently supporting immune tolerance^{18,51}. Curiously, while teplizumab has received specific approval for T1D, immunomodulatory doses are being studied for various other autoimmune diseases. For example, ablative chemotherapy with stem cell support and targeted immunotherapy using monoclonal antibodies such as rituximab have found their way in Multiple Sclerosis, systemic sclerosis, and Rheumatoid arthritis⁵². This suggests that essential principles will underline teplizumab effectiveness in broader contexts into autoimmune disease management.

Conclusion

In conclusion, the success of teplizumab in T1D treatment reflects the potential of targeted immunotherapy in dealing with autoimmune diseases. Its ability to delay the onset of disease and slow its progression demonstrates a paradigm shift in the management of T1D⁵³. As research progresses, the role of teplizumab may expand, and applications akin to it for other autoimmune conditions could develop, revolutionizing their treatment strategies.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Bastos MD, Kowalski TW, Mariath LM, et al. Polymorphisms and genetic susceptibility of type 1 diabetes mellitus and celiac disease. *Diabetol Metab Syndr.* 2015;7(S1):A216, 1758-5996-7-S1-A216. doi:10.1186/1758-5996-7-S1-A216
2. Assmann TS, Coutinho MKP, Tschiedel B, Canani LH, Crispim D. Circulating microRNAs as biomarkers for type 1 diabetes mellitus. *Diabetol Metab Syndr.* 2015;7(S1):A206, 1758-5996-7-S1-A206. doi:10.1186/1758-5996-7-S1-A206
3. Toren E, Burnette KS, Banerjee RR, Hunter CS, Tse HM. Partners in Crime: Beta-Cells and Autoimmune Responses Complicit in Type 1 Diabetes Pathogenesis. *Front Immunol.* 2021;12:756548. doi:10.3389/fimmu.2021.756548

4. D'Addio F, Ben Nasr M, Lunati ME, Fiorina P. Autoimmune (Type 1) diabetes. In: *The Rose and Mackay Textbook of Autoimmune Diseases*. Elsevier; 2024:585-601. doi:10.1016/B978-0-443-23947-2.00075-8
5. Long SA, Buckner JH. Clinical and experimental treatment of type 1 diabetes. *Clin Exp Immunol*. 2022;210(2):105-113. doi:10.1093/cei/uxac077
6. Dooley J, Tian L, Schonefeldt S, et al. Genetic predisposition for beta cell fragility underlies type 1 and type 2 diabetes. *Nat Genet*. 2016;48(5):519-527. doi:10.1038/ng.3531
7. Keam SJ. Teplizumab: First Approval. *Drugs*. 2023;83(5):439-445. doi:10.1007/s40265-023-01847-y
8. Sims EK, Bundy BN, Stier K, et al. Teplizumab improves and stabilizes beta cell function in antibody-positive high-risk individuals. *Sci Transl Med*. 2021;13(583):eabc8980. doi:10.1126/scitranslmed.abc8980
9. Buddhavarapu V, Dhillon G, Grewal H, Sharma P, Kashyap R, Surani S. Safety of teplizumab in patients with high-risk for diabetes mellitus type 1: A systematic review. *World J Diabetes*. 2024;15(8):1793-1801. doi:10.4239/wjd.v15.i8.1793
10. Ahmed N, Kumari A, T TSL, V HRSL. Teplizumab: The Newest Weapon in the Fight against Type 1 Diabetes. *J Clin Pharm Res*. Published online April 30, 2023:18-23. doi:10.61427/jcpr.v3.i2.2023.93
11. Pillemer SR. Preserving β Cells in Type 1 Diabetes mellitus: the role of immunological tolerance. *Drug Dev Res*. 2008;69(3):153-157. doi:10.1002/ddr.20240
12. Long SA, Thorpe J, Herold KC, et al. Remodeling T cell compartments during anti-CD3 immunotherapy of type 1 diabetes. *Cell Immunol*. 2017;319:3-9. doi:10.1016/j.cellimm.2017.07.007
13. Long SA, Thorpe J, DeBerg HA, et al. Partial exhaustion of CD8 T cells and clinical response to teplizumab in new-onset type 1 diabetes. *Sci Immunol*. 2016;1(5):eaai7793. doi:10.1126/sciimmunol.aai7793
14. Fanaropoulou NM, Tsatsani GC, Koufakis T, Kotsa K. Teplizumab: promises and challenges of a recently approved monoclonal antibody for the prevention of type 1 diabetes or

preservation of residual beta cell function. *Expert Rev Clin Immunol.* 2024;20(2):185-196. doi:10.1080/1744666X.2023.2281990

15. Herold KC, Gitelman SE, Ehlers MR, et al. Teplizumab (Anti-CD3 mAb) Treatment Preserves C-Peptide Responses in Patients With New-Onset Type 1 Diabetes in a Randomized Controlled Trial. *Diabetes.* 2013;62(11):3766-3774. doi:10.2337/db13-0345

16. Seewoodhary J, Silveira A. Teplizumab – preventative approaches to type 1 diabetes mellitus. *Pract Diabetes.* 2023;40(2):35. doi:10.1002/pdi.2448

17. Masharani UB, Becker J. Teplizumab therapy for type 1 diabetes. *Expert Opin Biol Ther.* 2010;10(3):459-465. doi:10.1517/14712591003598843

18. Thakkar S, Department of Endocrinology, Indraprastha Apollo Hospitals, New Delhi, India, Chopra A, et al. Teplizumab in Type 1 Diabetes Mellitus: An Updated Review. *TouchREVIEWS Endocrinol.* 2023;19(2):7. doi:10.17925/EE.2023.19.2.7

19. Long SA, Thorpe J, Herold KC, et al. Remodeling T cell compartments during anti-CD3 immunotherapy of type 1 diabetes. *Cell Immunol.* 2017;319:3-9. doi:10.1016/j.cellimm.2017.07.007

20. Vudattu NK, Herold KC. Treatment of new onset type 1 diabetes with teplizumab: successes and pitfalls in development. *Expert Opin Biol Ther.* 2014;14(3):377-385. doi:10.1517/14712598.2014.881797

21. Skelley JW, Elmore LK, Kyle JA. Teplizumab for Treatment of Type 1 Diabetes Mellitus. *Ann Pharmacother.* 2012;46(10):1405-1412. doi:10.1345/aph.1R065

22. Ma XL, Ge D, Hu XJ. Evaluation of teplizumab's efficacy and safety in treatment of type 1 diabetes mellitus: A systematic review and meta-analysis. *World J Diabetes.* 2024;15(7):1615-1626. doi:10.4239/wjd.v15.i7.1615

23. Kamrul-Hasan ABM, Mondal S, Nagendra L, Yadav A, Aalpona FTZ, Dutta D. Role of Teplizumab, a Humanized Anti-CD3 Monoclonal Antibody, in Managing Newly Diagnosed Type 1 Diabetes: An Updated Systematic Review and Meta-Analysis. *Endocr Pract.* 2024;30(5):431-440. doi:10.1016/j.eprac.2024.03.006

24. Nourelden AZ, Elshanbary AA, El-Sherif L, et al. Safety and Efficacy of Teplizumab for Treatment of Type One Diabetes Mellitus: A Systematic Review and Meta-Analysis. *Endocr Metab Immune Disord - Drug Targets*. 2021;21(10):1895-1904. doi:10.2174/1871530320999201209222921
25. Sharma N, Das DD, Chawla PA. Journey of Teplizumab: A Promising Drug in the Treatment of Type 1 Diabetes Mellitus. *Curr Diabetes Rev*. 2025;21(1):e250124226249. doi:10.2174/0115733998261825231026060241
26. Wang W, Wang E, Balthasar J. Monoclonal Antibody Pharmacokinetics and Pharmacodynamics. *Clin Pharmacol Ther*. 2008;84(5):548-558. doi:10.1038/clpt.2008.170
27. Tooley JE, Vudattu N, Choi J, et al. Changes in T \square cell subsets identify responders to FcR \square nonbinding anti \square CD3 mAb (teplizumab) in patients with type 1 diabetes. *Eur J Immunol*. 2016;46(1):230-241. doi:10.1002/eji.201545708
28. Ablamunits V, Bisikirska B, Herold KC. Acquisition of regulatory function by human CD8⁺ T cells treated with anti \square CD3 antibody requires TNF. *Eur J Immunol*. 2010;40(10):2891-2901. doi:10.1002/eji.201040485
29. Waldron-Lynch F, Henegariu O, Deng S, et al. Teplizumab Induces Human Gut-Tropic Regulatory Cells in Humanized Mice and Patients. *Sci Transl Med*. 2012;4(118). doi:10.1126/scitranslmed.3003401
30. Long SA, Thorpe J, DeBerg HA, et al. Partial exhaustion of CD8 T cells and clinical response to teplizumab in new-onset type 1 diabetes. *Sci Immunol*. 2016;1(5):eaai7793. doi:10.1126/sciimmunol.aai7793
31. Ceuppens JL, Meurs L, Van Vaeck F, Van Wauwe JP. CD3 modulation inhibits pokeweed mitogen-induced T-cell help for immunoglobulin production. *Cell Immunol*. 1986;102(1):144-151. doi:10.1016/0008-8749(86)90333-3
32. Long SA, Thorpe J, DeBerg HA, et al. Partial exhaustion of CD8 T cells and clinical response to teplizumab in new-onset type 1 diabetes. *Sci Immunol*. 2016;1(5):eaai7793. doi:10.1126/sciimmunol.aai7793

33. Lledó-Delgado A, Preston-Hurlburt P, Currie S, et al. Teplizumab induces persistent changes in the antigen-specific repertoire in individuals at risk for type 1 diabetes. *J Clin Invest.* 2024;134(18):e177492. doi:10.1172/JCI177492
34. Kokori E, Olatunji G, Ogieuhi IJ, et al. Teplizumab's immunomodulatory effects on pancreatic β -cell function in type 1 diabetes mellitus. *Clin Diabetes Endocrinol.* 2024;10(1):23. doi:10.1186/s40842-024-00181-w
35. Herold KC, Bundy BN, Long SA, et al. An Anti-CD3 Antibody, Teplizumab, in Relatives at Risk for Type 1 Diabetes. *N Engl J Med.* 2019;381(7):603-613. doi:10.1056/NEJMoa1902226
36. Białek J, Nitka K, Więckowska-Deroń M, et al. Teplizumab - current state of knowledge on the effects of teplizumab in preventing the development of type 1 diabetes in people at risk. *J Educ Health Sport.* 2023;44(1):288-299. doi:10.12775/JEHS.2023.44.01.018
37. Wagner D, Bleich D. Multiple Immune Pathways to Type 1 Diabetes Mellitus: Lessons Learned from Human Clinical Trials and Animal Models of Disease Clinical Trials, Lessons Learned. *Med Res Arch.* 2023;11(11). doi:10.18103/mra.v11i11.4568
38. LeFevre JD, Cyriac SL, Tokmic A, Pitlick JM. Anti-CD3 monoclonal antibodies for the prevention and treatment of type 1 diabetes: A literature review. *Am J Health Syst Pharm.* 2022;79(23):2099-2117. doi:10.1093/ajhp/zxac244
39. Fanaropoulou NM, Tsatsani GC, Koufakis T, Kotsa K. Teplizumab: promises and challenges of a recently approved monoclonal antibody for the prevention of type 1 diabetes or preservation of residual beta cell function. *Expert Rev Clin Immunol.* 2024;20(2):185-196. doi:10.1080/1744666X.2023.2281990
40. Heidari E, Shafiee A, Noorian S, et al. Efficacy of teplizumab for treatment of type 1 diabetes: A meta-analysis of randomized controlled trials. *Diabetes Metab Res Rev.* 2024;40(4):e3806. doi:10.1002/dmrr.3806
41. Ma XL, Ge D, Hu XJ. Evaluation of teplizumab's efficacy and safety in treatment of type 1 diabetes mellitus: A systematic review and

- meta-analysis. *World J Diabetes*. 2024;15(7):1615-1626. doi:10.4239/wjd.v15.i7.1615
42. Sherry N, Hagopian W, Ludvigsson J, et al. Teplizumab for treatment of type 1 diabetes (Protégé study): 1-year results from a randomised, placebo-controlled trial. *The Lancet*. 2011;378(9790):487-497. doi:10.1016/S0140-6736(11)60931-8
43. Salama RAA, Patni MAMF, Ba-Hutair SNM, Wadid NA, Akikwala MS. Exploring Novel Treatment Modalities for Type 1 Diabetes Mellitus: Potential and Prospects. *Healthcare*. 2024;12(15):1485. doi:10.3390/healthcare12151485
44. Garg P, Sangeeta Bhanwra, Rajiv Kumar. Teplizumab: a new glimmer of hope for type 1 diabetic patients. *Int J Basic Clin Pharmacol*. 2023;12(3):499-502. doi:10.18203/2319-2003.ijbcp20231135
45. Herold KC, Gitelman SE, Gottlieb PA, Knecht LA, Raymond R, Ramos EL. Teplizumab: A Disease-Modifying Therapy for Type 1 Diabetes That Preserves β -Cell Function. *Diabetes Care*. 2023;46(10):1848-1856. doi:10.2337/dc23-0675
46. Nourelden AZ, Elshanbary AA, El-Sherif L, et al. Safety and Efficacy of Teplizumab for Treatment of Type One Diabetes Mellitus: A Systematic Review and Meta-Analysis. *Endocr Metab Immune Disord - Drug Targets*. 2021;21(10):1895-1904. doi:10.2174/1871530320999201209222921
47. Bombaci B, Passanisi S, Pecoraro M, et al. Use of teplizumab in children and adolescents at risk of type 1 diabetes: perspectives of parents and caregivers from an Italian Pediatric Diabetes Center. *Acta Diabetol*. 2024;61(5):635-642. doi:10.1007/s00592-024-02245-w
48. Lledo Delgado A, Preston-Hurlburt P, Currie S, Clark P, Herold K. Reshaping immune cells and the antigen-specific repertoire by anti-CD3 mAb teplizumab in Type 1 diabetes. *J Immunol*. 2024;212(1_Supplement):0958_5059-0958_5059. doi:10.4049/jimmunol.212.sup.0958.5059
49. Hirsch JS. FDA approves teplizumab: a milestone in type 1 diabetes. *Lancet Diabetes Endocrinol*. 2023;11(1):18. doi:10.1016/S2213-8587(22)00351-5

50. Fanaropoulou NM, Tsatsani GC, Koufakis T, Kotsa K. Teplizumab: promises and challenges of a recently approved monoclonal antibody for the prevention of type 1 diabetes or preservation of residual beta cell function. *Expert Rev Clin Immunol*. 2024;20(2):185-196. doi:10.1080/1744666X.2023.2281990
51. Eerike M, Konda VGR. Immunomodulatory therapy in diabetes mellitus: A pharmacological approach. In: *Biochemical Immunology of Diabetes and Associated Complications*. Elsevier; 2024:121-139. doi:10.1016/B978-0-443-13195-0.00007-7
52. Cohen Y, Nagler A. Treatment of refractory autoimmune diseases with ablative immunotherapy. *Autoimmun Rev*. 2004;3(2):21-29. doi:10.1016/S1568-9972(03)00083-1
53. Novograd J, Frishman WH. Teplizumab Therapy to Delay the Onset of Type 1 Diabetes. *Cardiol Rev*. 2024;32(6):572-576. doi:10.1097/CRD.0000000000000563