

Genetic variability and correlation studies in Crossandra(*Crossandra in fundibuliformis*(L.) (Nees)

Abstract

The present investigation was carried out to study the performance of crossandra genotypes for morphological, flowering, yield, quality and screening for nematode resistance at the Department of Floriculture and Landscape Architecture, Tamil Nadu Agricultural University, Coimbatore during the year 2018-2019. Ten genotypes were investigated for ten quantitative characters viz., Plant height (cm), Plant spread both E-W & N-S (cm), No. of primary branches, Number of leaves, 100 flower weight (g), Number of flowers per spike, Number of spikes per plant, Rachis length and Yield per plant (g). Phenotypic coefficient of variation recorded was higher than the genotypic coefficient of variation for all the parameters which shows that greater influence over environment in plants. High GCV and PCV were recorded for plant height, number of leaves, rachis length, number of spikes per plant and yield per plant. High heritability with high genetic advance was noted in plant height, number of leaves, primary branches, number of spikes per plant, number of flowers per spike, 100 flower weight, rachis length and yield per plant. High heritability with high genetic advance is an important character for selection and these are traits controlled by additive gene action. From the correlation studies Yield per plant was positively and significantly correlated with number of primary branches, number of spikes per plant and rachis length. Positive and significant correlation was also recorded in rachis length, plant spread, number of spikes per plant and number of flowers per spike. Hence, simultaneous selection for the above traits would be more rewarding to bring improvement in crossandra.

Introduction

Crossandra(*Crossandra in fundibuliformis*(L.) (Nees) is a commercial loose flower crop mostly grown in southern region of India, The performance of genotypes exhibit wide range of variability with the climatic condition of their growing habitat and genetic potential of the individual genotype. The estimated area under crossandra cultivation in Tamil Nadu was 610 hectares and 2050 tonnes of production with productivity of 3.36 tonnes during 2016- 2017 (Data.gov.in, 2016-2017). It also has some considerable medicinal property to cure wounds, aphrodisiac and pyorrhoea (Malars of India, 2016). Genotypic and phenotypic

coefficient of variation shows the extent of variability in different traits due to environment and inherent capacity of genotype. Correlation also helps to improve the yield related components using association between yield and yield components. Based on these observations, genotypes will be selected for further breeding program. Therefore, the present study is focussed on to assess the genetic variability present in the population by using coefficient of variation and study the heritability of the character and the correlation among yield and component traits.

Materials and Methods

The present investigation was carried out to study the performance of crossandra genotypes for morphological, flowering, yield, quality and screening for nematode resistance at the Department of Floriculture and Landscape Architecture, Tamil Nadu Agricultural University, Coimbatore during the year 2018-2019. Ten genotypes were raised in randomized block design with three replications. The rooted cuttings collected from different locations were transplanted into the field in one side ridge with the spacing of 60 × 45 cm. Before transplanting, the rooted cuttings were dipped in carbendazim (1g/l of water) to avoid fungal infection. Planting was done in the early morning hours and irrigation was given immediately after transplanting to minimize the transplanting shock. After good establishment, irrigation was given once in a week and gap filling was done regularly. The data was collected on ten yield and yield contributing characters *viz.*, Plant height (cm), Plant spread both E-W & N-S (cm), No. of primary branches, Number of leaves, 100 flower weight (g), Number of flowers per spike, Number of spikes per plant, Rachis length and Yield per plant (g). Among different genotypes, five plants were randomly selected and tagged for taking observations and the mean value was calculated for analysis on three replications. The cultivation practices like irrigation, weeding, fertilization and pesticide application *etc.* were followed on proper times. The analysis of variance of RBD and their significance for all the characters were worked out as suggested by Panse and Sukhatme (1954). The various genetic parameters *viz.*, ECV, GCV, PCV, heritability and GAM were calculated by adopting the formulae given by Johnson *et al.*, (1955).

Results and Discussion

The mean performance of all the characters is presented in Table 1. Among the genotypes collected, Arka Shravya (54.66 cm) recorded maximum plant height followed by Arka Ambara (48.16 cm). The variation in genotypes might be due to the genetic makeup of the individual genotype (Kulkarni and Reddy, 2004). The increased variation in plant height may

also due to elongation in meristematic region, rapid cell division and maximum internodal length of the plant. Arka Shravya recorded maximum East- West (44.00 cm) and North-South (33.33 cm) plant spread. The maximum plant spread may be due to the production of more number of branches per plant, branching habit, internodal length, flowering habit and genetic character of each genotype. Variation in plant spread might be due to additive gene action of the genotype. Similar results were obtained in Arka Shravya by Priyanka *et al.* (2017) and Tejaswi *et al.* (2019) in crossandra. The maximum number of primary branches (4.60) was observed in genotypes Arka Shravya, Arka Kanaka and Arka Ambara. This trait directly contributes to growth parameters of genotypes. Genotype Arka Shravya (210.00) recorded the maximum number of leaves followed by Arka Kanaka (198.00). This trait might be depending on number of branches and genetical makeup of the individual genotype. Similar increased variation were observed by Poornima *et al.* (2006); Zosiamliana *et al.* (2013) in china aster, Uddin *et al.* (2015); Suvija *et al.* (2016); Singh *et al.* (2017) and Niki *et al.* (2016) in chrysanthemum.

Maximum flower yield per plant was recorded in genotype Arka Shravya. Generally, this character depends on various independent characters like plant height, plant spread, more number of leaves, leaf area, flower size, weight of the flower, number of spikes and flowers per plant which help in production of more photosynthates resulting in high yield. The difference in flower yield potential is related to the additive gene effect. Similar results were reported by Ramachandrudu and Thangam (2010); Priyanka *et al.* (2017); Tejaswi *et al.* (2019) in crossandra.

The presence and magnitude of genetic variability is prerequisite for any breeding program. It will be useful for quantifying the genetic variability, heritability and genetic advance of genotypes to identify for high yielding characters. Phenotypic coefficient of variation recorded was higher than the genotypic coefficient of variation for all the parameters which shows that greater influence over environment in plants. High GCV and PCV were recorded for plant height, number of leaves, rachis length, number of spikes per plant and yield per plant. It indicates that maximum variability is observed in these characters. Moderate GCV and PCV was noticed in number of primary branches and 100 flower weight. Low GCV and PCV was found in plant spread.

High heritability was found in all the characters recorded. High heritability value of a particular character had less influence of environment which is highly useful in selecting genetically good individual. High heritability with high genetic advance was noted in plant

height, number of leaves, primary branches, number of spikes per plant, number of flowers per spike, 100 flower weight, rachis length and yield per plant. High heritability with high genetic advance is an important character for selection and these are traits controlled by additive gene action. Higher heritability with moderate genetic advance was recorded in plant spread. It is due to additive and non-additive gene effect. This character can be improved through hybridization or by selection. Similar reports were reported by Kadam *et al.* (2014) in gladiolus, Poornima *et al.* (2006) and Sankari *et al.* (2019) in china aster.

Correlation studies

The correlation coefficient shows a strong association between plant morphological characters coupled with yield. A positive correlation between desirable characters is favorable to the plant breeder focusing the genotypes in improvement of both the characters. Yield per plant was positively and significantly correlated with number of primary branches, number of spikes per plant and rachis length. Positive and significant correlation was also recorded in rachis length, plant spread, number of spikes per plant and number of flowers per spike. Number of spikes per plant was positively and significantly correlated with plant height, plant spread, number of primary branches and number of leaves. Positive and significant correlation was observed in number of flowers per spike with plant spread. 100 flower weight was positively and significantly correlated with number of primary branches. Number of leaves was positively and significantly correlated with plant height, number of primary branches and plant spread. Number of primary branches was positively and significantly correlated with plant height and plant spread. Positive and significant correlation was noted in plant spread and plant height. Similar observation was reported by Poornima *et al.* (2006) in China aster and Kumar *et al.* (2012) in chrysanthemum.

Table.1 Mean performance of ten genotypes for different characters

Genotypes	Plant height (cm)	Plant spread (cm) 120 DAT		No. of primary branches	No. of Leaves	Days to flower	100 flower weight (g)	Rachis length (cm)	Number of spikes per plant	Number of flowers per spike	Flower yield per plant (g)
	120 DAT	N-S	E-W								
Arka Shravya	54.66	33.33	44.00	4.60	210.00	61.33	6.86	13.25	72.00	53.33	142.00

Arka Kanaka	47.33	31.05	35.00	4.60	198.00	69.33	7.18	8.60	51.00	33.33	112.00
Arka Shreeya	45.20	27.06	33.00	4.30	168.67	63.00	8.99	6.33	49.00	29.33	98.00
Arka Ambara	48.16	29.73	36.20	4.60	186.89	65.00	7.14	8.33	47.00	30.66	87.00
Arka Chenna	37.26	28.98	32.08	3.80	132.26	64.23	7.12	5.60	29.00	20.00	84.00
Nilakkottai local	43.00	31.00	36.50	3.60	146.33	67.66	6.02	7.33	23.00	29.33	41.00
Dharmapuri local	38.66	25.54	31.65	3.20	128.34	78.89	5.75	6.53	13.00	28.00	22.00
Delhi crossandra	24.45	27.98	32.65	3.40	72.25	64.02	6.58	8.40	20.00	38.33	72.00
Villupuram local	37.67	26.65	33.65	3.40	136.52	63.00	6.26	6.67	32.00	31.55	81.00
Neyveli local	35.65	27.02	33.82	3.60	112.53	68.42	5.98	6.40	25.00	29.01	80.00
Mean	41.20	28.83	34.85	3.90	149.17	66.48	6.78	7.74	36.10	32.28	81.90
S. Ed	2.09	1.44	1.73	0.20	7.45	3.35	0.35	0.36	1.80	1.52	4.10
CD (P=0.05)	4.17	2.88	3.46	0.39	14.89	6.70	0.70	0.71	3.60	3.03	8.19

Table 2. Mean, genotypic and phenotypic co-efficient variation, heritability and genetic advance as percent of mean for different parameters of crossandra genotypes

S.No	Characters	Mean	GCV	PCV	Heritability	GA (%) of mean
1	Plant height 120 DAT	41.20	20.05	20.90	92.00	39.62
2	Plant spread 120 DAT in N-S direction	28.83	7.74	9.70	63.74	12.74
3	Plant spread 120 DAT in E-W direction	34.85	9.79	11.38	74.00	17.34
4	No. of primary branches on 120 DAT	3.90	13.91	15.10	85.05	26.44
5	Number of leaves at 120 days	149.17	28.10	28.70	95.88	56.67

6	100 flower weight	6.79	13.32	14.66	82.70	24.96
7	Number of flowers per spike	32.29	26.76	27.31	95.97	54.00
8	Number of spikes per plant	36.10	50.05	50.38	98.66	102.40
9	Rachis length	7.74	28.05	28.57	96.44	56.76
10	Yield per plant	81.90	41.32	40.92	98.00	83.42

Table 3. Genotypic correlation coefficient for different characters of crossandra genotypes

	1	2	3	4	5	6	7	8	9	10
1	1	0.637*	0.717*	0.798**	0.977**	0.346	0.366	0.813*	0.539	0.530
2		1	0.835**	0.679*	0.649*	0.107	0.571	0.682*	0.765**	0.591
3		.	1	0.593	0.664*	0.006	0.814**	0.758*	0.906**	0.619
4				1	0.866**	0.635*	0.379	0.907*	0.555	0.788**
5					1	0.422	0.358	0.859*	0.537	0.611
6						1	-0.010	0.569	0.011	0.536
7							1	0.622	0.946**	0.585
8								1	0.719*	0.905**
9									1	0.634*
10									.	1

1-Plant height 120 DAT

2- Plant spread 120 DAT in N-S direction

- | | |
|--|--------------------------------|
| 3- Plant spread 120 DAT in E-W direction | 4- Primary branches in 120 DAT |
| 5- Number of leaves | 6- 100 flower weight |
| 7- Number of flowers per spike | 8- Number of spikes per plant |
| 9- Rachis length | 10- yield per plant |
| *Significant at 5% level | **Significant at 1% level |

References

- Data.gov.in. 2016-2017. <https://data.gov.in/resources/area-production-and-productivity-crossandra-flowers-district-wise-tamil-nadu-year-2016-17>
- Johnson, H.W., H. Robinson and R. Comstock. 1955. "Estimates of genetic and environmental variability in soybeans 1." *Agronomy Journal* 47 (7):314-318.
- Kadam, G.B., G. Kumar, T. Saha, A. Tiwari and R. Kumar. 2014. "Varietal evaluation and genetic variability studies on gladiolus." *Indian Journal of Horticulture* 71 (3):379-384.
- Kulkarni, B.S. and B. Reddy. 2004. "Vegetative growth, flower yield and quality of different chrysanthemum cultivars." *Journal of Ornamental Horticulture* 7 (3):32-36.
- Kumar, M., S. Kumar, M.K. Singh, S. Malik and A. Kumar. 2012. "Studies on correlation and path analysis in chrysanthemum (*Dendranthema grandiflora* Tzvelev)." *Vegetos-An International Journal of Plant Research* 25 (2):62-65.
- Malars of india. 2016. <https://malarsfindia.blogspot.com/2016/01/crossandra-infundibuliformis.html=1>
- Niki, D., K. Sunil, S. Swati and C. Susmita. 2016. "Evaluation of chrysanthemum (*Chrysanthemum morifolium* Ramat) genotypes under West Garo Hills District, Meghalaya." *HortFlora Research Spectrum* 5 (3):189-194.
- Panse, V.G. and P.V. Sukhatme. 1954. "Statistical methods for agricultural workers."
- Poornima, G., D. Kumar and G. Seetharamu. 2006. "Evaluation of China aster (*Callestephus chinensis* (L.) ness) genotypes under hill zone of Karnataka." *Journal of Ornamental Horticulture* 9 (3):208-211.
- Priyanka, T., B. Kamble, R. Anuradha, N. Kumar and A. Shirol. 2017. "Evaluation of different crossandra genotypes under ghataprabha command area." *Journal of Pharmacognosy and Phytochemistry* 6 (6):252-254.

- Ramachandrudu, K. and M. Thangam. 2010. "Characterization and evaluation of local germplasm of crossandra (*Crossandra undulaefolia* Salisb.)." *Journal of Ornamental Horticulture* 13 (2):138-141.
- Sankari, A., M. Anand and B. Anita. 2019. "Evaluation of China Aster Cultivars (*Callistephus chinensis* (L.) Nees.) under Nilgiris." *Advances in Floriculture And urban Horticulture* 7(3):1649-1652.
- Singh, D.D., S. Tyagi, S. Singh and P.K. Ray. 2017. "Studies on the Performance and Flower Characterization of Chrysanthemum (*Dendranthema grandiflora* L.) Genotypes under Uttar Pradesh Conditions."
- Suvija, N., J. Suresh, R. Kumar and M. Kannan. 2016. "Evaluation of chrysanthemum (*Chrysanthemum morifolium* Ramat.) genotypes for loose flower, cut flower and pot mums." *International Journal Innovative Research Advanced Studies* 3 (4):100-104.
- Tejaswi, R., z. salma, A.V.D.D. Rao and S. Naresh. 2019. "study on growth, floral and yield parameters of crossandra genotypes under coastal andhra aradesh conditions." *Bulletin of environment, pharmacology and life sciences* 8:100-104.
- Uddin, A., T. Taufique, A. Ona, S. Shahrin and H. Mehraj. 2015. "Growth and flowering performance evaluation of thirty two chrysanthemum cultivars." *Journal of Bioscience Agriculture Research* 4 (01):40-51.
- Zosiamliana, J., G. Reddy and H. Rymbai. 2013. "Study on the performance of some varieties of China aster (*Callistephus chinensis* Ness.) in Andhra Pradesh." *Progressive Horticulture* 45 (2):312-316.