

Ornamental plants of Bignoniaceae family: Source of bioactive compounds with therapeutic applications and ecological services

Abstract

The Bignoniaceae family, comprising over 100 genera and 800 species, is a rich source of ornamental plants with medicinal properties. This review focuses on the pharmacological uses, bioactive compounds, and future aspects of selected ornamental plants of the Bignoniaceae family, including *Pyrostegia venusta*, *Jacaranda mimosifolia*, *Tabebuia* spp., and others. These plants have been traditionally used to treat various ailments, and recent studies have confirmed their antimicrobial, anti-inflammatory, and antioxidant properties. The bioactive compounds responsible for these properties include alkaloids, flavonoids, and phenolic acids. This review highlights the potential of these plants as a source of new medicinal agents and their applications in sustainable agriculture and horticulture including their ecological significances. Further research into the cultivation, bioactive compounds, and medicinal properties of these plants is warranted to unlock their full therapeutic potential. This review aims to provide a comprehensive overview of the ornamental plants of the Bignoniaceae family and their potential uses in medicine and beyond.

Keywords: Ornamental plants, bioactive compounds, medicinal properties, horticulture

Introduction

Ornamental plants have long been appreciated for their aesthetic value, but they also hold a hidden treasure trove of bioactive compounds with therapeutic applications (Schmitzer et al., 2022). Plants have been an integral part of human civilization, serving not only as a source of food and shelter but also as a repository of medicinal agents (Dimri et al., 2024; Bhat et al., 2024). For centuries, plants have been used in traditional medicine to treat various ailments, from simple fever to complex diseases like cancer (Sethi et al., 2024; Jena et al., 2024). The Bignoniaceae family, comprising over 840 species (Cordeiro et al., 2020), is a rich source of medicinal plants that have been used in traditional medicine for their antiseptic, anti-inflammatory, and antimicrobial properties (Assanti et al., 2022). The Bignoniaceae family is

also known for its ornamental value, with many species being cultivated for their showy flowers and attractive foliage. However, the medicinal properties of these ornamental plants have been largely overlooked. Recent studies have shown that plants of the Bignoniaceae family possess a wide range of bioactive compounds, including alkaloids, glycosides, and phenolic acids (Benvenuti and Mazzoncini, 2021). These compounds have been shown to exhibit various biological activities, including antimicrobial, antioxidant, and anti-inflammatory effects (Devi et al., 2024). The discovery of these bioactive compounds has sparked interest in the Bignoniaceae family as a potential source of new medicinal agents. With the rise of antibiotic resistance and the need for new treatments, plants offer a valuable resource for the discovery of novel medicinal compounds. The Bignoniaceae family holds great promise as a source of bioactive compounds with medicinal properties. The common ornamental plants of Bignoniaceae family are *Crescentia cujete* (Figure 1), *Spathodea campanulata* (Figure 2), *Tecoma stans* (Figure 3), *Tabebuia rosea* (Figure 4), *Jacaranda mimosifolia* (Figure 5) etc., In this context, exploring the ornamental plants of the Bignoniaceae family as a source of bioactive compounds with therapeutic applications is of great significance (Mukherjee et al., 2024). This review aims to provide an overview of the medicinal properties of the Bignoniaceae family, with a focus on their ornamental plants, and to highlight their potential as a source of new medicinal agents including ecological services.

Some common ornamental plants of Bignoniaceae family

***Pyrostegia venusta* (Flame Vine)**

Medicinal uses: Antiseptic, anti-inflammatory, and antimicrobial properties (Kusmardiyani et al., 2021).

Bioactive compounds: Flavonoids, phenolic compounds, and alkaloids. (Coimbra et al., 2019).

Ecological significance: Attract butterflies.

***Jacaranda mimosifolia* (Jacaranda)**

Medicinal uses: Antimicrobial activity (Aguirre et al., 2020).

Bioactive compounds: Phenolic compounds and tannin (Leutcha et al., 2025).

Ecological significance: Attract small birds.

***Tabebuia* spp. (Trumpet Tree)**

Medicinal uses: Antimicrobial, anti-inflammatory, and antioxidant properties (Barrios et al., 2023).

Bioactive compounds: Phenolic compounds (Barrios-Nolasco et al., 2023).

Ecological significance: Attract butterflies and small birds.

***Campsis radicans* (Trumpet Vine)**

Medicinal uses: Anti-hyperglycemic activity (Islam et al., 2019).

Bioactive compounds: Phenolic compounds (Islam et al., 2019).

Ecological significance: Attract butterflies.

***Bignonia capreolata* (Crossvine)**

Medicinal uses: Bark is used to cure headache (Temperate Plants Database, Ken Fern. temperate.theferns.info. 2025-01-11)

Bioactive compounds: Indole alkaloids (Clark and Lund, 2012)

Ecological significance: Attract insects.

***Catalpa bignonioides* (Catalpa)**

Medicinal uses: Leaves are used to treat skin infections (Mingarro et al., 2003).

Bioactive compounds: Saponin and phenols (Mingarro et al., 2003).

Ecological significance: Attract butterflies.

***Tecoma stans* (Yellow Bells)**

Medicinal uses: Used in treating skin infections (Bakr et al., 2019).

Bioactive compounds: Flavonoids and phenolic compounds (Anand and Basavaraju, 2021).

Ecological significance: Attracts small pollinators.



Figure 1: Flower of *Crescentia cujete*



Figure 2: Flowers of *Spathodea campanulata*



Figure 3: Leaves and flowers of *Tecoma stans*



Figure 4: Flowers of *Tabebuia rosea*



Figure 5: Flowers and leaves of *Jacaranda mimosifolia*



Figure 6: Flowers of *Podranea ricasoliana*

***Podranea ricasoliana* (Pink Trumpet Vine; Figure 6)**

Medicinal uses: It has antibacterial and antioxidant potentials (Araujo et al., 2020).

Bioactive compounds: Essential oils (Araujo et al., 2020).

Ecological significance: Attracts butterflies.

***Dolichandra unguis-cati* (Cat's Claw Creeper)**

Medicinal uses: It is used in the treatment of snake bites and used as folk medicines (Calil et al., 2017).

Bioactive compounds: Chlorogenic acid (Brondani et al., 2020).

Ecological significance: Attracts butterflies.

***Crescentia cujete* (Calabash tree)**

Medicinal uses: Anti-inflammatory activities (Parvin et al., 2015).

Bioactive compounds: Flavonoids, saponins, tannins, alkaloids, cardenolides and terpenoids (Balogun and Sabiu, 2021).

Ecological significance: It attracts bats (Lima et al., 2022).

Pharmacological potential

The Bignoniaceae family exhibits significant pharmacological potential, with various species demonstrating antimicrobial, anti-inflammatory, antioxidant, and anticancer activities (Nabatanzi et al., 2020). The bioactive compounds present in these plants, including alkaloids, flavonoids, and phenolic acids, have been shown to inhibit the growth of microorganisms, reduce inflammation, and scavenge free radicals (do Nascimento et al., 2022). Additionally, some species have been found to possess antiproliferative and pro-apoptotic effects, making them potential candidates for cancer therapy (Lima et al., 2024; Ravikumara et al., 2024; Nkumah et al., 2024). Further research is needed to fully explore the pharmacological potential of the Bignoniaceae family and to develop novel therapeutic agents from these plants.

Conclusion

The Bignoniaceae family of plants offers a rich source of bioactive compounds with potential therapeutic applications. The ornamental plants of this family, such as *Pyrostegia venusta*,

Jacaranda mimosifolia, and *Tabebuia* spp., have been found to possess antimicrobial, anti-inflammatory, and antioxidant properties, making them valuable for the development of new medicines and therapies. Further research into the cultivation, bioactive compounds, and medicinal properties of these plants is warranted, as they hold great promise for addressing various health challenges and promoting sustainable agriculture and horticulture practices having ecological significances.

Future aspects

The plants of the Bignoniaceae family, particularly those mentioned earlier, hold great promise for future research and development. With the increasing demand for natural products and the need for new medicinal agents, these plants offer a rich source of bioactive compounds with potential therapeutic applications. Future research should focus on isolating and characterizing these compounds, as well as investigating their mechanisms of action and potential uses in medicine. Another important aspect of these plants is their potential for use in sustainable agriculture and horticulture. Many of these plants are ornamental and can be cultivated for their aesthetic value, providing a source of income for farmers and gardeners. Additionally, some of these plants have been shown to have pest-repellent and fertilizer properties, making them useful for organic farming practices. Further research into the cultivation and uses of these plants could lead to new opportunities for sustainable agriculture and horticulture. In the future, it is also expected that these plants will play a significant role in the development of new medicines and therapies. With the rise of antibiotic resistance and the need for new treatments, the bioactive compounds found in these plants offer a promising solution. Additionally, the use of these plants in traditional medicine provides a valuable source of knowledge and experience that can inform modern medical research. As research into these plants continues to advance, it is likely that we will see new and innovative uses for these plants in medicine and beyond.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Option 2

Author(s) hereby declares that, Meta AI is used in writing only for getting standard language not in data collection from the field and literature. The version is Meta Llama 3.1.and source is WhatsApp

References

- Aguirre-Becerra H, Pineda-Nieto SA, García-Trejo JF, Guevara-González RG, Feregrino-Pérez AA, Álvarez-Mayorga BL, Rivera Pastrana DM. (2020). Jacaranda flower (*Jacaranda mimosifolia*) as an alternative for antioxidant and antimicrobial use. *Heliyon*. 6(12):e05802. doi: 10.1016/j.heliyon.2020.e05802.
- Anand M and Basavaraju R. (2021). A review on phytochemistry and pharmacological uses of *Tecoma stans* (L.) Juss. ex Kunth. *Journal of Ethnopharmacology*. 265: 113270. Doi:10.1016/j.jep.2020.113270.
- Araujo, C.A., Nascimento, A.L.S., da Camara, C.A.G. *et al.* (2020). Composition of the Essential Oil of *Podranea ricasoliana*. *Chem Nat Compd*. 56: 551–552.
- Assanti G, Kaur R, Nizard S, Pollack-Blackwood E, Rafferty B, Priano C, Fernández Romero JA, Koroch AR. (2022). Biology, chemistry, and pharmacological activity of *Kigelia africana* (Bignoniaceae) and *Garcinia kola* (Clusiaceae) - a review. *Journal of Med Act Plants*. 11(1):1-21.
- Bakr RO, Fayed MAA, Salem MA, Hussein AS. (2019). *Tecoma stans*: Alkaloid Profile and Antimicrobial Activity. *J Pharm Bioallied Sci*. 11(4):341-347.
- Balogun FO, Sabiu S. (2021). A Review of the Phytochemistry, Ethnobotany, Toxicology, and Pharmacological Potentials of *Crescentia cujete* L. (Bignoniaceae). *Evid Based Complement Alternat Med*. 2021:6683708. doi: 10.1155/2021/6683708.
- Barrios-Nolasco A, Domínguez-López A, Miliar-García A, Cornejo-Garrido J, Jaramillo-Flores ME. Anti-Inflammatory Effect of Ethanolic Extract from *Tabebuia rosea* (Bertol.) DC., Quercetin, and Anti-Obesity Drugs in Adipose Tissue in Wistar Rats with Diet-Induced Obesity. *Molecules*. 2023 Apr 28;28(9):3801. doi: 10.3390/molecules28093801.
- Barrios-Nolasco A, Domínguez-López A, Miliar-García A, Cornejo-Garrido J and Jaramillo-Flores ME. (2023). Anti-Inflammatory Effect of Ethanolic Extract from *Tabebuia rosea* (Bertol.) DC., Quercetin, and Anti-Obesity Drugs in Adipose Tissue in Wistar Rats with Diet-Induced Obesity. *Molecules*. 28(9):3801. doi: 10.3390/molecules28093801.
- Benvenuti S, Mazzoncini M. (2021). The Biodiversity of Edible Flowers: Discovering New Tastes and New Health Benefits. *Front Plant Sci*. 11:569499. doi: 10.3389/fpls.2020.569499.

- Bhat SS, Tallur PN, Prameela HC, Thakur S, Singh B, Chauhan J, Kumar S and Singh R. (2024). Medicinal, phytochemical screening and cytotoxicity of *Pergularia daemia* (Forssk.) Chiov. African Journal of Biological Sciences. 6(Si4): 5867-5874; 10.48047/AFJBS.6.Si4.2024.5867-5874
- Brondani JC, Lima R, Machado MM, Manfron MP. (2020). Determination of phytochemical composition, cytotoxicity, genotoxicity and mutagenicity of the hydroethanolic extract of *Dolichandra unguis-cati* L. leaves in human leukocytes. Journal of Herbal Medicine. 22: 100333. [Doi:10.1016/j.hermed.2020.100333](https://doi.org/10.1016/j.hermed.2020.100333).
- Calil Brondani, J., Reginato, F. Z., da Silva Brum, E., de Souza Vencato, M., Lima Lhamas, C., Viana, C., da Rocha, M. I., de Freitas Bauermann, L., & Manfron, M. P. (2017). Evaluation of acute and subacute toxicity of hydroethanolic extract of *Dolichandra unguis-cati* L. leaves in rats. Journal of Ethnopharmacology. 202:147–153. <https://doi.org/10.1016/j.jep.2017.03.011>
- Clark T and Lund K. (2012). Presence of the Indole Alkaloid Reserpine in *Bignonia capreolata* L. International Journal of Pharmacognosy and Phytochemical Research. 4(3):89-91.
- Coimbra MC, Chagas RCR, Vilela MSP and Castro AHF. (2019). Growth, morphology and bioactive phenolic compounds production in *Pyrostegia venusta* calli. Biocatalysis and Agricultural Biotechnology. 18:101036. Doi:10.1016/j.bcab.2019.101036
- Cordeiro JMP, Kaehler M, Souza LG, Felix LP. (2020). Heterochromatin and numeric chromosome evolution in Bignoniaceae, with emphasis on the Neotropical clade *Tabebuia* alliance. Genet Mol Biol. 43(1):e20180171. doi: 10.1590/1678-4685-GMB-2018-0171.
- Devi RS, Satapathy KB, Manjula BL, Sharma BP and Kumar S. (2024). Evaluation of Ethnopharmacological Potential of *Oenanthe javanica* DC. Annals of Biology. 40(1):135-139.
- Dimri R, Das K, Devi RS, Kumar S, Kumar N and Das A. (2024). Ecological, Socioeconomic and Nutraceutical Values of *Zizania latifolia* (Kambong): A wetland Grass for Sangai deer & Human population. Indian Forester. 150 (10): 956-961.
- do Nascimento JR, de Jesus Alves Miranda A, Vieira FC, Rodrigues CDP, Vasconcelos LN, Filho JLP, Lopes ACCB, Tangerina MMP, Vilegas W, da Rocha CQ. (2022). A Review of the Phytochemistry and Pharmacological Properties of the Genus *Arrabidaea*. Pharmaceuticals. 15(6):658. <https://doi.org/10.3390/ph15060658>

- Islam, M., Jannat, T., Kuddus, M.R. et al. (2019). In vitro and in vivo evaluation of pharmacological potentials of *Campsis radicans* L. Clin Phytosci 5: 42. <https://doi.org/10.1186/s40816-019-0144-9>
- Jena N, Rout S, Devi RS and Kumar S. (2024). Phytochemical and cytotoxicity analysis of fruits of *Zanthoxylum asiaticum* (L.) Appelhans, Groppo & J.Wen: A minor fruit plant of Odisha, India. e-planet. 22(1): 62-70.
- Leutcha PB, M Hamadou, Ditchou YON, Ansari SA, Ngnoung GAA, Mujwar S, Taiga JD, Agrawal M, Nembot GM, Hamadou SB, Lannang AM and Noundou XS. (2025). Flavonoids and other constituents from *Jacaranda mimosifolia*: In vitro analysis, molecular docking, and molecular dynamic simulations of antioxidant and anti-inflammatory activities. Biomedicine & Pharmacotherapy. 182: 117768. Doi:10.1016/j.biopha.2024.117768.
- Lima KMM, Calandrini de Azevedo LF, Rissino JD, Vale VV, Costa EVS, Dolabela MF, Nagamachi CY and Pieczarka JC. (2024). Anticancer Potential and Safety Profile of β -Lapachone In Vitro. Molecules. 29(6):1395. <https://doi.org/10.3390/molecules29061395>.
- Mingarro DM, Acero N, Llinares F, Pozuelo JM, Mera AGD, Vicenten JA, Morales L, Alguacil LF and Perez C. (2003). Biological activity of extracts from *Catalpa bignonioides* Walt. (Bignoniaceae). Journal of Ethnopharmacology. 87(2–3): 163-167.
- Mukherjee S, Hebbar DR, Alam SKS, Krishnan R, Sharma BP, Mishra S and Kumar S. (2024). Medicinally Important Plants of Bignoniaceae Family and their Pharmacological Potential. African Journal of Biological Sciences. 6(Si4): 4856-4866; DOI: 10.48047/AFJBS.6.Si4.2024.4856-4866.
- Nabatanzi A, M Nkadimeng S, Lall N, Kabasa JD, J McGaw L. (2020). Ethnobotany, Phytochemistry and Pharmacological Activity of *Kigelia africana* (Lam.) Benth. (Bignoniaceae). Plants (Basel). 9(6):753. doi: 10.3390/plants9060753.
- Nkumah AO, Kehinde CT, Oluremi BB, Attah AF and Ogbole OO. (2024). Peptide-rich extracts from leaves of *Newbouldia laevis* (P. Beauv.) Seem. ex. Bureau (Bignoniaceae) with antimicrobial and brine shrimp lethality activities. Journal of Pharmacy & Bioresources. 21(2): 87-96.
- Parvin MS, Das N, Jahan N, Akhter MA, Nahar L, Islam ME. (2015). Evaluation of in vitro anti-inflammatory and antibacterial potential of *Crescentia cujete* leaves and stem bark. BMC Res Notes. 8:412. doi: 10.1186/s13104-015-1384-5.

- Ravikumara K, Kaveri TM, Harishkumar J and Umesha S. (2024). *Bignonia magnifica*: a comprehensive exploration of its morphology, phytochemistry, pharmacology and therapeutic potential. *Vegetos*. 1-16.
- SA Lima, UM Diniz and ICS Machado. (2022). A nectar oasis for urban Glossophaginae bats: Temporal resource dynamics of the chiropterophilous *Crescentia cujete* (Bignoniaceae). *Urban Forestry & Urban Greening*. 67: 127412. <https://doi.org/10.1016/j.ufug.2021.127412>.
- Schmitzer V, Choo WS and Urbanek Krajnc A. (2022). Editorial: Phytochemicals in ornamental plants: Synthesis, nutraceutical prospects and applied focus-Women in plant science series. *Front Plant Sci*. 13:1016968. doi: 10.3389/fpls.2022.1016968.
- Sethi J, Jena N and Kumar S. (2024). Medicinal Plants of Rourkela Forest Division, Odisha, India. *Asian Journal of Environment & Ecology*. 23 (12):69-84. <https://doi.org/10.9734/ajee/2024/v23i12635>.