


# Enhancing Agricultural Resilience Through Seed Priming: A Review

## ABSTRACT

Since India is an agriculture-dependent country, more than 80% of the rural population depends on agriculture and other industries for their livelihood. The world is currently struggling to increase food production per unit of land due to population growth and the impact of climate change on agriculture, among other factors. The main stages of a plant's life cycle include germination and emergence. The main problems in rice cultivation in regions with low rainfall are low yield and poor production. Farmers are more vulnerable than successful farmers because they do not have the resources to prepare seeds for planting. Good planting, on the other hand, can increase the resilience of plants, increase drought resistance, increase yields and avoid time-consuming and expensive reseeded. It is generally believed that startups can promote growth, shorten startup time, and stimulate business development. Seeds carry out metabolic processes that prevent seed spoilage, end dormancy and increase resistance to biotic and abiotic stress factors. Explains the physical and biochemical changes that genes undergo during preparation. In recent years, seed preparation has become an important tool in producing agricultural plants that can cope with many challenges. The present review shows the advantages of various seed preparation methods such as water priming, halo priming, osmotic priming, matrix priming, osmotic hardening, agronomic priming, hormonal primers, Nutrient priming, etc, for various crops along with plant germination and growth problems.

**Keywords:** Seed Priming; priming techniques; benefits; cereals; vegetable; fruit crops




Yield Improvement




Priming Techniques



Field/Crop



Priming Benefits



Priming Process

## Graphical abstract

### Introduction:

Cereals are annual grasses with long, slender stalks that are grown all over the world for their edible grain (Sarwar, 2013). Around 50% of all calories are provided by cereals, which are grown on 700 million hectares of land (Dunwell, 2014). Around 95% of the overall dietary demand is met by the main grains (Asgari *et al.*, 2017; Zargar *et al.*, 2017). As they are essential to human nourishment, vegetable and fruit plants are also in high demand everywhere. Global warming, heat stress, drought, salinity, and other abiotic stresses have also increased along with rising population, which has increased the demand for food and commodities (Maiti and Satya, 2014). As a result, the productivity of these crops is now in jeopardy. By 2050, there will be an additional 10 million people on the planet. We only have a little amount of agricultural area, and the field crops' ability to reproduce is all we have to guarantee food security (Sher *et al.*, 2019). Yet, the globe is working to increase the grain production per unit of land as a result of the worsening effects of climate change (Singer *et al.*, 2019). For a decent crop, quick seed germination and seedling development are necessary. Scientists and farmers agree that

maintaining dependable and high-quality food supply is a significant challenge (Chivasa *et al.*, 1998; Murungu *et al.*, 2004).

Seed preparation is a direct and convenient method to compare seed germination, promote germination and seed production, among other methods “(K.Ghassemi-Golezani *et al.*, 2012)”. Farmers have been removing biotic and abiotic stressors for a very long time thanks to a promising seed priming technology that provided ground-breaking media that supports close monitoring “(Sivasubramaniam *et al.*, 2011)”. In many field crops, including cereals, seed priming offers the significant benefit of promoting resilience to conditions like drought stress, heat stress, salt, etc. The outcomes of research on seed priming have clearly demonstrated the significance of priming for improved crop stand and emergence “(Nawaz *et al.*, 2013)”. There are several techniques for rejuvenating seeds and removing environmental stress, such as hydro priming, which hydrates seeds beforehand and triggers early germination activities “(Lutts *et al.*, 2016)”. Many field crops, including cereals, have benefited from priming treatments like hydro priming, which are efficient in increasing seed vigour “(Tian *et al.*, 2014; Prasad *et al.*, 2012; Bakht *et al.*, 2011)”. The effective establishment of the seedling is a result of quick and uniform root development “(Lutts *et al.*, 2016)”.

### **Seed priming:**

Before planting, a process called seed preparation requires adequate moistening of the seed to allow for metabolic processes prior to germination once radicle protection is established- (Nascimento *et al.*, 2004; Rehman *et al.*, 2011), Priming is a process that exposes seeds to different organic or inorganic substances and temperature (Kamithi *et al.*, 2016). To prevent radicle germination before planting, the seeds should be soaked in different solutions for a while under controlled conditions and then dried to normal moisture levels. In addition to preventing the negative consequences of seed spoilage (Ghassemi-Golezani *et al.*, 2012), this supports many metabolic processes and thus Improves the germination and emergence of different seed types, especially seed plants, small-seeded grass seeds and ornamental plants. species- (Tavili *et al.*, 2011). Seed preparation is considered simple, effective, inexpensive and risk-free. Seed treatment is beneficial because it has many advantages such as uniformity and quick and rapid emergence (Musa *et al.*, 1999)-planting crops, using water efficiently, improving deep roots, allowing germination in the latent period, promoting seed maturation and earlier, flowering by increasing metabolic activity and initiating reproductive development (Singh *et al.*, 2015), Improving plant

competition and abiotic stress tolerance (Elouaer *et al.*, 2013, Aghbolaghi & Seaghi 2014)- together with the protection of problematic soil-borne diseases (*Fusarium* spp., *Sclerotinia sclerotiorum*, *R. solani*, etc.).



a)



b)



c)

**Fig. 1:- a) The soaking of wheat seeds in different priming solutions, b) The emergence of seedlings of wheat up to first 20 days of sowing and c) The seedlings at 20 DAS (Kumar *et al.*, 2023)**

### **Priming techniques :**

Many priming strategies are currently being investigated to improve seed quality. Hydraulic priming, halo-priming, osmotic priming, matrix priming, osmotic hardening, on-farm priming, hormone or growth promoter priming, micronutrient seed priming (Nutripriming), and biological priming (or treating seeds with microorganisms such as *Pseudomonas aureofaciens*, *Trichoderma* (NPs) are examples of this process. Every crop needs a distinct, optimal priming

procedure. The factors that go into optimisation for each cultivar are standardised by trial and error. These parameters include the time needed for treatment, the priming or coating material, the vigour of the seeds, and the storage conditions (temperature, moisture, oxygen demand, etc.) (Selvarani *et al.*, 2011).

### 1. Hydro-priming:

Soak the seeds in hot or cold water for a while (7-14 hours) and then cure the seeds under water before spreading them into the field or other growing areas. As a result, the seeds are able to absorb moisture and gradually weaken their coating, allowing the plant to germinate quickly and easily. Although germination has started, no rootlets have formed. Cover the seeds with the primer solution and allow to dry. As a result, the number of intermediate metabolites (intermediate metabolites) involved in seed germination and the number of enzymes involved in the production of germination energy causes the seeds to constantly absorb more moisture than the oxygen in the air. Seeds should be weighed twice, banded and stored in jars at room temperature (25±2°C), depending on the application. To measure water absorption, the seeds are weighed after processing and removal of excess or excess water. The formula is as follows:

$$\text{Water imbibitions\%} = \frac{\text{Weight of seed priming} - \text{Weight of seed before priming}}{\text{Weight of seed before priming}} \times 100$$

Before spreading the wet seeds on the main field that has already been prepared, the seeds will be dried for roughly 45 hours in the bottom of the hut. In each treatment area, seeds are planted at the chosen crop's prescribed dates. Based on daily observations of static seeds, the following formula is used to determine the germination rate:

$$\text{Germination\%} = \frac{\text{Number of germinated seeds}}{\text{Number of total seed}} \times 100$$

One of the most effective, inexpensive, accessible, and ecologically friendly techniques for slowing down and promoting seed germination is hydro-priming. increases the uniformity and rate of germination. Additionally, it makes garden plant storage better. Shortly after germination,

it begins to grow quickly. enhances the emergence of seedlings and the germination process in a salty and saline environment (Srivastava *et al.*, 2022).

## 2 On farm priming:

On-farm priming of seeds means soaking the seeds in water for about 12 hours (Rehman *et al.*, 2011), 17 hours in semi-arid tropics, 8 hours in semi-arid regions or overnight in dry ground (Musa *et al.*, 1999), and sow the seeds next day.

## 3 Halo priming:

It occurs by soaking seeds in inorganic salt solutions such as sodium chloride (NaCl), potassium nitrate (KNO<sub>3</sub>), calcium chloride (CaCl<sub>2</sub>), and calcium sulfate (CaSO<sub>4</sub>). Use different types of salt to flavor the seeds. The use of germinations stimulating solutions promotes the emergence of seeds and seedlings even in adverse environmental conditions. A tolerable NaCl solution is used to treat the seeds. The results of extensive study on haloprimer regeneration include notable enhancements in seed germination, seedling emergence, and establishment, as well as the elimination of agricultural yields in salty soils as reprisal for halo-priming. When preparing seeds, sodium chloride (NaCl) and potassium chloride (KCl) were particularly effective in removing the negative effects of salt. (Srivastava *et al.*, 2022)



**Fig.2: Effect of seed priming in germination and seedling vigour of sunflower hybrid (Manonmani *et al.*, 2014)**

## 4 Osmotic priming:

Osmotic priming, also known as osmotic-priming or osmotic-conditioning, involves immersing seeds in an osmotic solution rather than pure water in shallow water. Due to the low water content of the osmotic solution, water gradually enters the seeds, increasing water absorption and promoting seed growth while preventing the production of free radicals. Inorganic salts such as sodium chloride (NaCl), potassium chloride (KCl), potassium nitrate

(KNO<sub>3</sub>), tripotassium phosphate (K<sub>3</sub>PO<sub>4</sub>), potassium dihydrogen phosphate (KH<sub>2</sub>PO<sub>4</sub>), and calcium chloride (CaCl<sub>2</sub>). Due to its unique properties, PEG is the most commonly used drug for osmotic priming. The size of the PEG molecule limits its ability to enter the seed, reducing the direction of damage to the seed and limiting the ability of the seed to penetrate. PEG has other disadvantages such as high viscosity which reduces the rate at which oxygen can diffuse into the solution. Therefore, mechanical aeration is often used during PEG installation (Srivastava *et al.*, 2022).



**Fig. 3:-** Caraway seedlings (25 days old) (A) from non-primed seeds and (B) from primed seeds with 5% PEG for 24 h. (Mirmazloum *et al.*, 2020)

### **5 Hormonal priming:**

Hormonal seed priming is a method of boosting plant growth and development by dispersing hormone-laced seeds ahead of time. Seeds take in plant growth inhibitors when hormones are being made, which eventually has a direct impact on seed metabolism. Gibberellic acid, auxin, gibberellin, kinetin, ethylene, polyamines, salicylic acid, and gibberellin are among the most often utilised priming agents. Because inorganic salts improve germination and the bounds of pure seed development, more seeds emerge and germinate. Garden and vegetable crops benefit from KNO<sub>3</sub> since it boosts production and fruit size as well as crop quality. Soybean germination and emergence are enhanced by gibberellic acid seed preparation. Cytokinins are primarily responsible for removing certain seeds from dormancy and are also employed as priming agents. (Srivastava *et al.*, 2022)

### **6 Solid Matrix priming:**

Unlike osmotic priming, solid matrix priming (SMP), also known as matrix-conditioning, employs seeds to inhibit water absorption. By managing the water content of the seeds, Solid Matrix Priming (SMP) enhances the biological control of *Trichoderma*. This enables the seed

area to be well colonised before the seeds are sown with the appropriate pH and matrix material. In order to make the mixture's water content slightly lower than what is necessary for seed germination, informed seed is mixed with an organic carrier in this technique. After being manufactured in a solid matrix, the seeds are mixed and placed in a solid water carrier for a period of time. After being taken out of the matrix solution, the seeds are carefully cleaned and dried once more. Strong media imitates the soil's natural absorption mechanism by allowing seedlings to absorb water more slowly. “(Srivastava et al., 2022)”

### **7. Nutri priming:**

Nutripriming involves soaking seeds in nutrient solutions prior to improve seed quality by increasing nutritional content. Micronutrients are necessary for plant growth because they enable photosynthesis and respiration, two vital plant functions that when impaired can diminish grain production and overall growth (Farooq *et al.*, 2012). Micronutrients can be applied in three different methods to address this issue: directly to seeds, as a soil amendment, or as a plant or leaf spray.

### **8. Bio- priming:**


Biopriming is a seed treatment that combines seed hydration (as part of the immune system) with injection into the seed (as part of seed control). Generally speaking, biopriming entails injecting microorganisms into seedlings. Bioagents with seed hydration results in seed biopriming. (Srivastava *et al.*, 2022)

### **9. Nano priming:**

Nano priming is a revolutionary technology for seed priming that employs nanoparticles (NPs) such as zinc oxide, iron oxide, titanium dioxide, silver nanoparticles, and others. Plants are unable to utilise fertilisers or nutrients that have been applied to them because they drain or degrade when exposed to light and water. Delivery of nutrients and nanoparticulate materials to plants allows for appropriate and controlled utilisation of macromolecules and nutrients at a particular spot needed to promote plant development (Alam *et al.*, 2015).



**Fig. 4:-** Tomato variety-8152 showing increased root length in halopriming compared to control and hydropriming (Maiti *et al.*, 2013)



**Fig. 5:-** Effect of different priming methods on the growth of wheat plants (Gul *et al.*, 2022)

### The following are the primary advantages of priming:

1. Priming has a constant and quick germination rate, seedling vigour, and crop establishment under adverse situations.
2. Priming is used to combat thermoodormancy. High temperatures (35° C) that limit the development of free amino acids and esterase activity required for radicle protrusion may be overcome when seeds are primed with kinetin, resulting in germination (Cantliffe *et al.*, 1984).
3. Crop production can be increased and negative impacts of stress (drought and salt) can be reduced by seed priming.
4. Age-related chromosomal damage is lessened by priming.
5. It lessens the damaging illnesses carried by the soil.
6. Low vigour seeds' viability is also improved by priming.

### Conclusion and future perspectives

Seed priming has been demonstrated to improve stand establishment, boost yield, and accelerate seedling emergence. Priming has been demonstrated to be beneficial for a variety of field crops. It is the finest remedy for germination-related issues, particularly when crops are produced in unfavourable circumstances. Numerous priming methods have developed and are currently used in a wide variety of crops. It can increase germination rates and percentages as well as seedling emergence, ensuring adequate stand establishment in a variety of environmental

circumstances. Seed priming, which improves tolerance to harmful environmental circumstances, undoubtedly activates a number of metabolic pathways. The seedlings that grew from prepped seeds had uniformly early germination. Additionally, seed priming treatments enhance the general growth of seedlings and plants, thus farmers or other crop producers should practise and implement the science of seed priming to maximise crop production in any challenging environmental conditions.

**Table 1:- Response to various seed priming strategies in various crops:**

S No.	Crop	Priming Technique	Results	Reference
1.	Wheat	Hormonal-priming	Reduces germination time and boosts fresh and dry weight	Afzal et al., 2006
		Hydro-priming	Increases germination rate, fasters growth and higher grain yield	Rajpar et al., 2006
		Seed priming	Enhances germination, crop-stand establishment, tillering, grain and straw yields, and the harvest index.	Farooq et al., 2008
		Osmo-priming (PEG)	Improves the seedling stand establishment parameters	Salehzade et al., 2009
		Hormonal-priming	Increases emergence count	Sharma et al., 2010
		KH <sub>2</sub> PO <sub>4</sub> and KCl	Enhances germination, emergence, growth and grain yield	Yari et al., 2010
		Hydro-priming	Improves germination indices, seedling growth and crop establishment	Meena et al., 2013
		Halo-priming	Increases speed of emergence, seedling length, dry weight	Kumar et al., 2023
2.	Maize	Potassium Nitrate	Increases germination rate, yield, harvest index and tassel weight.	Vazirimehr et al., 2014
		Hydro- priming	Recorded highest grain yield, produces more panicles and 1000 grain weight	Mahajan et al., 2011
		Nutrient seed priming	Enhances seedling growth and nutritional content in maize seeds	Imran et al., 2013
		1 % ZnSO <sub>4</sub>	Increases not only plant growth but also ultimate grain production and seed Zn content in plants cultivated in soil with low Zn availability.	Harris et al., 2007
		Chitosan	Improves germination and seedling development when exposed to low temperatures.	Guan et al., 2009
3.	Rice	Hydro-priming	Produces more panicles per meter square, filled grain per panicle, 1000 grain weight and spikelet sterility.	Mahajan et al., 2011
		Nitrate salts (Mg (NO <sub>3</sub> ) <sub>2</sub> and KNO <sub>3</sub> )	Increases plant height, leaf area, and number of leaves, as well as yield-related features.	Srivastava and Bose, 2012
		Ascorbic acid @ 200 ppm and	Because of its antioxidant capability, it improves rice germination under heat stress conditions.	Kata et al., 2014

		Salicylic acid @ 50 ppm		
4.	Sorghum	Halo-priming	Accelerates emergence, increases plant height	Shehzad et al., 2012
		Osmo-priming/ PEG	Helpful in mitigating the unfavourable impact of low water potential on seed germination in inadequate temperature circumstances.	Patane et al., 2009
5.	Millets	Halo-priming	Improves millet germination components under salinity stress circumstances	Aghbolaghi and Sedghi, 2013
6.	Chickpea	2% SSP	Records higher seed yield and significantly higher N uptake	Patter et al., 2014
		Water	Influences plant height and nodule dry weight	Gupta and Singh, 2012
7.	Cowpea	Osmo-priming/ KNO <sub>3</sub>	Increases seed germination, plant height, and dry matter buildup.	Singh et al., 2014a
8.	Soybean	Primed seeds	Good establishment, earlier flowering, higher yield	Harris et al., 2007
		KNO <sub>3</sub>	Enhances pods per plant, seeds per plant and seed yield per plant	Golezani et al., 2011
		CaCl <sub>2</sub> . 2H <sub>2</sub> O	Improves plant height, number of branches, number of pods per plant, number of seeds per pod, seed yield	Chavan et al., 2014
		Seed priming for 0-14 hrs.	Increases speed of germination	Assefa and Hunje et al., 2011
		Hormonal-priming/ GA <sub>3</sub> @ 50 ppm	Enhances emergence, germination and speed of germination	Bassi et al., 2011
9.	French bean	Hormonal-priming/ GA <sub>3</sub>	Improves seed quality, as evidenced by increased seedling length, dry weight, seedling vigour index, germination speed, and mean germination time.	Sarika et al., 2013
		Hydro-priming	Improve plant performance	Ghassemi-Golezani et al., 2010
10.	Okra	Solid matrix-priming/ Calcium aluminium silicate	Increases seed germination, seedling vigour, mean germination time and marketable fruit yield	Sharma et al., 2013
		Hydro priming	Greatest mean stem length and diameter.	Sharma et al., 2014
11.	Tomato	KNO <sub>3</sub>	Increases the NR activity, germination time, germination rate	Lara, 2014
		Hydro priming	Uniform seedling growth	Maiti, et al., 2009
		Hormonal-priming	Increased germination by 30.56%.	Soubhagyabehera, 2016
12.	Spinach	SSP+Na <sub>2</sub> CO <sub>3</sub>	Improves emergence, germination percentage, plant height, leaf area, leaf yield, and seed weight by 100 percent.	Alam et al., 2013
		Halo priming with ZnSO <sub>4</sub> .7H <sub>2</sub> O	Improved resistance to biotic and abiotic stimuli, as well as early seedling development.	Chen, et al., 2011
13.	Brinjal	Osmo-priming	Increase the amount of leaves, plant height, fruit yield, fruit length, and	Satishkumar, et al.,

			blooming days by 50%.	2005
14.	Cucumber	Hydro priming (CaCl <sub>2</sub> )	Physiological changes include total seedling weight, dry weight, and seedling vigour index. Total soluble sugars and proteins, dehydrogenase activity, amylase activity isoenzyme, protein profile are all biochemical markers.	Pandey, et al., 2017
15.	Pumpkin	Hydro-priming	Enhance metabolic activities as well as biochemical parameters.	Pritima, et al., 2017
16.	Fennel	0.25 and 0.5 Mm of Salicylic acid	Improves germination, germination rate, seed stamina index, hypocotyl length, radical length, seedling fresh and dry weight.	Farahbakhsh, 2012
17.	Fenugreek	4 g/l NaCl	Fenugreek seeds had significant effects on total germination %, mean germination time, germination index, and coefficient of velocity.	Soughir et al., 2012
18.	Canola	NaCl	Increases total emergence and dry weight	Farhoudi and Sharifzadeh, 2006
19.	Safflower	NaCl	Total germination %, mean germination time, germination index, and coefficient of velocity of safflower seeds were all significantly affected.	Aymen and Hannachi, 2012
		5g/L NaCl and Kcl	Increases germination by 8.66% and 5.06%, respectively	Elouaer and Hannachi, 2012
20.	Wolf apple	Hydro-priming	Enhances seed germination and seedling growth	Anese et al., 2011
21.	Cumin	Osmo-priming	Enhances quick and uniform emergence and contributes to high vigour and higher yields.	Nematollahi et al., 2009
		Seed priming	Improves germination characteristics	Rahimi 2013
22.	Linola	Osmo-priming	Reduces emergence time while increasing seedling fresh and dry weights.	Rehman et al., 2014
23.	Sunflower	NaCl, KNO <sub>3</sub>	Increases germination percentage and seedlings growth	Bajehbaj 2010
24.	Marigold	GA <sub>3</sub> , NaCl	Higher germination rate, seedling establishment, plumule length	Sedghi et al., 2010

#### REFERENCE:

1. Afzal S, Nadeem A, Zahoor A, Qaiser M, 2006. Role of seed priming with zinc in improving the hybrid maize (*Zea mays*) yield. American-Eurasian J Agric Environ Sci 13: pp 301-06.
2. Aghbolaghi MA, Sedghi M, 2014. The effect of Osmo and Hormone Priming on Germination and Seed Reserve Utilization of Millet Seeds under Drought stress. Journal of Stress Physiology and Biochemistry 10(1): pp. 214-221.
3. Alam A, Amin NU, Ara N, Ali M, Ali I, 2013. Effect of various sources and durations of priming on Spinach seeds. Pak J Bot. 45: pp 773-77.

4. Alam MJ, Sultana F, Iqbal MT, 2015. Potential of iron nanoparticles to increase germination and growth of wheat seedling. *Journal of Nanoscience with Advanced Technology* 1: pp 14-20.
5. Anese S, Silva D, Davide AC, Rocha, FariaJM, Soares GCM, Matos ACB, Toorop PE, 2011. Seed priming improves endosperm weakening, germination, and subsequent seedling development of *Solanum lycopersium* St. Hil. *Seed Sci Technol* 39: pp 125-139.
6. Asgari MJ, Gharineh MH, Ziaiyan AH, Asoudar MA, 2017. Effects of water stress on wheat nitrogen use under minimum and conventional tillage systems. *Crop Respiration* 52: pp 209-17.
7. Assefa MK, Hunje R, 2011. Standardization of seed priming duration in soybean [*Glycine max* (L.) Merrill]. *Seed Res.* 39: pp 1-4.
8. Aymen EM, Hannachi C, 2012. Effects of NaCl priming duration and concentration on germination behaviour of Tunisian safflower. *Eurasian J Bio Sci* 6: pp 76-84.
9. Bajehbaj AA, 2010. The effects of NaCl priming on salt tolerance in sunflower germination and seedling grown under salinity conditions. *African J Biotech* 9: pp 1764-70.
10. Bakht J, Shafi M, Jamal Y, Sher H, 2011. Response of maize (*Zea mays* L.) to seed priming with NaCl and salinity stress. *Spanish Journal of Agricultural Research* 9(1): pp 252-261.
11. Bassi G, Sharma S, Gill BS, 2011. Pre-sowing seed treatment and quality in-vigouration in soybean {*Glycine max* (L) Merrill}. *Seed Res* 31: pp 81-84.
12. Cantliffe DJ, Fischer JM, Nell TA, 1984. Mechanism of Seed Priming in Circumventing Thermodormancy in Lettuce. *Plant Physiology* 75(2): pp 290-294.
13. Chavan NG, Bhujbal GB, Manjare MR, 2014. Effect of seed priming on field performance and seed yield of soybean [*Glycine max* (L.) Merrill] varieties. *Bioscan* 9: pp 111-14.
14. Chen K, Arora R, 2011. Dynamics of the antioxidant system during seed osmopriming, post-priming germination, and seedling establishment in Spinach (*Spinacia oleracea*). *Plant Science* 180(2): pp 212-220.

15. Chivasa W, Harris D, Chiduza C, Nyamudeza P, Mashingaidze AB, 1998. Agronomic practices, major crops and farmers' perceptions of the importance of good stand establishment in Musikavanhu Communal Area, Zimbabwe.
16. Dunwell JM, 2014. Transgenic cereals: Current status and future prospects. *Journal of Cereal Science* 59(3): pp 419–434.
17. Elouaer MA, Hannachi C, 2013. Influence of seed priming on emergence and growth of coriander (*Coriandrum sativum L.*) seedlings grown under salt stress, *Acta Agriculture Slovenica* 101(1): pp. 42-47.
18. Elouaer MA, Hannachi C, 2012. Seed priming to improve germination and seedling growth of safflower under salt stress. *Eur Asian J Bio Sci* 6: pp 76-84.
19. Farahbakhsh H, 2012. Germination and seedling growth in primed and unprimed seeds of fennel as affected by reduced water potential induced by NaCl. *Inter Res J Applied Basic Sci* 3: pp 737-44.
20. Farhoudi R, Sharifzadeh F, 2006. The effects of NaCl priming on salt tolerance in canola (*Brassica napus L.*) seedlings grown under saline conditions. *Indian J Crop Sci* 1: pp 74-78.
21. Farooq M, Basra SMA, Rehman H, Saleem BA, 2008. Seed priming enhances the performance of late sown wheat (*Triticum aestivum L.*) by improving chilling tolerance. *Journal of Agronomy & Crop Science* 194: pp 55–60.
22. Farooq MA, Wahid, Siddique KHM, 2012. Micronutrient application through seed treatments- a review. *Journal of Soil Science and Plant Nutrition* 12(1): pp 125-142.
23. Ghassemi GK, Jabbarpour S, Zehtab, Salmasi S, Mohammadi A, 2010. Response of winter rapeseed (*Brassica napus L.*) cultivars to salt priming of seeds. *African J Biotech* 5: pp 1089-94.
24. Ghassemi GK, Hosseinzadeh MA, Zehtab SS, Tourchi M, 2012. Improving field performance of aged chickpea seeds by hydro-priming under water stress. *Int J Plant Animal Environ* 2: pp 168-176.
25. Golezani KG, Jafari SF, Kolvanagh JS, 2011. Seed priming and field performance of soybean *Glycine max (L.)* in response to water limitation. *Not Bot Hort Agro Bot.* 39: pp 186-89.

26. Guan YJ, Jin HU, Wang, Xian J, Chen, Xia S, 2009. Seed priming with chitosan improve maize germination and seedling growth in relation to physiological changes under low temperature stress. *J Zhejiang Univ Sci* 10: pp 427-43.
27. Gul, S., Hussain, A., Ali, Q., Alam, I., Alshegaihi, R.M., Meng, Q., Zaman, W., Manghwar, H. and Munis, M.F.H. 2022. Hydropriming and osmotic priming induce resistance against *Aspergillus niger* in wheat (*Triticum aestivum* L.) by activating  $\beta$ -1, 3-glucanase, Chitinase and Thaumatin like protein genes. *Life*, **12**(12): 2061.
28. Gupta V, Singh M, 2012. Effect of seed priming and fungicide treatment on chickpea (*Cicer arietinum*) sown at different sowing depths in *kandi* belt of low altitude subtropical zone of Jammu. *Applied Bio Res.* 14: pp 187-92.
29. Harris D, Rashid A, Hollington A, Jasi L, Riches C, 2007. Prospects of improving maize yield with on farm seed priming. 'Sustainable Maize Production Systems for Nepal'. NARC and CIMMYT, Kathmandu, pp: 180-185.
30. Harris D, Rashid A, Miraj G, Arif M, Shah H, 2007. On-farm seed priming with zinc sulphate solution, a cost effective way to increase the maize yields of resource poor farmers. *Field Crop Res.* 110: pp 119-27.
31. Imran M, Mahmood A, Romheld V, Neumann V, 2013. Nutrient seed priming improves seedling development of maize exposed to low root zone temperatures during early growth. *Eur J Agron* 49: pp 141-148.
32. Kamithi KD, Wachira F, Kibe AM, 2016. Effects of different priming methods and priming durations on enzyme activities in germinating chickpea (*Cicerarietinum* L.). *American Journal of Natural and Applied Sciences* 1: pp A1-A9.
33. Kata LP, Bhaskaran M, Umarani R, 2014. Influence of priming treatments on stress tolerance during seed germination of rice. *Inter J Agri Enviro Biotech* 7: pp 225-32.
34. Kumar V, Reena, Bansal KK, Srivastava H, Attri M, McCarty SJ, Lalrammawii C, 2023. Seed Priming on crop establishment of late sown wheat crop.
35. Lara TS, Lira JMS, Rodrigues AC, Rakocevic M, Alvarenga AA, 2014. Potassium nitrate priming affects the activity of nitrate reductase and antioxidant enzymes in tomato germination. *J Agri Sci.* 6: pp 72-80.
36. Lutts S, Benincasa P, Wojtyla L, Kubala S, Pace R, Lechowska K, Quinet M, Garnczarska M, 2016. Seed priming: new comprehensive approaches for an old empirical technique. In

- 'New challenges in seed biology: basic translational research driving seed technology'. (Eds S Araújo, A Balestrazzi) pp. 1–47.
37. Mahajan G, Singh J, Sharma N, 2011. Enhancing the performance of direct seeded basmati rice through seed priming and nitrogen management. *Indian J Agron (oryza)* 48: pp 380-82.
  38. Maiti RK, Gupta A, Umasahankar P, Raj Kumar D, Vidyasagar P, 2009. Effect of priming on seedling vigour and growth and productivity of few vegetable species: Tomato, Chilli, Cucumber and Cabbage. *International Journal of Environment, Agriculture and Biotechnology* 2(4): pp 368-374.
  39. Maiti, R., Rajkumar, D., Jagan, M., Pramanik, K. and Vidyasagar, P. 2013. Effect of seed priming on seedling vigour and yield of tomato. *International Journal of Bio-resource and Stress Management*, 4(2): 119-125.
  40. Maiti RK, Satya P, 2014. Research advances in major cereal crops for adaptation to abiotic stresses, *GM Crops Food* 5(4): pp 259-279.
  41. Manonmani, V., Begum, M.A.J. and Jayanthi, M. 2014. Halo priming of seeds. *Research Journal of Seed Science*, 7: 1-13.
  42. Mirmazloun, I., Kiss, A., Erdelyi, E., Ladanyi, M., Nemeth, E.Z. and Radacsi, P. 2020. The effect of osmopriming on seed germination and early seedling characteristics of *Carum carvi* L. *Agriculture*, 10(4): 94.
  43. Meena RP, Sendhil R, Tripathi SC, Chander S, Chhokar RS, Sharma RK, 2013. Hydro-priming of seed improves the water use efficiency, grain yield and net economic returns of wheat under different moisture regimes. *SAARC J. Agri.* 11: pp 149-59.
  44. Murungu FS, Chiduzo C, Nyamuga P, Clark LJ, Whalley WR, Finch WE, 2004. Effects of on-farm seed priming on consecutive daily sowing occasion on the emergence and growth of maize in semi-arid Zimbabwe. *Field Crop Res* 89: pp 49-57.
  45. Musa AM, Johansen C, Kumar J, Harris D, 1999. Response of chickpea to seed priming in the High Barind Tract of Bangladesh. *Inter Chickpea and Pigeonpea Newsletter* 6: pp 20-22.
  46. Nascimento WM, Cantliffe DJ, Huber DJ, 2004. Ethylene evolution and endo- $\beta$ -mannanase activity during lettuce seed germination at high temperature. *Scientia Agricola*, 61(2): pp 156-163.

47. Nawaz J, Hussain M, Jabbar A, Nadeem GA, Sajid M, Subtain M, Shabbir I, 2013. Seed Priming A Technique. *International Journal of Agriculture and Crop Sciences*, 6 (20): pp. 1373–1381.
48. Nematollahi E, Bannayan M, Souhani DA, Ghanbari A, 2009. Hydro-priming and osmo-priming effects on (*Cuminum cyminum* L.) seed germination. *World Acad Sci Eng Tech* 57: pp 526-29.
49. Pandey P, Bhanuprakash K, 2017. Effect of seed priming on biochemical changes in fresh and aged seeds of cucumber. *Journal of Agricultural Studies* 5(3): pp 62-74.
50. Patane C, Cavallaro V, Salvatore L, 2009. Germination and radical growth in unprimed and primed seed of sweetsorghum as affected by reduced water potential in NaCl at different temperatures. *Industrial Crops Products* 30: pp 1-8.
51. Pattar, P. S., Mansur, C. P., Alagundagi, S. C., Salakinkoppa, S. R. and Yenagi, B. S. (2014). Nutrients uptake, soil nutrient status and grain yield of chickpea as influenced by phosphorus sources and seed priming. *Bioinfolet* 11: pp 470-73.
52. Prasad S, Prasad B, Singh RK, 2012. Effect of Hydro priming duration on germination and seedling vigor of rice [*Oryza sativa* L.] cv. Prasad. *Journal of Crop and Weed* 8(1): pp 65–71.
53. Pratima Pandey, Bhanuprakash K, Umesh, 2017. Effect of Seed Priming on Seed Germination and Vigour in Fresh and Aged Seeds of pumpkin. *International Journal of Environment, Agriculture and Biotechnology* 2(4): pp 2261-2264.
54. Rafi H, Dawar S, Zaki MJ, 2015. Seed Priming with extracts of *Acacia nilotica* (L.) Willd. Ex Delile and *Sapindus mukorossi* (L.) plant parts in the control of root rot fungi and growth of plants. *Pakistan Journal of Botany* 47: pp 1129-113.
55. Rahimi A, 2013. Seed priming improves the germination performance of cumin (*Cuminum cyminum* L.) under temperatures and water stress. *Industrial Crops Products* 42: pp 454-60.
56. Rajpar I, Khanif YM, Memon AA, 2006. Effect of Seed Priming on Growth and Yield of Wheat (*Triticum aestivum* L.) Under Non-Saline Conditions. *Inter J Agri Res* 1: pp 259-64.
57. Rehman HU, Maqsood S, Basra A, Farooq M, 2011. Field appraisal of seed priming to improve the growth, yield and quality of direct seeded rice. *Turk J Agric.* 35: pp 357-65.

58. Rehman HU, Nawaz MQ, Maqsood S, Ahmad, Basra, Irfan, Afzal, Azra, Yasmeen, Hassan, FU, 2014. Seed priming influence on early crop growth, phonological development and yield performance of linseed (*Linum usitatissimum* L.). J Integr Agric 13: pp 990-96.
59. Salehzade H, Shishvan MI, Ghiyasi M, Forouzin F, Siyahjani AA, 2009. Effect of seed priming on germination and seedling growth of wheat [*Triticum aestivum* (L.)]. Res J Biol. Sci 4: pp 629-31.
60. Sarika G, Basavaraju GV, Bhanuprakash K, Chaanakeshava V, Paramesh R, Radha BN, 2013. Investigation on seed viability and vigour of aged seed by priming in French bean. Veg Sci 40: pp 169-73.
61. Sarwar H, 2013. The importance of cereals (*Poaceae: Gramineae*) nutrition in human health: A review. Journal of Cereals and Oilseeds 4(3): pp 32–35.
62. Satishkumar L, 2005. Influence of presoaking seed treatment and seed pelleting on storability in brinjal (*Solanum melongena* L.). Indian Horticulture 32: pp 78-82.
63. Sedghi M, Ali N, Esmailpour B, 2010. Effect of seed priming on germination and seedling growth of two medicinal plants under salinity. J Emir Food Agric. 22: pp 130-13.
64. Selvarani K, Umarani R, 2011. Evaluation of seed priming methods to improve seed vigour of onion (*Allium cepa* cv. *aggreratum*) and carrot (*Daucus carota*). Journal of Agricultural Technology 7(3): pp 857-867.
65. Sharma AD, Rathore SVS, Srinivasan K, Tyagi RK, 2014. Comparison of various seed priming methods for seed germination, seedling vigour and fruit yield in okra (*Abelmoschus esculentus* L. Moench). Scientia Horticulturae. 165: pp 75–81.
66. Sharma S, Gupta A, Shekhar J, 2010. Growth parameters in late sown rainfed wheat as influenced by seed priming and foliar applied growth regulators. J Res, SKUAST 1: pp 63-69.
67. Shehzad M, Ayub M, Ahmad AHU, Yaseen M, 2012. Influence of priming techniques on emergence and seedling growth of forage sorghum (*Sorghum bicolor* l.). J Ani Pl Sci. 22: pp 154-58.
68. Sher A, Sarwar T, Nawaz A, Ijaz M, Sattar A, Ahmad S, 2019. Methods of Seed Priming. In M. Hasanuzzaman V. Fotopoulos (Eds), Priming and Pretreatment of Seeds and Seedlings: pp 1-11.

69. Singer SD, Foroud N, Laurie JD, 2019. Molecular Improvement of Grain: Target Traits for a Changing World.
70. Singh A, Dahiru R, Musa M, Haliru BS, 2014a. Effects of osmo-priming duration on germination, emergence and early growth of cowpea (*Vigna unguiculata* (L.) Walp.) in the Sudan savanna Nigeria. Int J Agron.
71. Singh H, Jassal, RK, Kang JS, Sandhu SS, Kang H, Grewal K, 2015. Seed Priming techniques in field crops- A review. Agricultural Reviews 36(4): pp 251-264.
72. Singh S, Bawa S, Singh S, Sharma SC, Kumar V, 2014b. Effect of seed priming with molybdenum on performance of rainfed chickpea (*Cicer arietinum* L.). J Res Punjab Agric Univ 51: pp 124-27.
73. Sivasubramaniam K, Geetha R, Sujatha K, Raja K, Sripunitha A, Selvarani R, 2011. Seed Priming: Triumphs and Tribulations. Physiology 98: pp 197–209.
74. Soleimanzadeh H, 2013. Effect of seed priming on germination and yield of corn.
75. Soubhagya B, 2016. A study on the effect of hormonal priming ( $Ga_3$ ) on seed quality parameters of solanaceous vegetables. International journal of agricultural science and research. 6(3): pp 337-348.
76. Soughir M, Elouaer MA, Cherif H, 2012. Effects of NaCl priming duration and concentration on germination behaviour of fenugreek. Albanian J Agric Sci. 11: pp 193-198.
77. Srivastava AK, Bose B, 2012. Effect of nitrate seed priming on phenology, growth rate and yield attributes in rice (*Oryza sativa* L.). Vegetos 25: pp 174-81.
78. Srivastava M, Devi SA, Konthoujam G, Lovely K, Das P, Thangjam R, 2022. Benefits of Seed priming in different vegetables. The Pharma Innovation Journal SP-11(6): pp 399-403.
79. Tavili A, Zare S, Moosavi SA, Enayati A, 2011. Effects of Seed priming on Germination characteristics of Bromus Species under Salt and Draught Conditions, American- Eurasian Journal of Agricultural and Environmental Sciences, 10(2): pp 163- 168.
80. Tian Y, Guan B, Zhou D, Yu J, Li G, Lou Y, 2014. Responses of seed germination, seedling growth, and seed yield traits to seed pretreatment in maize (*Zea mays* L.). Scientific World Journal.

81. Vazirimehr MR, Ganjali H R, Rigi K, Keshtehgar A, 2014. Effect of seed priming on quantitative traits corn. *International Journal of Plant Science*, 4: 134-40.
82. Yari L, Aghaalikani M, Khazaei F, 2010. Effect of seed priming duration and temperature on seed germination behaviour of bread wheat (*Triticum aestivum* L.). *Agric Biol Sci*.5:1.
83. Zargar M, Rebouh N, Pakina E, Gadzhikurbanov A, Lyashko M, Ortskhanov B, 2017. Impact of climate change on cereal production in the highlands of eastern Algeria. *Research on Crops* 18(4): pp 575–582.

UNDER PEER REVIEW