

First report of Banana streak virus infecting banana in Burkina Faso

ABSTRACT

Banana plants (*Musa* spp), both food and economic crops, are potential hosts for a various range of badnavirus species considered a major constraint to banana improvement and a threat to *Musa* production worldwide. A survey and sample collection were carried out in two main banana-growing regions of Burkina Faso. The samples collected were subjected to biological, serological and molecular diagnosis of the reverse transcriptase/ribonuclease H (RT/RNase H) region using Badna FP/RP primers, followed by sequence comparison with the Genbank database. Analyses confirmed the presence of BSV in all symptomatic samples tested in both regions. Amplification bands of the expected size were obtained for the symptomatic samples tested positive in the serological test. These partial RT/RNase H gene sequences shared highest nucleotide identity ranging from 85.80% to 99.05% with DBV isolates in GenBank. These results are proof of the existence of BSV in Burkina Faso. It is therefore important to undertake studies that will provide basic information on the virus for the development of effective control strategies.

INTRODUCTION

Banana (*Musa* spp) is herbaceous plant belonging to the family *Musaceae* of the genus *Musa* and is native to the tropical region of South East Asia. Viruses are important constraints to the movement and propagation of plant germplasm, especially for vegetatively propagated crops such as banana and plantain. Banana production is threatened by the Banana streak disease (BSD), and its pathogen belongs to the genus Badnavirus, family *Caulimoviridae* (Alangar *et al.*, 2016). BSV is widely distributed in the main planting areas of banana industry in Southeast Asia and Africa, and it had seriously affected the yield and quality of bananas resulted in huge economic losses (Kumar *et al.*, 2015). BSD can manifest with wide-ranging symptoms, from complete lack of visible impacts to plant death, depending on BSV isolates, host cultivars, and environmental conditions (Dahal *et al.*, 2000). However, the major symptoms of BSD are chlorotic and necrotic streaks.

In Burkina Faso, banana is produced in all agro-ecological zones and help combat food and nutritional insecurity and unemployment. From 2012 to 2014, production rose from 50.571

tonnes to 79.561 tonnes (TFB, 2015). In spite of this increase, production is struggling to cover national consumption needs. This is due to abiotic and biotic constraints including viral diseases causing drastic yield losses. Unfortunately, the viruses infecting bananas are not well documented in Burkina Faso's banana-growing areas. BSV has never been reported in banana growing areas in Burkina Faso. Though, viral diseases symptoms similar to badnavirus symptoms was observed on banana plant in many regions.

MATERIALS AND METHODS

Twenty-five leaf samples were collected from symptomatic banana plants in two main production regions, Boucle du Mouhoun and Hauts-Bassins. among those, 22 showed disease symptoms and 3 were symptomless. All samples were assayed to confirm BSV infection by Indirect Antigen Coated Plate Assay-ELISA with BSV polyclonal antisera (developed in-house at CIRAD, kindly provided by Serge GALZI). BSV was detect in all samples tested. Total DNA was extracted by the CTAB protocol (Permingeat *et al.*, 1998) from 25 samples. The fragment of RT/RNase H region was amplified by PCR using primers Badna FP (5'-ATGCCITTYGGIAARAAYGCICC-3') and Badna RP (5'-CCAYTTRCAIACISCICCCCAICC-3') (Yang *et al.*, 2003). Amplification products were sent for sequencing to Macrogen (Amsterdam, Netherlands). Sequences were compared with other viral sequences in the NCBI database using BLAST (BLAST, <http://www.ncbi.nlm.nih.gov/blast>).

RESULTS AND DISCUSSION

During the survey, plants showing symptoms of discontinuous and continuous streaks, chlorotic streaks and necrotic streaks were observed in the plantations (Fig. 1). Based on the nature of the symptoms observed, infection by BSV was suspected. These symptoms are similar to the usual BSV symptoms reported by many authors (Gayral and Iskra-Caruana, 2009; Furuya *et al.*, 2012).

All symptomatic samples were positives for serological assay but any symptomless samples were positives. An amplification product of the expected size was obtained for all 22 symptomatic sample tested positive (Fig. 2). Five PCR products were directly sequenced. These five sequences showed identity ranging from 85.80% to 99.05% with each other BSV isolates. All these facts are evidence of the existence of BSV in the banana-growing areas surveyed in Burkina Faso. The presence of BSV in bordering countries such as Côte d'Ivoire (Kouadio *et al.*, 2016), Benin (Pasberg-Gauhl *et al.*, 1996), Togo (Lockhart, 1995) and Ghana,

Nigeria (Agindotan *et al.*, 2006) has already been reported. Exchanges of plant material between these countries could be cause the spread of the virus in the sub-region.



Fig 1. Symptoms observed in the surveyed banana plantations. (A) Healthy leaves, (B) Discontinuous and discontinuous chlorotic streaks, (C) Chlorotic and necrotic streaks




Fig 2. Agarose gel electrophoresis (1%) showing PCR amplified products: lane M = 100pb DNA ladder (Solis Biodyne); lanes 1, 2 = BSV infected samples, lane T- = negative control; lane T+ = Positive control

CONCLUSION

To the best of our knowledge, this is the first report of BSV infecting banana in Burkina Faso. This report is the primary step to initiate research on the impact of the virus in banana production and germplasm exchange. Further research is needed to elucidate epidemiology and impact of the virus in banana production and germplasm exchange.

REFERENCES

1. Agindotan B., Winter S., Lesemann D., Uwaifo A., Mignouna J., Hugues J. and Thottapilly G., 2006. Diversity of Banana Streak-Inducing Viruses in Nigeria and Ghana: Twice as Many Sources Detected by Immunoelectron Microscopy (IEM) than by TAS-ELISA or IC-PCR. *African Journal of Biotechnology*, 5, 1194-1203.
2. Alangar B., Thomas H. and Ramasamy S., 2016. Badnaviruses: the current global scenario. *Viruses* 8(6): Article 177 DOI 10.3390/v8060177.
3. Alangar B., Thomas H. and Ramasamy S., 2016. Badnaviruses: the current global scenario. *Viruses* 8(6): Article 177 DOI 10.3390/v8060177.
4. Dahal G., Hughes J., Gauhl F., Pasberg-Gauhl C. and Nokoe K.S., 2000. Symptomatology and development of banana streak. a disease caused by banana streak badnavirus. under natural conditions in Ibadan. Nigeria. *Acta Horticulturae*. 540:361-375.
5. Furuya N., Suastika G. and Natsuaki K., 2012. First Report and Molecular Characterization of Exogenous Banana Streak Mysore Virus from Banana in Indonesia. *Asian Journal of Plant Pathology*, 6, 41-47. <https://doi.org/10.3923/ajppaj.2012.41.47>
6. Gayral P. and Iskra-Caruana M.L., (2009). Phylogeny of Banana Streak Virus Reveals Recent and Repetitive Endogenization in the Genome of Its Banana Host (*Musa* sp.). *Journal of Molecular Evolution*, 69, 65-80. <https://doi.org/10.1007/s00239-009-9253-2>
7. Kouadio K.T., De Clerck C., Agneroh T.A., Lassois L., Parisi O., Massart S., Lepoivre P. and Jijakli M.H., 2016. Prevalence of Viruses Infecting Plantain (*Musa* sp. AAB Genome) in the Major Growing Regions in Cote d'Ivoire. *African Journal of Agricultural Research*, 11, 4532-4541. <https://doi.org/10.5897/AJAR2016.11421>
8. Kumar P.L., Selvarajan R., Iskra-Caruana M.L., Chabannes M. and Hanna R., 2015. Biology. etiology and control of virus diseases of banana and plantain. *Advance in Virus Research*, 91:229-269.
9. Lockhart B.E., 1995. Banana Streak Badnavirus Infection in *Musa*: Epidemiology, Diagnosis, and Control. Food and Fertilizer Technology Center, Taiwan, Technical Bulletin 143, 1-11
10. Pasberg-Gauhl C., Gauhl F., Schill P., Lockhart B.E.L., Afreh-Nuamah K., Osei J.K. and Zuofa K., 1996. First Report of Banana Streak Virus in Farmers' Fields in Benin,

Ghana and Nigeria. *West African Plant Disease*, 80, 224. <https://doi.org/10.1094/PD-80-0224H>

11. Permingeat H. R., Romagnoli M.V and Vallejos R. H., 1998. A simple method for isolating high yield and quality DNA from (*Gossypium hirsutum* L.) leaves. *Plant Molecular Biology Reporter*, 16, pp 1–6.
12. TFB, 2015. Lettre de demande de renouvellement des matières premières de la banane (semences de bananiers sains produites par PIF) adressée au Ministère chargé de l'agriculture le 18/03/15. 2p
13. Yang I.C., Hafner G.J., Revill P.A., Dale J.L. and Harding R.M., 2003. Sequence diversity of South Pacific isolates of Taro bacilliform virus and the development of a PCR based diagnostic test. *Archives of Virology*, 148: 1957–1968.

UNDER PEER REVIEW