

EFFECT OF FOLIAR APPLICATION OF GROWTH RETARDANTS ON REPRODUCTIVE EFFICIENCY OF GROUNDNUT (*Arachis hypogaea* L.) GENOTYPES

ABSTRACT

A field experiment entitled “Study on foliar application of Plant growth regulators for improving reproductive efficiency and yield of Groundnut (*Arachis hypogaea* L.) genotypes” was carried out during *Rabi* (Summer season), 2023-24 at wetland farm of S. V. Agricultural College, Tirupati campus of Acharya N. G. Ranga Agricultural University, Andhra Pradesh. This experiment was laid out in randomized block design with factorial concept and replicated thrice, with four genotypes as factor-1 (G₁-Nityaharitha, G₂-Dharani, G₃-K-6, G₄-Vishista) and four plant growth regulators applied on as Factor-2 (T₁- Control + RDF of NPK, T₂- Prohexadione calcium@105g a.i/ha, T₃- Putrescine@100 ppm, T₄- Paclobutrazol@ 100 ppm applied before and at peak flowering stage of crop). The reproductive efficiency attributes *viz.*, number of flowers, pegs and pods per plant, flower to peg ratio, flower to pod ratio and length of peg were recorded and analyzed to understand the influence of plant growth regulators on groundnut. G₁ recorded highest number for pegs per plant (54.6), total pods per plant (36.8), mature pods per plant (28.5) and lowest flower to peg ratio (2.8), lowest length of peg (5.0cm). whereas G₂ showed lowest flower to pod ratio (4.3). Treatment of T₂ resulted in highest number of pegs per plant (51.8), total pods per plant (36.9), mature pods per plant (29.7), lowest flower to peg ratio (2.6), minimum length of peg (5.2 cm) lowest flower to pod ratio (3.6) and lowest number of flowers per plant (134.9) compared to the other treatments. Nityaharita and Prohexadione calcium@105g a.i/ha resulted in better reproductive efficiency in groundnut.

KEYWORDS: Groundnut, plant growth regulators, growth retardants, Maleic hydrazide, Paclobutrazol, reproductive efficiency.

INTRODUCTION

Groundnut belongs to the family fabaceae and order fabales. It is a widely cultivated annual crop that exhibits dicotyledonous behaviour and self-pollination. Groundnut is grown in tropical, sub-tropical, and warm climate zones between the latitudes of 40° N and 40° S. Groundnut is a rich source of edible oil (47-54%), high-quality protein (22-30%), starch (6-24%), cellulose (1-2%), minerals (2-3%) and calories (5-6%). Globally, Groundnut is cultivated under 327 lakh hectares, producing 539 lakh tonnes with the productivity of 1648 kg per hectare (FAOSTAT, 2021). India has 54.2 Lakh Ha area under Groundnut cultivation with a productivity of 1863 kg per hectare. In Andhra Pradesh, groundnut is cultivated in 8.23 lakh hectares with a production of 5.19 lakh tones. Groundnut seed contains 44 to 56% oil and 22 to 30% protein. Kernels are rich sources of riboflavin, thiamine, nicotinic acid and

Commented [1]: Generally this manuscript as a whole has grammatical mistakes that need correction. So I recommend to make a thorough investigation pertaining the grammatical errors.

Commented [2]: It would be good to add a commonly used season like during summer or spring just after Rabi depending on what Rabi means.

Commented [3]: Since the results obtained from the biostimulants is to be compared against the control, the control should be specified individually, like T1- control. If RDF of NPK is considered as control then you do not have to write Control + RDF of NPK, instead you can write as (T1-Control).

Commented [4]: S replaced by Z

Commented [5]: Use either of them. They can be used interchangeably

Commented [6]: You need to omit it for it is not mentioned in the abstract section.

Commented [7]: You may need to replace Lakh by thousand to make it understandable in all the parts of the manuscript where the word appears.

vitamin E and is rich in minerals like P, Ca, Mg and K and vitamins.

Reproductive efficiency of groundnut depends primarily on light absorption, assimilates production, production of viable flowers, pegs, flower to peg ratio, conversion of and pegs to filled pods (Swethasree *et al.*,2021). Putrescine is involved in various physiological processes, including growth, development and stress responses and results in increased crop yields (Deotale *et al.*,2018). Paclobutrazole is widely proven to inhibit the biosynthesis of gibberellins, which promote stem elongation, resulting in compact plants with shorter internodes with stouter stem, increasing root growth and causing better fruit set (Berova and Zlatev, 2000). Prohexadione Calcium controls excessive vegetative growth, resulting in shorter internodes and a more compact plant structure. It suppresses apical dominance, leading to increased branching and lateral bud development.

MATERIAL AND METHODS

A field experiment entitled “Study on foliar application of plant growth regulators for improving reproductive efficiency and yield of Groundnut (*Arachis hypogaea* L.) genotypes” was conducted on sandy clay loam soils of wetland farm of S.V. Agricultural College, Tirupati, Acharya N.G. Ranga Agricultural University. The experiment was laid out in a randomized block design with factorial concept and replicated thrice. The experiment consisted of 16 combinations with four groundnut genotypes *viz.*, Nityaharitha, Dharani, K-6 and Vishista and four growth regulator treatments *viz.*, control, Prohexadione Calcium @105 g a.i./ha, Putrescine @ 100ppm and Paclobutrazol @100ppm.

Pre harvest chemical treatment sprays were prepared and applied before and during flowering. In order to prepare 105 g a.i./ha Prohexadione calcium, 105 g of chemical active ingredient is dissolved in 500 lit of water and applied per hectare. In order to prepare 100 ppm of Putrescine, 100 mg of chemical was dissolved in 1 litre of water. And in order to prepare 100 ppm of Paclobutrazole, 100 mg of chemical is dissolved in 1 litre of water.

Number of flowers was computed by summing up the number of flowers produced per plant during entire flowering period. The number of pegs from the five plants tagged were counted and the average was expressed as number of pegs per plant. The number of pods from the five plants tagged were counted and the average was expressed as total number of pods per plant. The number of mature pods from the five plants tagged were counted and the average was expressed as number of mature pods per plant. It is the ratio of total number of flowers produced per plant to the number of pegs from the five plants tagged were counted and the average was expressed as flowers to peg ratio.

It is the ratio of total number of flowers produced per plant during entire flowering period to the number of pods from the five plants tagged which is calculated as flowers to pod ratio. Length of peg (cm) of the five randomly selected and tagged plants in each plot was measured from the base of

Commented [8]: The purpose and specific objectives of the study should be mentioned in the introduction. Besides this you need to specify the motive for selecting the current title, in other words, having conducted a solution oriented research, the problems encountered by the farming society that initiated the researcher to conduct the study should be mentioned.

Commented [9]: The climatic and soil conditions of the area or field where the study was conducted should be mentioned.

Commented [10]: It is good to have consistency while using the word plant growth regulators in place of growth retardants though they refer to the same idea. In the title it is referred as growth retardants.

Commented [11]: You need to avoid abbreviations if they are not described.

Commented [12]: How many days before?

Commented [13]: liters

Commented [14]: Avoid a sentence starting with “and”

Commented [15]: Instead of mentioning “the five plants” in every parameter, you can describe it only once in a way where it represents every other parameter. Better to avoid unnecessary repetition of words

Commented [16]: Total number of plants in each experimental plot? Number of rows? Spacing between rows and between plants?

the flower to the growing point of the peg using a scale which contains numerical values in centimeters (cm). statistical analysis

Commented [17]: What was the statistical package used to analyze the data? Present all the statistical parameters that need to be displayed.

RESULTS AND DISCUSSION

The number of flowers per plant of groundnut was significantly influenced by the genotypes. Among the genotypes the of study, the highest (158.8) and lowest (142.3) number of flowers per plant of groundnut was recorded with the G₃ and G₄ respectively. The maximum and minimum number of flowers per plant was observed with T₁ (173.7) and T₂ (134.9) respectively. And among all of the interactions of genotypes with treated plant growth regulator treatments, the highest number flowers per plant (181.7) found in G₂, treated with T₁. This treatment could result in reduced vegetative growth and translocate those resources to flowers and thereby enhancing the maximum number of flowers and seed filling percentage. The application of PBZ to peanut plants at the stage of pod formation, seed yield was increased (Senoo and Isoda, 2003).

Commented [18]: The added

Commented [19]: Replaced with "were"

Commented [20]: Replaced with "from"

Commented [21]: It is better to mention the name of the treatments instead of writing T₁, G₂,..... etc

The maximum number of pegs per plant of groundnut was recorded with the G₁ (54.6) which indicates efficiency of the genotype to transform flowers into pegs which could led to better pegs production and yield. Whereas the lowest number recorded with G₃ (42.3). Among the PGRs studied, the maximum number of pegs per plant was observed with T₂ (51.8), and minimum number of pegs per plant recorded with T₁ (44.1). The maximum (57.1) and minimum (36.2) number of pegs per plant were recorded with interaction of G₁ when sprayed with T₂ and G₃ (36.2) with T₁ among all interactions. These results are in similarity with Hua *et al.*(2014) and Barman *et al.*(2017).

Commented [22]: Better to write it in full form i.e plant growth regulators.

The maximum number of pods per plant recorded with the G₁ (36.8) followed by G₂ (36.6) and T₂ (36.9) and lowest number of total pods per plant recorded with G₄ (25.6) and T₁ (21.0). G₁ applied with T₂ recorded highest (40.9) and G₄ when treated with T₁ recorded minimum number of total pods (21.0). The number of mature pods per plant of groundnut was recorded maximum with the G₁ (28.5), T₂ (29.7) and G₁ treated with T₂ (32.5). This could be attributed to increased pegs production and better assimilate translocation efficiency of G₁ which ultimately leads to greater mature pods production potential. The minimum mature pods per plant recorded with G₄ (22.6) and T₁ (20.3). These results were in accordance with Gaudin, 2022.

Commented [23]: You need to specify between which treatments there existed significant differences. In all the parameters, you need to show whether the individual effect had significant effect on every parameter or not.

The minimum of flower to peg ratio of groundnut was recorded with the G₁ (2.8), T₂ (2.6). And highest flower to peg ratio recorded in genotype G₃ (3.8), T₁ (4.0) and G₃ treated with T₁ (5.0) because of inherent genetic characteristics of genotypes.

The highest of flower to pod ratio was recorded with the G₄ (5.6), T₁ (6.3) and G₄ treated with T₁ (7.5). And lowest flower to peg ratio recorded in genotype G₂ (4.3) and T₂ (3.6).

Minimum length of peg of groundnut was recorded with the G₁ (5.0 cm) and T₂ (5.2 cm). Maximum length of peg recorded in genotype G₃ (6.0 cm), T₃ (6.2 cm) which increases plant height by cell division, cell elongation and cell differentiation due to induction of auxins and

cytokinin. By treatment with TIBA, peg length inhibited by 25.08% than the control due to reduced plant height results in pegs travel less distance to reach earth surface (Peng *et al.*, 2013).

CONCLUSION

The foliar application of Prohexadione Calcium, Paclobutrazolin at specified concentrations applied to groundnut just before flowering stage, had been proved better in reducing excessive vegetative growth and increased the flower and pod number of groundnut. The present study concludes that, the reproductive efficiency obtained with Prohexadione Calcium@ 105ga.i/ha in Groundnut, on sandy clay loam soils of Southern Agro-climatic Zone of Andhra Pradesh during Rabi season.

Commented [24]: The issue of the cultivar is missing in the conclusion

Commented [25]: The results were written on the basis of the cumulative application of the PGRS before and during flowering, however in the conclusion it is written only before flowering. So avoid contradictions. You need to maintain consistency.

REFERENCES

- Barman, M., Gunri, S.K., Puste, A.Mand Paul, S.2017. Effect of paclobutrazol on growth and yield of kharif groundnut (*Arachis hypogaea* L.). *International Journal of Agriculture, Environment and Biotechnology*. 10(4):513-518.
- Berova, Mand Zlatev, Z.2000. Physiological response and yield of paclobutrazol treated tomato plants (*Lycopersicon esculentum* Mill.). *Plant Growth Regulation*. 30(2):117-123.
- Deotale, R.D., Shinde, R.D and Meshram, S.D. 2018.Impact of Putrescine and indole-3-butyric acid in enhancing chemical-biochemical parameters and yield of groundnut.*Journal of Soils and Crops*.28(2):359-365.
- FAOSTAT, Food and Agricultural Organization of United Nations Database, 2021.
- Gaudin, A.B.2022. Novel applications in Mississippi cotton and peanut. M.Sc. (Ag.) thesis, StateUniversity, Mississippi. 33 (50): 5567.
- Hua, S., Zhang, Y., Yu, H., Lin, B., Ding, H., Zhang, D., Ren, Yand Fang, Z. 2014. Paclobutrazol application effects on plant height seed yield and carbohydrate metabolism in canola. *International Journal of Agriculture and Biology*. 16: 471-479.
- Peng, Q., Wang, H., Tong, J., Kabir, M.H., Huang, Zand Xiao, L., 2013. Effects of indole-3-acetic acid and auxin transport inhibitor on auxin distribution and development of peanut at pegging stage. *Scientia horticultrae*.162:76-81.
- Senoo, Sand Isoda, A. 2003. Effects of paclobutrazol on dry matter distribution and yield in peanut. *Plant production science*.6(1):90-94.
- Swethasree, M., Sudhakar, P., Umamahesh, V., Prathima, T and T Giridhara, K.2024. Effect of planting density on yield and architecture suitability of groundnut (*Arachis hypogaea*L.) varieties. *The Indian Journal of Agricultural Sciences*. 94(3):297-302.

UNDER PEER REVIEW




Table 1: Total Number of Flowers, Pegs and Length of Peg of Groundnut genotypes as influenced by PGRs

Treatments	Total Number of Flowers					Total Number of Pegs					Length of Peg (cm)				
	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean
G1	171.5	137.3	159.3	148.7	154.2	51.2	57.1	54.3	55.7	54.6	5.1	4.6	5.6	4.7	5.0
G2	181.7	138.8	161.8	149.2	157.9	49.3	56.7	52.9	54.9	53.5	5.7	5.2	6.1	5.3	5.6
G3	179.0	139.5	164.4	152.1	158.8	36.2	46.6	42.0	44.2	42.3	6.1	5.6	6.5	5.8	6.0
G4	162.5	124.0	144.3	138.3	142.3	39.5	46.8	43.8	45.3	43.8	5.9	5.4	6.5	5.7	5.9
Mean	173.7	134.9	157.5	147.1		44.1	51.8	48.2	50.0		5.7	5.2	6.2	5.4	
	G	T	GXT			G	T	GXT			G	T	GXT		
SE m	3.53	3.53	7.06			0.35	0.35	0.70			0.038	0.038	0.076		
CD(P=0.05)	10.201	10.201	NS			1.01	1.01	NS			0.109	0.109	NS		

Table 2: Total Number of pods per plant and mature pods per plant, Flower to Peg Ratio and Flower to Pod Ratio of Groundnut genotypes as influenced by PGRs

Treatments	Total number of pods per plant					Total number of mature pods per plant					Flower to Peg Ratio					Flower to Pod Ratio				
	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean	T1	T2	T3	T4	Mean
G1	30.7	40.9	37.0	38.7	36.8	22.3	32.5	28.7	30.3	28.5	3.4	2.4	2.9	2.7	2.8	5.7	3.4	4.4	3.9	4.4
G2	30.0	40.0	38.2	38.3	36.6	21.0	31.7	29.5	29.3	27.9	3.7	2.5	3.1	2.7	3.0	6.0	3.4	4.2	3.8	4.3
G3	29.0	38.0	36.3	37.0	35.1	20.0	29.0	27.4	28.3	26.2	5.0	3.0	3.9	3.4	3.8	5.9	3.5	4.3	3.9	4.4
G4	21.0	28.7	25.9	26.7	25.6	18.0	25.7	23.0	23.7	22.6	4.1	2.7	3.3	3.1	3.3	7.5	4.2	5.4	5.1	5.6
Mean	27.7	36.9	34.4	35.2		20.3	29.7	27.2	27.9		4.0	2.6	3.3	3.0		6.3	3.6	4.6	4.2	
	G	T	GXT			G	T	GXT			G	T	GXT			G	T	GXT		
SEm+	0.45	0.45	0.91			0.47	0.47	0.94			0.084	0.084	0.169			0.117	0.117	0.233		
CD (P=0.05)	1.31	1.31	NS			1.36	1.36	NS			0.243	0.243	NS			0.337	0.337	NS		

Commented [26]: The value of coefficient of variance should be mentioned