

Original Research Article

New chromogenic reagent for spectrophotometric determination of ethylene bis-dithiocarbamate pesticides in vegetables

ABSTRACT

Aim: The use of pesticides, mainly ethylene bis-dithiocarbamates (EBDCs) has been reported to be abusive in the North West Region (Cameroon) but due to limited analytical facilities there are very limited data on their concentrations in foods from the local milieu. This study aimed to develop a simple spectrophotometric method to monitor residue levels of EBDCs in vegetables in order to improve food safety measures.

Methodology: The technique of analysis was based on the degradation of EBDCs in acidic medium in the presence of tin (II) chloride to liberate carbon disulfide absorbed by a new chromogenic reagent (silver nitrate/diethanolamine) which was subsequently quantified spectrophotometrically.

Results: A good linearity ($R^2=0.999$), detection limit ($0.02 \mu\text{g/mL}$), limit of quantification ($20.04 \mu\text{g/mL}$) were established. The precision of the method was within the generally accepted interval for pesticides analysis. The analysis of selected vegetables revealed that all the sampled vegetables were contaminated and the EBDC residue levels of 2.96, 6.34 and 7.92 mg/kg for parsley, huckleberry and tomato, respectively exceeded the maximum residue levels recommended by EFSA (2019) and FAO/WHO (2020).

Conclusion: The developed method was simple, rapid, and easy to use for the estimation and control of residue levels of EBDC in vegetables.

Key words: carbon disulfide, chromogenic, ethylene bis-dithiocarbamates, spectrophotometry, vegetables

1. INTRODUCTION

Most economies in Africa are largely agrarian-based with about 66% of wetlands used for agriculture [1]. In Cameroon, agriculture is predominantly rural and it plays a key role in employment, gross domestic product level and poverty reduction [2]. The crop diversity is very

rich, impressive and contributes to the country's food security [3] and reduction of hunger which is one of the United Nations sustainable goals. The Western highlands of Cameroon are made up of wetlands. Wetlands are favorable for agriculture which is the most practiced activity for about 70% of the population throughout the year [4]. Agricultural activities especially crop production is one of the major means of income generation for the population of Santa in the North West Region (NWR) of Cameroon. Vegetable production represents about 50% of the total regional farming production estimated at 185,542 tons spanning throughout a surface area of 123,769 hectares [5]. Vegetables play a significant role in human nutrition, especially as sources of vitamins (C, A, B1, B6, B9, E), minerals, dietary fiber and phytochemicals. Vegetables in the daily diet have been strongly associated with improvement of gastrointestinal health, good vision, and reduced risk of heart disease, stroke, chronic diseases such as diabetes and some form of cancer [6].

The ever-increasing demand for vegetables due to their importance in human nutrition unfortunately still requires a widespread use of pesticides. Residues of pesticides such as dithiocarbamates (DTCs) contaminate crops causing food safety issues with significant health consequences in consumers (animals and humans). The major concern in the non-target organisms are their toxic effects such as interfering with the reproductive systems and fetal development as well as their capacity to cause cancer and asthma [7]. Though maximum residue levels (MRLs) of DTCs have been defined in vegetables, their residue levels in these crops may exceed MRLs due to farmers' malpractices and non-respect of good agricultural practice, habits commonly observed within our communities [8, 9]. Among DTCs/EBDCs used in Cameroon, fungicides especially Mancozeb/mancozam, maneb, pencozeb, and zineb are highly prevalent [8, 10]. Mancozam and pencozeb are non-systemic preventive fungicides that belong to the EBDC class. They form complexes with different metal ions and exit in that form. For example when a complex is formed with manganese, it is referred as Maneb, zinc (Zineb), manganese and zinc (Mancozeb or mancozam), and sodium (Nabam). EBDCs possess a carbon disulfide (CS_2) moiety in their structure. Acid hydrolysis of EBDCs releases CS_2 which can be quantified using appropriate methods [11-13].

One of the sustainable development goals of the United Nations is to provide safe food to all. Hence, there is need to screen foods to control residue levels. Accurate but sophisticated methods to determine EBDC residues as CS_2 [11,13,14] exist but they are expensive and need trained

personnel to operate while their maintenance is costly. Some chromogenic reagents [13] have been used to determine EBDC residues in food commodities but they are very expensive and not available in limited resource settings. Hence the determination of EBDC residues in vegetables is still problematic for many developing countries including Cameroon. Therefore a simple sensitive, rapid and accurate spectrophotometric method to determine mancozam and pencozeb in vegetables was proposed in this study. It is based on using a new chromogenic reagent of silver nitrate/diethanolamine that reacts with CS₂ liberated from vegetables leading to the formation of a yellow product (complex) having a maximum absorption at λ_{max} (410nm).

2. MATERIALS AND METHODS

2.1 Plant material

Experimental vegetable samples (*Solanum nigrum*, *Solanum lycopersicum* and *Petroselinum crispum*) were cultivated on one of the experimental fields of the University of Bamenda by the researchers from April to August 2022. These vegetables were used as control samples. Field vegetables were purchased from a local market in Santa, North West Region of Cameroon. A quantity of 2 kg of each sample was collected and separately packaged in polythene bags. The samples were then placed in separate compartments of a cooler containing ice blocks. Once in the laboratory the samples were manually cut into ~ 0.5 cm pieces immediately before analysis. The selection of vegetables was based on a survey conducted by the researchers on types of vegetables cultivated and commonly consumed by the inhabitants.

2.2 Apparatus and reagents

A UV-Vis Spectrophotometer UV 752 (D) PEC Medical USA was used for all the measurements. Absorbance was measured with a 10.00 mm quartz cuvette (EN ISO 9001) which matched with the cells of the spectrophotometer. A digital scale and 500 mL Soxhlet equipment were used to weigh reagents and extract carbon disulfide, respectively.

Chemicals used were of analytical grade reagents. Carbon disulfide (>99 % purity) was supplied by Sigma Aldrich, Germany. Other chemicals including silver nitrate, diethanolamine, tin (II) chloride, hydrochloric acid, and ethanol were purchased from SGC LABO (Douala, Cameroon).

2.2.1 Carbon disulfide extraction equipment

This constituted a 500 mL soxhlet apparatus. Each chopped vegetable was separately transferred into 500 mL round bottomed flask of the soxhlet apparatus initially containing 100 mL of

hydrolyzing solution (0.1 M SnCl₂/HCl). The flask was then heated at 90°C for 45 min. The condensate was collected in the thimble and later transferred into a 50 mL conical flask.

2.3 Preparation of experimental solutions

2.3.1 Silver nitrate solution and the chromogenic reagent/solution

A 0.03 M solution of silver nitrate was prepared by dissolving accurately 0.8 g of silver nitrate in 150 mL of distilled water. For the chromogenic reagent a 10 mL/150 mL chromogenic solution was prepared by diluting 10 mL of diethanolamine in 150 mL of 0.03 M silver nitrate solution.

2.3.2 Standard carbon disulfide solution

A standard stock solution was prepared by accurately measuring 0.1 mL of carbon disulfide that was diluted with 25 mL absolute ethanol to obtain 5.01 µg/mL (Stock 1). Then, working standard solutions were prepared by transferring increasing volumes of stock 1 into separate 50 mL calibrated volumetric flasks and 25 mL of ethanol added to cover the concentration range of (0-160.32 µg/mL) and the flasks were homogenized.

2.3.3 Hydrolyzing reagent

A solution of tin (II) chloride/hydrochloric acid (0.1M) was prepared by dissolving 20 g of stannous (II) chloride dihydrate in 500 mL concentrated hydrochloric acid in a 1000 mL volumetric flask and filled to the designated mark with distilled water [11]. The solution was used immediately.

2.4 Determination of wavelength (λ_{max})

The UV-VIS spectrophotometer UV 752 (D) power button was turned on and then allowed to calibrate for 15 min. About 3 mL of the chromogenic solution/reagent blank was transferred in a 10 mm quartz cuvette and the wavelength varied from 380 to 650 nm.

2.5 Optimization of experimental conditions

2.5.1 Determination of the concentration of the chromogenic solution

In order to establish the concentration of the chromogenic solution used in the study, different concentrations of silver nitrate were investigated. Accurately measured amounts of silver nitrate (0.6 to 0.9 g) were separately dissolved in distilled water in 250 mL volumetric flasks. Each amount was weight in duplicates and dissolved in 100 and 150 mL water respectively. After the preparation of each solution, 10 mL of diethanolamine were added and the absorbance of the resulting mixture was measured at 410 nm.

2.5.2 Effect of order of addition of reagents

The order of addition of reagents silver nitrate (A), diethanolamine (B), and distilled water (C) was evaluated in order to ascertain the effect on absorbance. Firstly, the order of addition was A+B+C. Secondly it was B+A+C and thirdly, C+A+B. Lastly, the chromogenic solution (D) and diethanolamine (E) were added and absorbance recorded at 410 nm.

2.5.3 Stability of the chromogenic solution

The stability of the chromogenic solution was studied over a period of time (2 h). Briefly, 2 mL of the chromogenic solution was transferred in a 10.00 mm quartz cuvette (EN ISO 9001) and absorbance measured at 410 nm. A new aliquot was obtained from the flask and placed in the spectrophotometer for determination of absorbance after every 5 min.

2.5.4 Stability of the complex

The stability of the complex was studied. A specified volume, 10 mL of the chromogenic solution was transferred in a 50 mL conical flask. Then 0.1 mL stock 1 was added and after homogenization the absorbance was measured at 410 nm at 5 min interval.

2.5.5 Quantity of sample, volume of hydrolyzing reagent, time and temperature of heating

The amount of sample, time and temperature of heating, and the volume of the hydrolyzing reagent depend on each other. Hence, the parameters were not investigated individually. This experiment was carried out using blank sample from the field. A scale was used to accurately measure 10, 15 and 20 g of parsley. Parsley was transferred separately in a 500 mL round-bottomed flask of the soxhlet equipment. Then, 0.1 mL of stock 2 was added. Thereafter 50 mL of the hydrolyzing solution were added. The flask was placed on a heating system and heated at 50 - 90 °C for 45 min to 1h. In order to measure the optical density at 410 nm, the volume of the condensate (carbon disulfide) was taken into consideration. The optical density was measured separately after heating at 50, 60, 70, 80, and 90°C for 45 min to 1 h.

2.6 Method validation

2.6.1 Calibration curve, linearity and range

In order to quantitatively analyze EBDC fungicides (mancozam and pencozeb) in the form of carbon disulfide it was necessary to establish a calibration curve. The latter was presented as a variation of absorbance which was recorded in triplicate against the concentration of carbon disulfide to prove linearity. Calibration parameters were generated from a stock solution of concentration 5.01 µg/mL. Aliquots of 20.04 - 160.32 µg/mL were transferred in separate 50 mL

volumetric flasks containing 10 mL of the chromogenic solution. Then the reading was performed at 410 nm using a UV-Vis Spectrophotometer UV 752 (D).

2.6.2 Limit of detection

The limit of detection (LOD) was established practically and theoretically. Practically it was determined by the successive gradual dilution of the minimum concentration of the standard solution that was used in the preparation of the calibration curve [15]. In this case, 0.1 mL of stock 2 was pipetted into a calibrated 50 mL conical flask containing 10 mL of the chromogenic solution. Then 2 mL ethanol solution was added sequentially and after every addition absorbance was measured at 410 nm after 5 min. Theoretically LOD was calculated according to ICH guidelines as $LOD = 3.3 \times \sigma/S$, where σ is the standard deviation of y-intercept of regression lines (standard deviation of response) and S is the slope of calibration curve [16].

2.6.3 Limit of quantification

The limit of quantification (LOQ) was determined according to the ECDGHFS guideline which states that the limit of quantification of a method is defined as the lowest level of fortification that met the validation criteria (recovery between 70 and 120% and relative standard deviation (RSD) lower than 20%) [17]. Lowest level of fortification was taken from the analytical range which was used to plot the graph of absorbance against the concentration of carbon disulfide. The lowest level of fortification that produced satisfactory results for the recovery of carbon disulfide was 20.04 $\mu\text{g/mL}$. In order to realize the recovery of carbon disulfide, 0.1 mL of corresponding to 20.04 $\mu\text{g/mL}$ of carbon disulfide were diligently mixed with 20 g of blended parsley and 100 mL of tin (II) chloride/hydrochloric acid added in a 500 mL round-bottomed flask of soxhlet equipment. The flask was placed on a heating system and then heated at 90 °C for 45 min. Absorbance was measured at 410 nm. The percent recovery was calculated as amount of carbon disulfide recovered (g) divided by the amount spiked multiplied by one hundred.

2.6.4 Precision

In other to evaluate the precision of the proposed method, stock 2 was used. Briefly, 0.1 mL of stock 2 was transferred in a 50 mL conical flask containing 10 mL of the chromogenic solution. The absorbance was measured three times during the same day (intra-day precision) and three consecutive days (inter-day precision) [9,18,19].

2.6.5 Reproducibility

The reproducibility of the method, also called the day to-day precision or intermediate precision, or the inter-day precision was assessed by performing analyses of stock 2 by two different analysts. To validate the reproducibility of the method, 1.0 mL of the stock 2 was pipetted into a 50 mL conical flask contained 10 mL of the chromogenic solution. The absorbance was measured at 410 nm in triplicate by each analyst.

2.6.6 Accuracy and specificity

The accuracy (trueness) and specificity of the proposed method were assessed by performing triplicate recovery experiments at one concentration level (20.04 $\mu\text{g/mL}$). The experiment was performed on a control sample. A 0.1 mL of carbon disulfide working standard corresponding to 20.04 $\mu\text{g/mL}$ was pipetted into a 500 mL round-bottomed flask of the soxhlet equipment containing 20 g of blended control parsley and the flask manually homogenized. Then 100 mL of the hydrolyzing mixture were added and after homogenization the mixture was heated at 90 °C for 45 min. The condensate in the thimble was collected and transferred to a 50 mL conical flask containing 10 mL of the chromogenic solution and the yellow complex formed was homogenized and the absorbance measured at 410 nm after 5 min.

2.6.7 Recovery analysis

The recovery analysis was performed on one control sample (parsley). In this case the control sample was blended and 20 g of sample was introduced in a 500 mL flask. Then 0.1 mL solution of carbon disulfide stock 2 was added and the mixture homogenized and 100 mL tin (II) chloride/ hydrochloric acid solution were added. The mixture was heated at 90°C for 45 min. Thereafter the condensate was collected and transferred in a 50 mL conical flask containing 10 mL chromogenic solution. After homogenizing the absorbance of the solution was measured three times at 410 nm after 5 min.

2.6.8 Application of the method

The developed method was applied to determine EBDC residues as carbon disulfide in some common vegetables (huckleberry, tomatoes, parsley) through chopped extraction. In summary, 20 g of each sample was separately transferred to a 500 mL flask containing 100 mL of the acid digestion solution and placed in a heating system. After heating for 45 min at 90°C the content of the flask was then transferred to a 50 mL conical flask containing 10 mL of the complexing

solution. Thereafter the flask was homogenized and the yellow complex was measured in a spectrophotometer at 410 nm after 5 min.

3. RESULTS AND DISCUSSION

3.1 Spectral characteristics

The proposed method involved the reaction of silver nitrate/diethanolamine solution with carbon disulfide hydrolyzed by hydrochloric acid in the presence of tin (II) chloride to give a yellow colored product that was measured against a blank scanned from 380-650 nm for absorbance versus wavelength. The maximum absorbance was recorded at 410 nm. However the absorbance decreased as the wavelength increased starting from 410 nm (Figure 1). Therefore, λ_{max} of 410 nm was used in all subsequent experiments.

3.2 Optimization of experimental conditions

Various parameters affected the intensity of the colored complex. Therefore, the effects of those parameters were studied and the reaction conditions were optimized.




Figure 1: Absorption spectrum of 410 nm.

3.2.1 Concentration of chromogenic compound

The effect of different concentrations (0.02 – 0.03 M) of the chromogenic compound on the absorbance after preparing separately in 100 and 150 mL of distilled water was studied. Absorbance was recorded at 410 nm against time that ranged from 0 to 20 min. The 100 mL solution was not stable. The 150 mL solution was stable for 1 h 25 min. A 0.023 M silver nitrate

solution did not affect the response of the detector. The mean values of absorbance of 0.027 M and 0.035 M, respectively did not increase with time. The absorbance of 0.035 M silver nitrate solution decreased with time. Only 0.031 M of silver nitrate in 150 mL of distilled water had absorbance values (nm) that increased with time. A 0.031 M silver nitrate solution (Table 1) was suitable for the study.

Table 1: Effect of the concentration of chromogenic compound.

Concentration of chromogenic compound (M)	Time interval (min)	Absorbance
0.023	0	0.000
	5	0.000
	10	0.000
	15	0.000

	20	0.000
	0	0.345
	5	0.341
	10	0.346
0.027	15	0.456
	20	0.462
	0	0.613
	5	0.631
0.031	10	0.634
	15	0.635
	20	0.652
	0	0.401
	5	0.347
0.035	10	0.288
	15	0.253
	20	0.231

3.2.2 Effect of order of addition of individual reagents and solutions

Firstly, 0.8 g of silver nitrate (A) pellet was measured and transferred in a clean and dry calibrated 250 mL volumetric flask. Then, 10 mL diethanolamine (B) and 150 mL of distilled water (C) were added, respectively. The measurements obtained indicated that the order of addition of the chemicals (A+B+C, or B+A+C, or C+A+B) involved in the preparation of the chromogenic solution has an effect on the sensitivity of the detector (lowering effect) and intensity of the yellow complex product formed as shown by the response of the detector. The disparity in absorbance (Table 2) could be due to the formation of different complex ions with varying colors. This order of addition (A+B+C) was not recommended.


When silver nitrate solution (D) was prepared separately and diethanolamine (E) added, the order of addition did not have an effect on the absorbance (Table 2). This could be due to the fact that the diethanolamine did not form any new complex ions with the silver ion which resulted in no significant change in the absorbance of the solution at 410 nm. Thus, the recommended addition, any order (D+E or E+D) was used throughout this work.

Table 2: Effect of order of addition of individual reagents and solutions on absorbance.

Reagents	A+B+C			B+A+C			C+A+B		
Absorbance (nm)	0.337	0.257	0.216	0.343	0.295	0.151	0.105	0.056	0.054
Mean \pm SD	0.270 \pm 0.062			0.260 \pm 0.100			0.071 \pm 0.029		
Solutions	D+E			E+D					
Absorbance (nm)	0.442	0.442	0.441	0.441	0.442	0.442			
Mean \pm SD	0.441 \pm 0.001			0.441 \pm 0.001					

3.2.3 Stability of the chromogenic solution

The stability of the chromogenic solution was studied. The results of the experiment indicated that the absorbance values were 0.000 from 0 to 80 min. There was an increase in absorbance from 85 – 100 min then constant from 105- 110 min (Figure 2). The chromogenic solution was stable for at least 1 h 20 min. The solution was stable during the first 85 min probably because there was no significant change in its properties. This could be attributed to the formation of a colorless complex between silver nitrate and diethanolamine. The increase in absorbance after 85 min can be attributed to the decomposition of the complex as the concentration of the products and reactants change over time. However, the reaction reaches equilibrium, and the rate of complex decomposition and the formation of the product become equal. It was recommended that the chromogenic solution be used within 80 min of the preparation.

**Figure 2: Stability of the chromogenic solution.**

3.2.4 Stability of the complex (colored product)

The yellow product obtained by the proposed method was studied at different time intervals in order to ascertain its stability. The results indicated that the absorbance values of the colored product were not constant within the time interval of 0-120 min. A maximum absorption was recorded after 5 min. Thereafter the absorbance decreased with time. The formation of the yellow complex begins immediately when the reactants are mixed leading to an increase in absorbance as more of the complex is formed. The peak concentration of the complex is around 5 min (Figure 3) when the reaction likely reaches its optimum absorbance. The complex starts to decompose after reaching the maximum absorbance. This phenomenon leads to a decrease in absorbance over time. Therefore it was recommended that absorbance should be measured after 5 min.




Figure 3: Stability of the complex (colored product).

3.2.5 Quantity of sample, volume of hydrolyzing reagent, time and temperature of heating

The quantity of sample, the volume of hydrolyzing reagent, time and temperature of heating were studied and results presented in Table 3. No condensate was formed with a 10 g sample after heating in a 50 mL tin (II) chloride/hydrochloric acid solution at 50 °C for 45 min. At lower temperature and smaller volume of tin (II) chloride/hydrochloric acid solution, the reaction proceeds at a slower rate due to reduced kinetic energy of the molecules involved in the reaction. Hence no condensate was formed. Aside, a black ring was formed in the heating flask when 50 ml of hydrolyzing reagent was used hence no absorbance was obtained. After 10 g of sample was used while time and volume of hydrolyzing reagent were varied from 45 – 60 min and 50 – 100

mL respectively and heating from 60 – 90 °C no condensate was formed. When 15 g sample was heated for 45 min at 90°C in 100 mL tin (II) chloride/hydrochloric acid solution there was a significant volume of the condensate. When the latter was reacted with a chromogenic solution, a white colored product was formed. Hence no absorbance was obtained since a yellow complex is detected instead of white. With 20 g sample a significant absorbance was recorded when the sample was heated at 90 °C for 45 min in 100 mL tin (II) chloride/hydrochloric acid solution. The increase in the quantity of sample, volume of tin (II) chloride/hydrochloric acid solution and temperature increased the kinetic energy of the molecules involved in the reaction. This led to an increase in the volume of the condensate formed. Hence, 20 g of sample, 100 mL hydrolyzing reagent, heating time and temperature of 45 min and 90 °C, respectively were used throughout the study.

Table 3: Effect of the amount of sample, volume of hydrolyzing reagent, heating time and temperature on the absorption.

Sample(g)	Time(min)	Concentration of CS ₂ (µg/mL)	Temperature (°C)	SnCl ₂ /HCl (mL)	Condensate (mL)	Absorbance (nm)
10	45	20.04	50	50	0	0.000
	45		60	100	0	0.000
	45		70	100	0	0.000
	60		60	100	0	0.000
	60		70	100	0	0.000
	60		90	100	0	0.000
15	45		90	100	8	0.000
20	45		50	100	0	0.000
	45		60	100	0	0.000
	45		70	100	8	0.000
	45		80	100	9	0.000
	45		90	100	17	0.056

3.2.6 Reaction mechanism

Ethylene bis-dithiocarbamates are a derivative of dithiocarbamic acid. Upon heating in tin (II) chloride/hydrochloric acid solution, the hydrochloric acid provides the proton (H⁺) which initiates the process of hydrolysis by protonating the nitrogen to release carbon disulfide. Tin (II) chloride acts as a reducing agent which stabilized the transition state during the hydrolysis process. The sulfur atoms in the released carbon disulfide form bonds with silver in the silver-diethanolamine complex leading to changes in the electronic structure of the complex.

Alterations in the d-orbital electron transition of silver ions due to the coordination of sulfur to silver results in the formation of a yellow coloration complex probably due to particular arrangement and interactions among silver, sulfur and carbon disulfide. The complex formed is measured at 410 nm. The proposed reaction is illustrated in Figure 4.




Figure 4: Proposed acid hydrolysis and complexation reactions for EBDCs forming a yellow complex.

3.3 Method validation

3.3.1 Calibration curve and quantification

Fixing all the achieved parameters in the previous parts, a series of sequential dilutions were prepared from the standard solution and a calibration curve for the variation of intensity of colored product with the concentrations of carbon disulfide was prepared over the concentration range 0.00-160.32 $\mu\text{g/mL}$.

The limit of detection (LOD) was determined through two methods (practically and theoretically). It was determined practically by the successive gradual dilution of the minimum concentration of the standard solution that was used in the preparation of the calibration graph. Practical and theoretical limits of detection were 0.02 and 0.03 $\mu\text{g/mL}$, respectively. The limit of quantification (20.04 $\mu\text{g/mL}$) was established from the lowest concentration of the standard solution that was used to spike vegetables in the recovery analysis which met the validation

criteria (recovery between 70 and 120% and RSD lower than 20%) [12]. The limit of detection obtained in this study was similar or lower than that obtained by previously reported spectrophotometric techniques for the determination of ethylene bis-dithiocarbamates in fruits and vegetables [12, 13, 20]. The Beer's law plot and analytical parameters are given in Table 4.

Table 4: Calibration parameters.

Parameter	Value
Wavelength (nm)	410
Beer's law linear range ($\mu\text{g/mL}$)	0.00 – 160.32
Molar absorptivity ($\text{L.Mol}^{-1}.\text{cm}^{-1}$)	7.3×10^{-6}
Sandell's sensitivity ($\mu\text{g}.\text{cm}^{-2}$)	5.2×10^{-2}
LOD ($\mu\text{g/mL}$)	0.02
Practical determination	0.02
Theoretical determination	0.03
LOQ ($\mu\text{g/mL}$)	20.04
Straight line equation	$\text{Abs}=0.0072x$
Slop	0.0072
Intercept	0.0
Correlation coefficient (r)	0.999

3.3.2 Accuracy, specificity and recovery analysis of the proposed method

Accuracy and specificity were determined based on the recovery of known amounts of carbon disulfide reference standard added to a control sample (parsley) at a selected concentration, $20.04 \mu\text{g/mL}$ ($20.04 \times 10^{-6} \text{ g}$) within the calibration range (0.0 – $160.32 \mu\text{g/mL}$). The accuracy and specificity were assessed as the percentage of CS_2 recovered and also expressed as relative standard deviation of replicate experiments between the measurements. The recovery and percent relative standard deviation were 84.62 % and 0.054 %, respectively (Table 5). The recovery indicates that most of the analyte added to the sample has been detected while the method produced consistent results under the same conditions. The results were quite satisfactory as the percent recovery was between 70 and 120, intervals established by European Union [21].

Recovery analysis was carried out on a single pesticide free vegetable. Exactly 20 g of parsley were spiked with $20.044 \times 10^{-6} \text{ g}$ reference CS_2 and transferred in a 500 mL flask containing the hydrolyzing solution. Then the flask was heated at 90°C for 45 min. As presented in Table 5, the percentage of CS_2 recovered was 84.62 %. The good solubility of CS_2 in tin (II) chloride/hydrochloric acid solution allowed for efficient extraction of CS_2 from parsley while the high temperature of 90°C accelerated the reaction rate, leading to a faster extraction and higher recovery. A higher sample size contributed to a good percentage recovery while 45 min allowed

for an adequate amount of time for the extraction process to take place. This result indicated that the recovery was within the general accepted interval (70 – 120 %).

Table 5: Accuracy, specificity and recovery of CS₂

Spiked (g)	Absorbance		Mean±SD	RSD	%RSD	%Recovery
	1.220	1.221	1.222	1.221±0.001		
20.04x10⁻⁶	1.221	1.221	1.222	1.221±0.0005	0.164	0.055
	1.222	1.221	1.222	1.221±0.0005		84.62

3.3.3 Precision

The precision of the proposed method was determined as intra-day and inter-day precision. Intra-day precision was evaluated in terms of repeatability and analyzed in triplicate on the same day in the morning (6 am), afternoon (12 noon) and evening (6 pm) by same analyst. As presented in Table 6, the percent relative standard deviation (%RSD) for intra-day precision was 0.168 and 0.173 in the morning and noon, respectively and 0.174 in the evening. %RSD values for the analysts were identical (0.187) in the evening but slightly lower at noon (0.174). These results demonstrate that the precision of the method was within the generally accepted interval for pesticides [21].

Table 6: Intra-day and inter-day precision of the method.

Time frame	Intra-day precision		Inter-day precision/reproducibility			
	Mean (nm)	%RSD	Day/Analyst	Mean (nm)	%RSD	
Morning	0.296	0.168	Day 1	1 st Analyst	0.275	0.181
				2 nd Analyst	0.269	0.185
Noon	0.289	0.173	Day 2	1 st Analyst	0.286	0.174
				2 nd Analyst	0.286	0.174
Evening	0.287	0.174	Day 3	1 st analyst	0.267	0.187
				2 nd analyst	0.266	0.187

3.4 Analysis of vegetable samples

The developed method was applied to determine mancozam and pencozeb residues levels in some selected vegetables including huckleberry, tomatoes, and parsley. The results presented in Table 7, showed that the vegetables were contaminated by both pesticides or related residues. Tomato (*Solanum lycopersicum*) had the highest concentration level (7.92 mg/kg) of mancozam and pencozeb residues. The concentration level of mancozam and pencozeb residues in huckleberry (*Solanum nigrum*) was 6.34 mg/kg while parsley (*Petroselinum crispum*) had the lowest level (2.96 mg/kg). For the tomatoes the environmental protection agency (EPA) and the food safety authority in the USA recommend 2.5 mg/kg (or ppm) [22] and 4 ppm of CS₂ respectively as MRLs [23]. The European Union Commission accepts 0.2 mg/kg for parsley [24]. These concentrations surpass the MRLs established by the European Commission and the Environmental protection agency. To the best of our knowledge MRLs for huckleberry do not exist. The alarming results for the contaminated vegetables may be due to agricultural malpractices such as misuse of pesticides, frequency of application and non-adherence to a minimum of two weeks for the sales of vegetables after pesticides application. The obtained results suggest exposure of the population to the carbamates pesticides.

Table 7: EBDC residue in terms of CS₂ (mg/kg) in vegetable samples.

S/N	Sample	EBDC residue level	EPA MRLs	EFSA MRLs	FAO/WHO MRLs
1	Tomato	7.916 (mg/kg)	2.5 (mg/kg)	0.03 (mg/kg)	/
2	Huckleberry	6.338 (mg/kg)	Not available	Not available	Not available
3	Parsley	2.962 (mg/kg)	/	/	0.023 (mg/kg)

4. CONCLUSION

The spectrophotometric method developed for determination of EBDCs in vegetables was satisfactory with LOD. A new chromogenic reagent (silver nitrate/diethanolamine) was successfully used to quantify EBDCs in vegetables. All the vegetables were contaminated with pesticide residue levels above the European Commission, the EPA and food safety authority of the United States of America. Although EBDC residues have been determined by different techniques, the method described here is viable, low turnaround time (5 min), simple, convenient, requires minimum sample quantity and does not require special working conditions

unlike other methods. The new chromogenic reagent can be used for routine monitoring of EBDC residues in different food samples.

Disclaimer (artificial intelligence)

Author(s) hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

Ethical approval

It is not applicable.

UNDER PEER REVIEW

REFERENCES

1. IWMI (International Water Management Institute). Working wetlands: a new approach to balancing agricultural development with environmental protection. Water policy briefing issue, Colombo, Sri Lanka; 2006
2. World Bank. Cameroon emergency food crisis response project (P177782); 2022. Accessed April 16 2023 Available: <http://www.worldbank.org/projects>
3. Tata P. I., Ngome F., Ntsomboh N.G., Ngayong J.W. Baseline survey for setting up smallholders' sustainable vegetable seed supply and distribution system in humid tropics areas of Cameroon. Consultancy report on vegetable seeds; 2013.
4. Suh F.C., Manfo T.F.P., Nantia A.E., Dionysius N.N., Fombele N.J., Suh L.D., et al. Use of agrochemicals resulted in anemia and inflammation among farmers in Fako Division, Cameroon, *Drug Chem. Toxicol.* 2024; DOI: 10.1080/01480545.2024.2429629
5. AGRI-STAT. Surfaces and crop production. In: Animal production: Year book of statistics for agricultural sector, campaign 2009 & 2010. DESA, MINADAR, Yaoundé, Cameroon.2012;17:1621
6. Keating, J.D.H., Waliyar, F., Jammadass, R.H., Moustafa, A., Andrade, M., Dreschsel, P., Hughes, J d'A., Kardivel, P. and Luther, K. Relearning old lessons for the future of food: By bread alone no longer—diversifying diets with fruits and vegetables. *Crop Sci.* 2010;50(Suppl.):51-62
7. Gilden, R. C.; Huffling, K. & Sattler B. 2010.Pesticides and Health Risks. *JOGNN*, 2010;39: 103–110.
8. Manfo, T. F. P., Moundipa P.F., Déchaud H., Tchana A.N.,Nantia E.A., Zobot M.T., and Pugeat M. Effect of agropesticides use on male reproductive function: A study on farmers in Djutitsa (Cameroon). *Environ. Toxic.* 2012;27:423–32. doi:10.1002/tox.20656
- 9.Nantia, E.A., González, D.M., Manfo, F.P.T., Sonchieu, J., Moundipa, P.F., GarcíaCampañab, A.M., and Gámiz-Gracia, L.Characterization of carbamate pesticides in natural water from Cameroon. *Anal. Let.* 2017a;50(9), 1397–1409

10. Fissou, H.Y., Kebfene, M., Ephraïm, D., Kodbesse, B., Azoukalne, M., Isreal, D.K., Clement, K.H. Perception de risques du paludismeetutilisation des moustiquaires au Tchad. *Int. J. Biol. Chem. Sci.* 2016;10(6): 2646-2654
11. Keshavayya J, Jithan K.R, Rai S.K. A new method for the estimation of dithiocarbamates using Vapodest. *Int. J. Chem. Tech. Res.* 2012;4(3), pp 867-869
12. Kavatsurwa, S.M., Kiremire, B., Wasswa, J and Mpiana, P.T. Dithiocarbamates residues level in selected vegetables from Bukavu, Democratic Republic of Congo. *J. Physic. Chem. Sci.* 2014;1(3) ISSN: 2348 – 327X. <http://scienceq.org/Journals/JPCS.php>
13. Mozzaquatro, J.O., Mello, D.C., Reginaldo, C.S.O., Raul, C.C.R., Costa, A.M., and Caldas, E.D. Dithiocarbamate residues in fruits and leaves of passion fruit (*Passifloraedulis*) from different Brazilian regions. *J. Braz. Chem. Soc.*, 2019;30(9), 1834-1840.doi.org/10.21577/0103-5053.20190091
14. Türker, A.R and Sezer, B. Indirect determination of dithiocarbamate fungicides (zineb and ferbam) in some foodstuff by flame absorption spectrometry, *Turkish J. Pharm. Sci.*, 2005;2(1) pp. 35-42.SANTE/11945/2015
15. Mohammed.J.H.AL-K and Ahamed.M.S.AL-A. New chromogenic reagent for the spectrophotometric determination of chlorpromazine HCl in aqueous solutions and pharmaceutical formulations. *Int. J. Pharm. Pharm. Sci.* 2013;5(3), 606-611
16. Miller J.N and Miller J.C. Statistics and chemometrics for analytical chemistry. 4th ed. Prentice Hall; 2005
17. ECDGHFS (European Commission Directorate General for Health and Food Safety). Guidance document on analytical quality control and method validation procedures for pesticide residues and analysis in food and feed, SANTE/11813/2017, 21 and 22 November 2017 rev.0, https://ec.europa.eu/food/sites/food/files/plant/docs/pesticides_mrl_guidelines_wrkdoc_11813.pdf.2017 Accessed April 2023.
18. Sameer A.M., Abdulrahman and Kanakapura, B. Sensitive and selective spectrophotometric determination of gabapentin in capsules using 2-nitrophenol as chromogenic agents. *Int. J. Anal. Chem.* 2011;9 pages

19. Nantia, E.A., González, D.M., García-Campañab, A.M., and Gámiz-Gracia, L. High-throughput methodology for the determination of carbamates in food supplements by UHPLC–MS/MS. *Chromatographia* 2017b;80, 63-70
20. Ahoudi H., Gnandi K., Tanouayig., Kamilou Ouro-sama K., Yorke J.C., Creppy E.E., Moesch C. Assessment of pesticides residues contents in the vegetables cultivated in urban area of Lome (southern Togo) and their risks on public health and the environment, Togo. *Int. J. Biol. Chem. Sci.* 2018;12(5): 2172-2185. Available online at <http://www.ifgdg.org>
21. EU (European Commission). Guidance document on analytical quality control and method validation procedures for pesticides residues analysis in food and feed. 2015 SANTE/11945/2015
22. EPA (Environmental Protection Agency). Mancozeb, Maneb, Metiram, and Thiram; Proposed Tolerance Actions. Federal Register, 2009;74(178), 47507-47517
23. FSA (Food Safety Authority). 2005. Pesticide MRL Database. NZ Food Safety Authority, 1-49
24. European Commission (EC). EU regulations on pesticides maximum residue limits in or on certain products. *Off. J. EU.* 2016;5(1)