

Characterization and classification of soils in semiarid region of Varathuru watershed area of Andhra Pradesh

ABSTRACT

Eight typical pedons representing Varathuru watershed of Chittoor district, viz Pachigunta (P1 and P2), Kalepalli (P3 and P4), Ramapuram (P5 and P6) and Oggavaripalli (P7 and P8) in semi-arid ecosystem were studied for morphological, physical, chemical characteristics and their classification. Results indicated that, the colour of the soils varied from dark reddish brown to strong brown with sandy loam to sandy clay loam texture. These soils were slightly acid to moderately alkaline (6.03 to 8.35) in reaction and non-saline (0.02 to 0.36 dSm⁻¹). The bulk density, water holding capacity, particle density, organic carbon (OC), cation exchange capacity (CEC) and base saturation were varied from 1.34 to 1.61 Mgm⁻³, 28.44 to 48.96%, 2.24 to 2.58 Mgm⁻³, 0.09 to 0.74 %, 3.68 to 20.32 cmol (p⁺) kg⁻¹ and 51.61 % to 91.86 %, respectively. Due to the presence of ustic soil moisture regime, isohyperthermic temperature regime and low organic matter content, the soils were classified into Typic Hapustepts (Pedons 2, 3, 4, 6 and 8), Typic Haplustalfs (Pedons 1 and 7) and Ultic Haplustalfs (pedon 5).

Key words: Characteristics, Typic Haplustepts, Typic Haplustalfs, Ultic Haplustalfs, isohyperthermic, Pedon, Watershed

1. INTRODUCTION

Utilization of soil and water resources is the major concern in these arid and semi-arid regions accounting for more than 60 per cent of area in the country. Over 75 percent of the cropped area falls in semiarid tropical region and most of the drought prone districts are concentrated in Andhra Pradesh, Maharashtra, Tamil Nadu, Karnataka and Rajasthan. Hence, management of soil resources on scientific principles is essential to maintain the present level of soil productivity and to prevent its degradation. Comprehensive use of soil rationalizes present soil usage but also shows how they can be used and managed best for agriculture. Further, food security is a challenging task faced by the populous nations like India besides huge demand for water. Hence, concept of watershed based holistic development in rainfed areas is a potential approach, which can lead to higher productivity and sustainability.

Characterization and classification of soils are important to understand the nature of soil resources and the data generated based on the profile characteristics will serve as a benchmark for monitoring and evaluating the likely changes in soil characteristics periodically under different cropping systems because of the introduction of different soil management techniques (Ravikumar *et al.* 2009). More than ever before, a renewed attention is being given to soils due to rapidly declining area for agriculture, declining soil fertility, increasing soil degradation, unsystematic land use policies and irrational and imbalanced use of inputs (Kanwar 2004). Hence, studies on characterization and classification of soils is needed to realize the concept of watershed approach successfully

Comment [U1]: distance

Comment [U2]: distance

Comment [U3]: distance

Comment [U4]: distance

Comment [U5]: distance

Comment [U6]: distance

Comment [U7]: distance

Comment [U8]: distance between words soil management techniques

Comment [U9]: Distance between words

Comment [U10]: Distance between words

Comment [U11]: Distance between words

(Sitanggang *et al.* 2006). Keeping the above facts in view, the present study was undertaken to characterize and classify the soils of Varathuru watershed of Chittoor district.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted in 2018 in Varathuru watershed of Chittoor district in Andhra Pradesh situated between 13° 16' to 13° 12' N latitudes and 79° 15' to 79° 19' E longitudes, with an average elevation ranging from 324 to 356 m msl. The total geographical area of watershed is about 2695 ha. The mean annual soil temperature was about 33.37°C. The average rainfall in the watershed was 923.48mm.

2.2 Natural Vegetation

The natural vegetation of the study area was *Acacia nilotica*, *Acacia auriculata*, *Azadirachta indica*, *Palmyra palm*, *Parthenium hysterophorus*, *Pogamia pinnata*, *Phoenix Sylvestris*, *Pterocarpus santalinus*, *Tamarindus indica*.

2.3 Survey Methods

For soil resource inventory digital cadastral map (1:10000) prepared by using Toposheet and False colour composite were used for site selection and profile excavation. Based on geomorphic-soil relationship, eight profiles were exposed and studied (Fig 1).

2.4 Soil Sampling and Preparation for The Laboratory Analysis

Morphological characteristics such as different horizons, depth, soil colour, texture were demarked in each pedon in the field and were recorded according to Soil Survey Staff (1998) for soil profile description. Sampling of each pedon was done under pedological horizons and 2-3 kg of soil was collected. Soil colour was determined by using Munsell colour chart. Particle size distribution, particle density and bulk density of the soils was determined by International pipette method (Piper 1950), Pycnometer method (Means and Parcher 1963) and core method (Blake 1965), respectively. The pH and electrical conductivity of the soils in soil: water (1:2.5) suspension was measured by glass electrode and conductivity meter, respectively (Jackson 1973). Calcium carbonate in soil samples was estimated by rapid titration method (Puri 1949). CEC was determined by neutral normal ammonium acetate extraction (Bower *et al.* 1952). Exchangeable calcium and magnesium were also determined in neutral normal ammonium acetate extract by Versanate titration method (Chopra and Kanwar 1991). The soils of study area were classified according to Keys to Soil Taxonomy (Soil Survey Staff 2014).

3. RESULTS AND DISCUSSION

Morphological characteristics: Morphological characteristics of study area are presented in Table 1. The studied pedons were moderately shallow to deep and exhibited A-B-C horizons except P2, P3 and P6 which exhibited transitional horizon (AB and BA respectively) in profile development. Horizon boundaries of the all the pedons varied from clear to diffuse in distinctness and smooth to wavy in topography. The colour of the studied pedons varied from dark reddish brown (2.5YR) to strong brown (10YR) with dominant hue of 10YR, value (2.5-5) and chroma (1-6). This variation in the soil colour was a function of textural makeup, topographic position, mineralogy, chemical composition and moisture regimes of the soil (Sujatha *et al.* 2021).

3.1 Physical Characteristics: The results on physical characteristics of soils are presented in Table 1. The texture of the studied pedons varied from sandy loam to sandy clay loam. The soils of P1, P2 and P8 were sandy clay loam in texture in all horizons, whereas pedons P3, P4, P5, P6 and P7 were sandy loam at surface horizons and sandy clay loam at sub-surface horizons. These differences were caused by topographic position, nature of parent material, *in-situ* weathering and translocation of clay and age of soils (Deepika *et al.*, 2021). The water holding capacity of the soils was higher in pedon P1 (48.96 to 46.55 %) whereas the low water holding capacity was observed in P3 (33.94 to 29.20%).

Comment [U12]: Material

Comment [U13]: Distance between words

Comment [U14]: the map of the situation study area reference to region

Comment [U15]: please indicate the local nomination of natural vegetation if possible

Comment [U16]: distance between words

Comment [U17]: distance between words

Comment [U18]: distance between words

Comment [U19]: distance between words

Comment [U20]: distance between words

Comment [U21]: distance between words

Comment [U22]: distance between words

Comment [U23]: distance between words

Comment [U24]: distance between words

Comment [U25]: distance between words

Comment [U26]: distance between words

Comment [U27]: distance between words

Comment [U28]: distance between words

Comment [U29]: distance between words

Bulk density of pedons varied from 1.34 to 1.61 Mgm^{-3} and increasing pattern in pedons 1, 3, 6 and 8 were observed with depth, which may be ascribed to progressive compaction and lower organic matter. However, P2, P4, P5 and P7 exhibited an irregular trend with depth. Similarly, increase in bulk density down the profile due to low organic matter and compaction of soil aggregates in Yerpedu mandal in Chittoor district in Andhra Pradesh (Leelavathi *et al.* 2009). The lower bulk density values in the surface horizons of pedons could be related to the structural aggregation of the soils as a result of relatively high organic matter content in surface horizons than sub-surface horizons (Meghana *et al.*, 2024).

Comment [U30]: distance between words

Comment [U31]: distance between words

The particle density and total porosity in studied pedons of the watershed were varied from 2.40 to 2.58 Mgm^{-3} and 38.52 to 65.09% respectively. P1, P2, P3 and P7 showed an increasing trend of particle density with depth. And remained more or less same. Water holding capacity of watershed was varied from 28.44 % to 48.96 %. P1, P2, P3 and P8 were showed and increasing trend of water holding capacity with depth. The remaining pedons showed no particular trend with depth. These differences in water holding capacity values were due to variation in the depth, clay, silt and organic carbon content of the pedons. (Lakshmi *et al.* 2020).

Comment [U32]: distance between words

3.2 Chemical Characteristics: Data on the chemical properties are presented in the Table 2. pH of the watershed ranged from 6.03 to 8.35, according to pH classes these were slightly acid to moderately alkaline in reaction. The P2, P3, P4 and P7 showed an increasing trend with depth whereas pedon 5, 6 and 8 which showed decreasing pattern and pedon 1 did not show any particular trend with depth. The electrical conductivity varied between 0.02 to 0.36 dSm^{-1} in the studied pedons indicating that the soils were non-saline. It may be due to excess leaching of salts and due to free drainage conditions, which favoured the removal of released bases by percolating and drainage water (Suguna *et al.*, 2023). The soils were low in organic carbon content (0.09 to 0.74%). The low organic carbon content in these pedons may be ascribed to ustic soil moisture regime, isohyperthermic temperature regime and poor vegetation (Leelavathi *et al.* 2009). Calcium carbonate content was varied from 1.55 to 8.68 % and could be due to ustic soil moisture regime, which is quite compatible for carbonate formation. Semi-arid climate and less rainfall as compared to annual evapotranspiration in the watershed and less available water for the leaching of insoluble carbonates and bicarbonates of the calcium, facilitated the accumulation of CaCO_3 in these soils.

Comment [U33]: distance between words

CEC of the soils was low (3.68 to 20.32 $\text{cmol}(\text{p}^+)\text{kg}^{-1}$) and depth wise distribution of CEC in soils of the profiles followed the pattern of clay distribution which influences, a significant positive correlation between CEC and clay ($r = 0.861^{**}$) and silt ($r = 0.511^{**}$) signifying that clay and silt were the principal factors that influenced CEC (Table 3). Contrarily, negative and significant correlation was observed between CEC and sand content ($r = -0.724^{**}$).

Comment [U34]: distance between words

Comment [U35]: distance between words

The exchangeable bases exhibited different regular and irregular trends because of topographic position. Among exchangeable cations, calcium was dominant (2.2 to 10.6 $\text{cmol}(\text{p}^+)\text{kg}^{-1}$) in all the pedons followed by magnesium (0.4 to 6.8 $\text{cmol}(\text{p}^+)\text{kg}^{-1}$), sodium (0.05 to 0.43 $\text{cmol}(\text{p}^+)\text{kg}^{-1}$) and potassium (0.03 to 0.22 $\text{cmol}(\text{p}^+)\text{kg}^{-1}$). The base saturation percentage (BSP) was ranging from 51.61 to 91.86% and high BSP was due to the occurrence of high exchangeable calcium (Vibha *et al.* 2019).

Comment [U36]: distance between words

Comment [U37]: distance between words

Comment [U38]: distance between words

Comment [U39]: distance between words

Comment [U40]: distance between words

Comment [U41]: distance between words

Comment [U42]: distance between words

Comment [U43]:

Comment [U44]: distance between words

Soil classification: The soils of the watershed had ustic soil moisture regime (rainfall 300-1000mm). The mean annual soil temperature was much higher than 22°C and the difference between mean summer and mean winter soil temperature was less than 6°C. Thus, the soil of the study area qualify for ustic soil moisture regime and isohyperthermic soil temperature regime. Based on the CEC / clay ratio for all the pedons was recorded as <0.7. Hence, mineralogy class for all the pedons was mixed.

Comment [U45]: distance between words

Comment [U46]: distance between words

Comment [U47]: distance between words

The soils of the watershed were placed under the order Alfisols and Inceptisols. P2, P3, P4, P6 and P8, which have cambic (Bw) sub-surface diagnostic horizon only were classified under Inceptisols and these pedons were grouped under Ustepts at sub-order level due to presence of

Comment [U48]: give examples

Comment [U49]: distance between words

ustic soil moisture regime and Haplustepts at greatgroup level because these pedons did not have either duripan or calcic horizon and base saturation was more than 60% at a depth between 0.25 to 0.50 m from the surface and these pedons did not exhibit any intergradation with other taxa or an extra gradation from the central concept. Hence, these pedons were logically classified as Typic Haplustepts at sub-group level (Leelavathy *et al.* 2014). P1, P5 and P7 have argillic (Bt) sub-surface diagnostic horizon which were classified under Alfisols and these pedons were grouped under Ustalfs at sub-order level due to presence of ustic soil moisture regime and Haplustalfs at great group level because these pedons did not have either duripan or calcic horizon and base saturation was more than 60% at a depth between 0.25 to 0.50 m from the surface and these pedons were classified into Typic Haplustalfs at sub-group level because of absence of lithic or paralithic contact within 100 cm from the surface horizon and absence of pumice or pumice like fragments and volcanic ash, absence of lamellae, sandy particles size except P5 which was classified under Ultic Haplustalfs at sub-group level due to the base saturation of argillic (Bt) horizon in P5 was < 75%. Based on particle size distribution, these soils were classified as fine loamy at family level.

Comment [U50]: distance between words

4. CONCLUSION:

The study indicated that there were considerable variations in the morphological, physical and chemical characteristics of the soils depending on their geomorphic position in the area. The texture of soils varied from sandy loam to sandy clay loam. Among exchangeable cations, calcium was dominant followed by magnesium, sodium and potassium. Soil classification (Inceptisols and Alfisols) revealed that sprinkler irrigation and growing of crops, viz. groundnut, red gram, oilseeds, millets, guava, custard apple and pomegranate are the best suggestions for these soils.

Comment [U51]: distance between words

Comment [U52]: distance between words

Comment [U53]: distance between words
why do you propose this kind of crops to cultivate in this study area, from all your paper you haven't mentioned the crops in the started introduction or object even as an examples

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

- Blake, G. R. (1965). Bulk density. *Methods of Soil Analysis*, American Society of Agronomy. 374-90. Madison, Wisconsin
- Bower, C.A., Reitemeyer, R.F., & Fireman, M. (1952). Exchangeable cation analysis of saline and alkaline soils. *Soil Science* 73, 251-261
- Chopra, S.L. & Kanwar, J.S. (1991). *Analytical Agricultural Chemistry* (4th Ed.). Kalyani Publishers, New Delhi.
- Deepika Devdas, Srivastava, L. K. & Mishra, V. N. (2021). Characterization and Classification of different Soil Type of Gariyaband District of Chhattisgarh. *International journal of current micro biology and applied sciences* 10(02), 1667-1677.
- Lakshmi, Ch. S. R., Sreelatha, T., Sireesha, A. & Bharatha Lakshmi, M. (2020). Characterisation and classification of sugarcane growing soils of North coastal district of Andhra Pradesh. *Journal of the Indian Society of Soil Science*. 68(4), 357-366.
- Leelavathy, G. P., Naidu, M.V.S., Giridhara Krishna, T. Ramana, K.V., Chandrika, V.

&Sudhakar, P. (2023). Land suitability for major crops in Varathuru watershed in Andhra Pradesh using RS and GIS. *Ecology, Environment & Conservation*. 29 (2),877-884

Kanwar, J. S. (2004). Address by the Guest of Honour: 69th Annual Convention of the Indian Society of Soil Science, Acharya N G Ranga Agricultural University, (ANGRAU), Hyderabad. *Journal of the Indian Society of Soil Science*.52(4), 295–6

Leelavathi, G.P., Naidu, M.V.S., Ramavatharam,N.& Sagar, G.K. (2009). Studies on genesis, classification and evaluation of soils for sustainable land use planning in Yerpedu mandal of Chittoor district, Andhra Pradesh. *Journal of the Indian Society of Soil Science*. 57, 109-120.

Means, R. E. & Parcher, J. V. (1963). *Physical Properties of Soils*. Charles E. Merrill Books Inc., Columbus, OH, 464.

Meghana, B.S., Leelavathy, G.P., Vajantha, B., Rao, G.P.& Naidu, M.V.S.(2024). Assessment of Soil Fertility Status of Rayal Cheruvu Village in Ramachandrapuram Mandal of Tirupati District, Andhra Pradesh,India. *International Journal of Plant & Soil Science*. 36(4), 287-292.

Piper C S. (1950). *Soil and Plant Analysis*. Academic Press, New York.

Puri, A. N. (1949). *Soils, their Physics and Chemistry*. Reinhold Publishing Corp.New York, USA.

Ravikumar, M. A., Patil, P. L.& Dasog, G. S. (2009). Characterization, classification and mapping of soil resources of 48A distributary of Malaprabha right bank command, Karnataka for land use planning. *Karnataka Journal of Agricultural Sciences*.22(1), 81–8

Sitanggang M., Rao, Y. S., Ahmed, N.& Mahapatra, S. K. (2006). Characterization and classification of soils in watershed area of Shikohpur, Gurgaon district, Haryana. *Journal of the Indian Society of Soil Science*.54(1), 106–10

Soil Survey Staff (2014). *Keys to Soil Taxonomy*, 14th Edition, USDA/NRCS, Washinton D.C.

Soil Survey Staff. (1998). *Keys to Soil Taxonomy*(8th ed.). Natural Resource Conservation Service, USDA, and Blacksburg, Virginia.

Sujatha, D.V., Naidu, M.V.S., Bhaskar, B.P., Subramanyam, D., Reddy, B.R.& Krishna, T.G. (2021). Relation of soil properties to landscape position: a transect study in a part of Pinneru River basin, YSR Kadapa district, Andhra Pradesh. *Arabian Journal of Geosciences*. 14, 1648.

Suguna, B., Vajantha, B., Leelavathy, G.P., Vineetha, U.& Naidu, M.V.S.(2023). Assessment of soil physical properties under major croppingsystems of SPSR Nellore District, Andhra Pradesh, India. *International Journal of Plant and Soil Science*. 35(20), 688-695.

Vibha Singh, Singh, R.P., Chandel, S.K.S.&Patel, S.K. (2019). Physico-Chemical Properties and Available N, P, K and S in Soils of Harahua Block of Varanasi district, Uttar Pradesh. *Agropedology*. 29(01), 72-75




Fig 1 Horizon differentiation of the profiles

Table 1 Morpho-physical characteristics of different pedons

Pedon No. & Horizon	Depth (m)	Horizon boundary	Colour (moist)	Sand (%)	Silt (%)	Clay (%)	Textural	Bulk density (Mg/mg3)	Particle density	Water holding capacity (%)	Pore space (%)	Volume expansion (%)	WHC/ Clay
				(2.0-0.05 mm)	(0.05-0.002 mm)	(<0.002 mm)							
Pedon 1 Fine loamy, mixed, iso hyperthermic, Typic Hapustalfs													
Ap	0.00– 0.23	c-s	2.5 YR 2.5/4	64.79	10.99	24.22	scl	1.54	2.26	48.96	65.09	11.11	2.02
Bt1	0.23– 0.54	d-w	5 YR 3/4	61.12	12.48	26.4	scl	1.45	2.37	46.55	59.51	10.35	1.76
Cr	0.54			Weathered gneiss									
Pedon 2 Fine loamy, mixed, iso hyperthermic, Typic Haplustepts													
Ap	0.00– 0.24	c-s	5 YR 4/1	63.56	11.53	24.91	scl	1.41	2.31	39.15	51.88	4.1	1.57
BA	0.24– 0.56	d-w	10 YR 5/4	66.73	9.22	24.05	scl	1.43	2.48	31.46	40.63	4.62	1.31
Cr	0.56			Weathered gneiss									
Pedon 3 Fine loamy, mixed, iso hyperthermic, Typic Haplustepts													
Ap	0.00– 0.23	c-s	10 YR 3/6	70.55	9.97	19.48	sl	1.47	2.45	33.94	44.18	7.22	1.74
AB	0.23– 0.55	d-w	10 YR 4/3	68.91	11.65	19.44	sl	1.52	2.58	29.20	41.49	3.25	1.5
Cr	0.55			Weathered gneiss									
Pedon 4 Fine loamy, mixed, iso hyperthermic, Typic Haplustepts													
AP	0.00– 0.20	d-s	7.5 YR 4/4	68.86	13.16	17.98	sl	1.5	2.5	28.44	38.52	2.43	1.58
Bw1	0.20– 0.54	d-w	7.5 YR 4/6	57.95	19.49	22.56	scl	1.49	2.40	45.49	56.3	12.78	2.02
Bw2	0.54– 0.86	d-w	10 YR 5/3	61.03	18.66	20.31	scl	1.54	2.45	39.99	53.04	8.2	1.97
Cr	0.86			Weathered gneiss									
Pedon 5 Fine loamy, mixed, iso hyperthermic, Ultic Haplustalfs													
Ap	0.00– 0.23	c-s	10 YR 3/2	71.76	10.71	17.53	sl	1.50	2.42	31.55	42.62	6.26	1.8
1Bt1	0.23– 0.31	c-s	5 YR 4/4	65.51	14.18	20.31	scl	1.34	2.5	38.81	47.55	5.36	1.91
11BC	0.31– 0.80	c-s	7.5 YR 5/6	89.21	4.03	6.76	s	1.35	2.57	48.15	39.91	2.71	7.12
111Bt2	0.80– 1.41	c-s	7.5 YR 4/4	7.84	42.87	49.29	sic	1.55	2.24	42.51	50.99	8.25	0.86
Cr	1.41			Weathered gneiss									
Pedon 6 Fine loamy, mixed, iso hyperthermic, Typic Haplustepts													
Ap	0.00– 0.17	c-s	7.5 YR 4/6	69.23	15.25	15.52	sl	1.38	2.46	31.54	45.68	3.01	2.03
BA	0.17– 0.54	d-w	5 YR 3/4	63.39	9.69	26.92	scl	1.44	2.41	34.74	44.09	5.87	1.29
Cr	0.54			Weathered gneiss									

Pedon 7 Fine loamy, mixed, iso hyperthermic, Typic Haplustalfs

Ap	0.00–0.27	c-s	5 YR 3/4	73.82	8.75	17.43	sl	1.51	2.36	29.46	41.88	3.66	1.69
Bt1	0.27–0.56	d-w	5 YR 3/4	64.77	12.22	23.01	scl	1.45	2.38	40.26	51.17	9	1.75
Bt2	0.56–0.85	d-w	5 YR 4/4	63.87	14.77	21.36	scl	1.55	2.44	36.27	49.33	7.41	1.7
Cr	0.85												

Weathered gneiss

Pedon 8 Fine loamy, mixed, iso hyperthermic, Typic Haplustepts

A1	0.00–0.24	c-s	7.5 YR 3/4	71.77	7.05	21.18	scl	1.45	2.42	38.02	50.75	5.45	1.79
Bw1	0.24–0.71	d-w	10 YR 3/6	67.98	8.04	23.98	scl	1.61	2.41	33.63	48.49	6.16	1.4
R	0.71												

Hard Granite Gneiss

Min				7.84	4.03	6.76		1.30	2.24	28.00	39.00	2.43	0.86
Max				89.20	42.90	49.29		1.60	2.58	49.00	65.00	12.78	7.12

UNDER PEER REVIEW

Table2. Chemical properties of different pedons

Pedon No. & Horizon	Depth (m)	pH (1:2.5)		EC (dS m ⁻¹)	OC (%)	CaCO ₃ (%)	CEC	Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	BSP (%)	ESP (%)
		H ₂ O	1N/KCl										
Pedon 1													
Ap	0.00–0.23	6.57	5.46	0.12	0.65	4.4	16.01	8.1	4.7	0.24	0.19	82.69	1.53
Bt1	0.23–0.54	6.03	5.24	0.1	0.18	2.98	12.14	5.1	3.8	0.19	0.11	75.86	1.61
Cr	0.54												
Weathered gneiss													
Pedon 2													
Ap	0.00–0.24	7.51	6.67	0.33	0.74	5.83	16.48	9.2	5.5	0.22	0.22	91.86	1.34
BA	0.24–0.56	8.14	7.61	0.25	0.15	7.25	14.96	6.5	3.7	0.22	0.08	70.16	1.46
Cr	0.56												
Weathered gneiss													
Pedon 3													
Ap	0.00–0.23	6.88	6.18	0.02	0.46	2.5	11.45	6.4	3.4	0.21	0.11	88.39	1.80
AB	0.23–0.55	7.18	6.15	0.19	0.09	3.45	10.38	5.2	3.2	0.25	0.08	84.14	2.44
Cr	0.55												
Weathered gneiss													
Pedon 4													
Ap	0.00–0.20	6.11	5.47	0.1	0.53	2.03	11.61	6.2	2.4	0.33	0.16	78.32	2.87
Bw1	0.20–0.54	7.14	6.41	0.08	0.37	3.93	13.83	8.2	3.6	0.24	0.22	88.61	1.72
Bw2	0.54–0.86	7.57	6.79	0.13	0.32	2.98	12.57	6.6	3.6	0.32	0.16	84.98	2.55
Cr	0.86												
Weathered gneiss													
Pedon 5													
Ap	0.00–0.23	8.12	7.34	0.36	0.44	6.3	10.06	5.2	2.6	0.19	0.17	81.09	1.89
1Bt1	0.23–0.31	8.35	7.52	0.27	0.32	8.68	12.13	6.9	2.1	0.17	0.12	76.66	1.44
11BC	0.31–0.80	8.14	7.36	0.06	0.19	5.83	3.68	2.2	0.4	0.05	0.03	72.84	1.34
111Bt2	0.80–1.41	7.92	6.56	0.21	0.12	3.45	20.32	10.6	6.8	0.25	0.14	87.56	1.22
Cr	1.41												
Weathered gneiss													
Pedon 6													
Ap	0.00–0.17	6.21	5.56	0.23	0.34	2.5	9.67	3.8	2.4	0.19	0.17	67.86	1.97
BA	0.17–0.54	6.14	5.35	0.19	0.28	5.35	17.33	5.5	4.8	0.15	0.12	61.00	0.85
Cr	0.54												
Weathered gneiss													
Pedon 7													
Ap	0.00–0.27	6.81	5.93	0.17	0.25	1.55	11.04	3.2	1.9	0.43	0.16	51.61	3.94
Bt1	0.27–0.56	6.09	5.64	0.09	0.34	2.98	15.07	5.1	2.7	0.42	0.13	75.43	2.79
Bt2	0.56–0.85	6.67	5.92	0.12	0.21	2.5	13.61	6.9	3.5	0.18	0.12	78.63	1.34
Cr	0.85												
Weathered gneiss													
Pedon 8													
A1	0.00–0.24	6.51	5.66	0.21	0.34	3.93	13.63	4.7	3.5	0.21	0.19	63.09	1.53
Bw1	0.24–0.71	6.23	5.75	0.29	0.11	6.78	15.81	6.2	3.4	0.13	0.12	62.28	0.82
R	0.71												
Hard Granite Gneiss													
Min		6.03	5.24	0.02	0.09	1.55	3.68	2.2	0.4	0.05	0.03	51.61	0.82
Max		8.35	7.61	0.36	0.74	8.68	20.32	10.6	6.8	0.43	0.22	91.86	3.94

Table.3. Correlation matrix among physico-chemical properties

Parameter	pH	EC	OC	CaCO ₃	CEC	Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	BSP	ESP	Sand	Silt	Clay
pH	1													
EC	0.97*	1												
OC	-0.08	-0.09	1											
CaCO₃	0.56*	0.62*	-0.03	1										
CEC	-0.17	-0.25	0.09	0.07	1									
Ca²⁺	0.20	0.10	0.29	0.13	0.77	1								
Mg²⁺	0.00	-0.13	0.18	-0.01	0.88*	0.82*	1							
Na⁺	-0.24	-0.25	0.18	0.57**	0.21	0.06	0.07	1						
K⁺	-0.20	-0.25	0.67*	-0.20	0.39*	0.42*	0.40*	0.36*	1					
BSP	0.45*	0.36*	0.38*	0.03	0.08	0.65*	0.42*	-0.18	0.23	1				
ESP	-0.15	-0.14	0.09	0.60**	-0.32*	-0.37*	-0.38*	0.84*	0.15	-0.23	1			
Sand	-0.12	0.03	0.16	0.12	0.72*	0.75*	0.75*	-0.17	-0.22	0.32*	0.21	1		
Silt	0.21	0.07	-0.16	-0.21	0.51*	0.67*	0.59*	0.19	0.22	0.42*	-0.08	0.95**	1	
Clay	0.01	-0.13	-0.15	-0.02	0.86*	0.75*	0.83*	0.13	0.19	0.19	0.31*	0.95**	0.79**	1

UNDER PEER REVIEW