

In-vitro* Nematicidal Activity of Different Solvent Extracts of *Solanum torvum* Fruit against *Meloidogyne incognita

ABSTRACT

Root-knot nematodes are harmful plant parasites that significantly reduce agricultural productivity, affecting about 2,000 plant species and causing 5% of global crop losses. Due to concerns about the environmental impact of chemical nematicides, plant-based alternatives are gaining attention. This study, conducted in September-October 2023 at the University of Mysore, tested four solvent extracts (aqueous, petroleum ether, ethanol, and methanol) from *Solanum torvum* fruit at varying concentrations (10-100%) for their effects on egg hatchability and juvenile mortality of *Meloidogyne incognita*. Results showed that the methanolic extract was most effective, achieving 99% inhibition of egg hatching and 100% juvenile mortality at 100% concentration. This suggests that *Solanum torvum* extract could be an eco-friendly and economical method for managing root-knot nematodes. Further research is needed to evaluate its field efficacy and to identify the active compounds responsible for its nematicidal properties.

Keywords: Root-knot nematode, *Meloidogyne incognita*, *Solanum torvum*, egg hatching, juvenile mortality.

INTRODUCTION

Plant parasitic nematodes, particularly Root-Knot Nematodes (*Meloidogyne* species), pose a significant threat to global crop plants, causing an estimated annual loss of 125 billion dollars (Chitwood, 2003). These nematodes, obligate root parasites, are responsible for substantial agricultural productivity reduction, affecting over 3,000 plant species worldwide and resulting in estimated crop losses of nearly USD 100 billion annually (Abad et al., 2003; Dejene, 2014; Hunt and Handoo, 2009). Their impact is pervasive, affecting most cultivated crop plants (Sasser and Carter, 1985), leading to root dysfunction, reduced root volume, and compromised nutrient and water uptake efficiency (Noling, 2002). In response to the environmental and health hazards associated with chemical nematicides, there is a growing interest in botanical alternatives. However, in India, efforts to utilize botanicals for Root-Knot Nematode control on crops are limited. Botanical nematicides are seen as safer alternatives with biodegradability, selective toxicity to pests, and minimal impact on non-target organisms. This study focuses on *Solanum torvum* Swartz, commonly known as Turkey berry, as a potential botanical nematicide. Distributed widely in Asia and Tropical America, *S. torvum* has been traditionally used for medicinal purposes. Rich in alkaloids, flavonoids, saponins, tannins, and glycosides, its extracts have shown antimicrobial properties (Fui, 2012; Biney et al., 2019). The objective of this research is to evaluate the nematicidal activity of various solvent extracts (Aqueous, Methanol, Ethanol, and Petroleum ether) of *S. torvum* fruit against root-knot nematodes under in-vitro conditions.

MATERIALS AND METHODS

Collection of Plant material: The experiments were carried out during September-October 2023 at the Department of Studies in Botany, Manasagangothri, University of Mysore, Mysuru, Karnataka – 570006. Fruits of *Solanum torvum* were collected during the field survey around the Mysore district of Karnataka state. Collected samples were kept in polythene bags, tagged, and brought to the laboratory, then the fruits were cut and shade dried for up to 10 days inside the laboratory.

Preparation of *Solanum torvum* fruit extracts: Shade-dried fruits of *S. torvum* are finely ground to powder with the help of a grinder and passed through a sieve. Weighed 10gm of powder in 100ml of methanol, ethanol, petroleum ether, and distilled water in conical flasks and kept in an incubation shaker for two days to prepare crude extracts treated as stock. From this stock, required concentrations viz., 10%, 25%, 50%, 75%, and 100% were prepared.

Preparation of Nematode inoculum: The culture of *M. incognita* was maintained on tomato plants (*Solanum lycopersicum* L.). The infected plants were uprooted, roots were thoroughly washed in a running tap, and then cut into 2-3 cm in length. The roots were then placed in a jar with about 300 ml of 1% Sodium hypochlorite (NaOCl) and agitated vigorously for about 3-5 minutes. The agitation in NaOCl solution dissolved the egg masses and eggs were released in the solution. The egg mass of *M. incognita* was gently picked using forceps under a stereo zoom-microscope. Egg masses were incubated for 48 hours by the modified Baermann funnel method to obtain infective second-stage juveniles. Population density of J₂ was calculated from 5 replications of one ml aliquots of inoculum suspension.

In-vitro analysis for Juvenile Mortality: Freshly hatched 150 second-stage juveniles (48 hours old) were taken in 2.5 cm diameter petri plates containing 5 ml of each treatment solution. J₂s kept in distilled water treated as control. Plates were covered with a lid and incubated at room temperature (25±2 °C) during the experiment period. Each treatment was replicated three times. Data on mortality was recorded after 12, 24, 48, and 72 hours after incubation. Mortality of the J₂ was assessed by observing the mobility of the J₂ under a stereo zoom-microscope at 60 X magnification and expressed as the percentage of the total population. The mobile and non-mobile J₂ were prodded using a fishing needle to check for mobile responses (Das et al., 2011). The percentage of mortality was calculated using the formula;

$$\% \text{ Juvenile mortality} = \frac{\text{Number of dead juveniles}}{\text{Total number of juveniles taken}} \times 100$$

In-vitro analysis for Egg Hatching: A single egg mass of *M. incognita* was kept in 2.5 cm diameter petri plates containing 5 ml solution of each treatment. Egg mass kept in distilled water was treated as a control. Each treatment was replicated three times. Petri plates were kept at room temperature (25±20 °C). The number of hatched J₂ was counted under a microscope and the solution of each treatment was replaced every counting. Data was recorded at every 24-hour interval and continued up to 72 hours. Percent egg hatching inhibition over control was calculated using the formula (Mahesha et al., 2017);

$$\% \text{ Egg hatching} = \frac{\text{Number of eggs hatched}}{\text{Total number of eggs taken}} \times 100$$

RESULT AND DISCUSSION

In-vitro analysis for Egg Hatching: Four solvent extracts viz., Aqueous, Petroleum ether, Ethanol, and Methanol of *S. torvum* fruit were selected at different concentrations were tested for egg hatchability of root-knot nematode *M. incognita* (Table 1). The different concentrations (10, 25, 50, 75, and 100%) were tested among the four solvents. The results showed a decrease in egg hatchability as increasing concentrations of the extracts. The increase in exposure period and increasing concentration also decreases the egg hatchability. The methanolic extract is the most toxic when compared to other solvent extracts. The egg hatchability has been found to decrease with increasing concentration from 10 to 100% after 72 hours of exposure time. The inhibition of egg hatching was observed in the decreasing order of Aqueous > Ethanol > Petroleum ether. Significant results were found in Methanolic extract at 75 and 100% concentration rates.

Similarly, petroleum ether extracts and ethanolic extracts were found to be more effective than aqueous extracts (Table-1). A similar result was reported by Irdani *et al.* (2023) and Azhagumurugan and Rajan (2015) in different plant leaf extracts. They also reported that decreased egg hatchability and increased juvenile mortality were found in higher concentrations of plant extracts. 99% of egg hatching was inhibited after 48 hours of exposure time was found in methanolic extract followed by petroleum ether extract (92%) and ethanolic extract (89.66%) at their 100% concentration rate after 72 hours of exposure time respectively. Aqueous extract at 50, 75, and 100% concentration rates also showed inhibitory action of egg hatching and it resulted in 49.66, 63.33, and 79.66% inhibition of egg hatching respectively (Table 1 and fig 3 to 5).

In-vitro analysis for Juvenile Mortality: The second-stage juveniles were exposed for 24-, 48- and 72-hours exposure time in different concentrations (10,25,50,75 and 100%) and the juvenile mortality was found to increase with increasing concentrations of different solvent extracts of which aqueous (78%), petroleum ether (86%), ethanol (88.33%) and methanol (100%) were found effective after 72 hours of exposure period at their highest concentration rate of 100% (Table-2 and fig- 6 to 8). The increase in juvenile mortality was found associated with an increase in concentrations of the extracts was also reported by Irdani *et al.* (2023) and Azhagumurugan and Rajan (2015), where they found the same in leaf extracts of different plants. The characteristic shape of nematodes killed by extracts of the fruits of *S. torvum* was found with straight shapes. The highest percentage of inhibition of egg hatching and juvenile mortality (99% and 100% respectively) was obtained at a higher concentration of methanolic extract at a 100% concentration rate. Similarly, applying different solvent extracts at all concentrations in different time intervals significantly inhibited egg hatching and caused the juvenile mortality of *M. incognita* compared to the control (Table 1 and 02). However, there were some variations among treatments in reducing egg hatching and juvenile mortality. Treatments applied at lower concentrations were less effective than higher concentrations in all tested treatments, a similar result was found by Terefe, (2015).



Figure - 1: *Solanum torvum* plant



Figure –2: Dried fruits of *Solanum torvum*

Treatment	Concentration	Percentage of egg hatched at different exposure time		
		24 hrs	48 hrs	72 hrs
Control	-	28.66	52.33	97.33
	10%	23.33	44.66	68
	25%	21	38.66	53.66
	50%	17.66	31.33	50.33
	75%	17	25.66	37.66
	100%	7.66	15	21.33
Petroleum ether extract	10%	12.33	28.66	58.33
	25%	12	18.33	28.33
	50%	8.66	14.33	17.66
	75%	7.33	12.33	12.33
	100%	3.66	6.33	8
Ethanol extract	10%	22.66	39.66	57.66
	25%	17.66	30	45.33
	50%	14.33	24.33	37.33
	75%	10.66	16	25.33
	100%	6	8.33	10.33
Methanol extract	10%	6.33	10.33	14
	25%	4	6.66	7.66
	50%	2.66	4.66	6.33
	75%	0.66	1	1
	100%	0.33	1	1

Table 1: Percentage of egg hatching in different solvent extracts of *Solanum torvum* fruit.

Treatment	Concentration	Percentage of juvenile mortality at different
-----------	---------------	---

		exposure time		
		24 hrs	48 hrs	72 hrs
Control	-	00	0.33	2.33
	10%	2.66	5.66	8
Aqueous extract	25%	9	15	21
	50%	16.66	28.33	41.66
	75%	24.33	39.66	53
	100%	31.33	57.66	78
	10%	4	8	14
Petroleum ether extract	25%	9.33	15.33	21.33
	50%	13.33	21.33	30.66
	75%	17.66	29.33	45.66
	100%	29.33	53.66	86
	10%	7.33	15.33	20.66
Ethanol extract	25%	11	19.33	29
	50%	16.33	26.66	37
	75%	19.66	28	42.66
	100%	27.66	48.66	88.33
	10%	12	21.33	37.66
Methanol extract	25%	25.33	38.33	54.66
	50%	44.33	64	71.66
	75%	81.66	92.33	98
	100%	98	100	100

Table 2: Percentage of juvenile mortality in different solvent extracts of *Solanum torvum* fruit.




Fig 3: Inhibition of egg hatching after 24 hours treated in different solvents




Fig 4: Inhibition of egg hatching after 48 hours treated in different solvents




Fig 5: Inhibition of egg hatching after 72 hours treated in different solvents.




Fig 6: Juvenile mortality after 24 hours treated in different solvents




Fig 7: Juvenile mortality after 48 hours treated in different solvents




Fig 8: Juvenile mortality after 72 hours treated in different solvents.

CONCLUSION

The present study showed the fruit extracts of *Solanum torvum* in different solvents at different concentrations gave significant results of inhibition of egg hatching and juvenile mortality and may be useful for nematode management, which will be an environmentally and economically safe option and also recommended for the promotion of organic agriculture. However, further study on field efficacy trials and identification of the active chemical compounds of these extracts are needed.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Abad P, Favery B, Rosso MN, Castagnone-Sereno P (2003) Root-knot nematode parasitism and host response: molecular basis of a sophisticated interaction. *Molecular plant pathology* 4(4), 217–224. <https://doi.org/10.1046/j.1364-3703.2003.00170.x>
2. Azhagumurugan, C., & Rajan, M. K. 2015. Effect of *Meloidogyne incognita* on the growth characteristics of *Vigna mungo* treated with leaf extract of *Mimusops elengi*. *Pakistan Journal of Nematology*, **33**(1), 79-86.
3. Biney EE, Nkoom M, Darkwah WK, Puplampu JB. 2019. High-performance liquid chromatography analysis and antioxidant activities of extract of *Azadirachta indica* (neem) leaves. *Pharmacognosy Res.* 11. doi: <https://doi.org/10.4103/pr.pr.14.19>
4. Chitwood, D.J. 2003, Research on plant-parasitic nematode biology conducted by the United States Department of Agriculture–Agricultural Research Service. *Pest. Manag. Sci.*, **59**: 748-753.
5. Das S, Wadud A, Khokon MAR (2021) Evaluation of the effect of different concentrations of organic amendments and botanical extracts on the mortality and hatching of *Meloidogyne javanica*. *Saudi Journal of Biological Sciences* 28(7), 3759-3767. <https://doi.org/10.1016/j.sjbs.2021.03.041>
6. Dejene TA (2014) Opportunities for biological control of root-knot nematodes in organic farming systems: a review. *Int. J. Org. Agric. Res. Dev.* 9, 87–107
7. Fui LH. 2012. Knowledge and use of forest product as traditional medicine: the case of the forest-dwelling communities. *Int J Pharmacy Biol Sci.* 3(4):104–111.
8. Hunt DJ, Handoo ZA (2009) Taxonomy, identification and principal species. In: Perry RN, Moens M, Starr JL (eds) *Root-knot nematodes*. CABI, Wallingford, pp 55–97. <https://doi.org/10.1079/9781845934927.0055>
9. Irdani T, Sala T, Cutino I, Tarchi F. *Solanum torvum* mediates protection against the nematode *Meloidogyne incognita* in neighboring plants. *Journal of Plant Diseases and Protection*. 2023 Dec;130(6):1301-15.
10. Noling, J. W. 2002. The practical realities of alternatives to methyl bromide: concluding remarks. *Phytopathology*, **92**(12), 1373-1375. <http://dx.doi.org/10.1094/PHYTO.2002.92.12.1373>
11. Mahesha HS, Ravichandra NG, Rao MS, Narasegowda NC (2017) Bio-efficacy of different strains of *Bacillus* spp. against *Meloidogyne incognita* under in vitro. *Int. J. Current Microbiol. Appl. Sci.* 6 (11), 2511–2517. <https://doi.org/10.20546/ijcmas.2017.611.295>
12. Sasser, J.N. & Carter, C.C. 1985. Overview of the International *Meloidogyne* Project 1975-1984. In: Sasser, J.N. & Carter, C.C. (Eds.). *Advanced Treatise on Meloidogyne Vol. 1. Biology and control*. North Carolina State University, Graphics, Raleigh, USA. pp. 19-24.
13. Terefe, M. 2015. Evaluation of nematicidal action of some bio-agents and botanicals for eco-friendly management of root-knot nematodes, *Meloidogyne incognita* on tomato. *Science, Technology and Arts Research Journal*, 4(3), 71-78.