

Quality Attributes and Consumer Acceptability of *Chinchin* Produced from Blends of Rice, Cowpea, and Sweet Potato Flours

Abstract

This study investigated the proximate composition, mineral content, vitamin content, amino acid profile and sensory attributes of *chinchin* produced from blends of rice flour (RF), cowpea flour (CPF), and sweet potato flour (SPF). All analyses were carried out using standard methods. Proximate analysis revealed moisture content ranging from 5.11 % to 7.15 %, with the highest in the 60:35:5 blend. Protein content increased significantly (9.76 to 16.25 %) with increase in CPF, while fat content ranged from 12.35 to 16.21 %, ash content from 1.60 % to 5.11 %, and fiber content from 1.87 % to 5.02 %. Carbohydrate content varied from 50.26 % to 68.67 %, and energy content from 411.90 to 427.43 kcal/100g. Addition of CPF and SPF significantly altered the mineral content of the *chinchin* samples as follows; Potassium (95.82 to 139.29 mg/100 g), Calcium (89.00 to 158.57 mg/100 g), Magnesium (59.06 to 93.59 mg/100 g), Phosphorous (100.14 to 162.90 mg/100 g), Iron (65.14 to 82.13 mg/100 g), Zinc (10.40 to 34.76 mg/100 g). Analysis of vitamins (A, B1, B2, B3) showed significant differences among *chinchin* samples. Amino acid composition improved with increased CPF and SPF. Sensory evaluation rated 90:5:5 and 80:15:5 blends favorably compared to control.

Keywords

Chinchin, composite flours, nutritional composition, sensory evaluation, amino acids, consumer acceptability

Introduction

Chinchin just like many other snacks is traditionally prepared from cereal flours which have been known to be deficient in some vital nutrients like Sulphur-containing amino acids. There is therefore need to complement their nutrient content by incorporating other food sources. The majority of snacks are made using low-nutrient cereal flour most especially wheat flour one of the reasons being its unique content of gluten. It is necessary to supplement these snacks' nutritious value with various food sources. Most of the time, wheat has only been utilized to make *chinchin* and other pastries. However, there have been advancements recently in what the food processing industry refers to as composite flour technology—the addition of additional flours derived from fruits, vegetables, legumes, or tubers (Abioye *et al.*, 2020; Akubor, 2004).

Rice flour is a valuable ingredient with significant nutritional benefits and economic importance. Research has shown that rice flour is a gluten-free alternative with high nutritional value, containing a balanced complex of proteins, lipids, minerals, and vitamins (Marques *et al.*, 2023). It is particularly rich in high-grade protein with a good balance of amino acids, making it a suitable dietary option for various populations, including the elderly, who may be at risk of protein-energy malnutrition (Tumenova & Myktabayeva, 2023). Additionally, rice flour has functional and technological properties that make it a versatile ingredient for food processing, allowing for the development of products that cater to specific nutritional needs while maintaining taste and quality (Matsuda, 2019). Economically, rice flour plays a crucial role in reducing wheat flour imports in non-producing countries and is increasingly used in gluten-free bakery products, contributing to the growth of the healthy food market (Cristina *et al.*, 2021). Moreover, the protein composition of rice flour makes it a valuable component in enhancing the nutritional quality of bakery products, especially in the context of dietary restrictions or health conditions that require specific protein intake adjustments (Andrey, 2023). With its complete amino acid profile and low dietary fiber content, rice flour offers a promising solution for increasing the nutritional value of food products while addressing dietary limitations, such as the need for complete proteins with minimal fiber for patients with gastrointestinal issues. Overall, the nutritional richness and functional properties of rice flour make it a versatile and economically significant ingredient in the food industry, supporting the development of diverse and nutritious food products for various consumer needs and preferences.

Cowpeas are major sources of proteins, some vital amino acids, minerals, vitamins, and dietary fibre, being a significant legume in West African diets. They give diversity to meals in the tropics, especially

ones that consist mostly of starchy staples like plantains, grains, and tubers. They are readily available sources of protein due to their high yield and drought tolerance, which allow them to flourish even in areas unsuitable for other legumes. Cowpeas have a high protein content, which is utilised to improve the nutritional value of food. Cowpea has therefore been added to *chinchin* and other similar products to supplement foods and make them more nutritious (Alamu *et al.*, 2020).

Sweet potatoes are a nutrient-dense root vegetable renowned for their rich nutritional composition. Packed with complex carbohydrates, they serve as an excellent energy source, while their high dietary fibre content supports digestive health and helps regulate blood sugar levels. Sweet potatoes are abundant in essential vitamins, particularly vitamin A in the form of beta-carotene, which is crucial for vision, immune function, and skin health (Aburime *et al.*, 2021). Additionally, sweet potatoes provide a spectrum of B vitamins, including B6, necessary for metabolism and neurological function. These tubers also offer minerals like potassium, manganese, and copper, contributing to cardiovascular health, bone density, and antioxidant defence. With a low glycemic index and various antioxidants, sweet potatoes are a versatile and nutritious food that can be a valuable part of a well-balanced diet (Ndife *et al.*, 2020).

2.0 Materials and Methods

2.1. Sample preparation and flour blend formulation

Rice flour, sweet potato flour and cowpea flour were produced as in Figures 1, 2 and 3. These flours were then blended in different ratios as in Table 1. *Chinchin* was prepared from different blends of rice flour, sweet potato and cowpea flours. Refined wheat flour was used as Control (reference) standard

2.2. Chinchin Preparation

Chinchin was prepared using the method outlined by Bongjo *et al.* (2023). Hundred grams (100 g) of the composite flour was weighed and sieved (250-micron particle size). The recipe is as shown in Table 2. All other ingredients (sugar, salt, and baking powder) were also added. Margarine, eggs, and milk was turned in and mixed to form dough. The dough was briefly kneaded, rolled out (2 cm thick) and cut into small squares (2 cm by 2 cm). It was deep fried till it turns golden brown. It was then drained, cooled, packaged (in Low Density Polyethylene-LDPE) and stored at room temperature.

2.3. Analyses

2.3.1 Proximate analysis

The proximate analysis of the composite flours were determined by the official methods of AOAC (2012). Energy was calculated using Atwater factor (fat x 9 + carbohydrate x 4 + protein x 4 kcal/100 g).

Table 1: Flour based formulation for *chinchin*

| Samples | Rice flour | Ingredient (%) | |
|---------|-------------------|----------------|--------------------|
| | | Cowpea flour | Sweet potato flour |
| 100:0:0 | 100 | 0 | 0 |
| 90:5:5 | 90 | 5 | 5 |
| 80:15:5 | 80 | 15 | 5 |
| 70:25:5 | 70 | 25 | 5 |
| 60:35:5 | 60 | 35 | 5 |
| CTRL | 100 % Wheat flour | | |

Table 2: Recipe used for the preparation of *chinchin*

| Ingredient | Composition (g) |
|-----------------|-----------------|
| Composite flour | 100 |
| Margarine | 25 |
| Sugar | 25 |
| Egg | 1 egg |
| Milk | 10 |
| Baking Powder | 5.6 |

UNDER PEER REVIEW




Figure 1: Flow Chart for rice flour Production
Source: (Osuji et al., 2020)




Figure 2: Flow chart for sweet potato flour production
Source: (Ndife et al., 2020)




Figure 3: Flow Chart for the Production of cowpea flour
Source: (Ibidapo et al., 2017)

2.3.2 Determination of Minerals

The minerals Ca, K, Mg, Fe, P and Zn were determined. The optimum range for each element was prepared and all the operational instruction for setting up the instrument for the analysis of specific element was strictly followed. The ash residues were digested with 5 mL of concentrated nitric acid, filtered and the filtrate was transferred 100 mL volumetric flask and diluted with distilled water to 100 mL volume. This was done for all the samples, and stored at room temperature pending **Atomic absorption spectroscopy (AAS)** analysis (AOAC, 2012).

2.3.3. Determination of Vitamin Content

The method of AOAC (2012) using the colourimeter was adopted. Vitamins A, B1, B2 and B3 were determined

2.3.4. Amino acid analysis of the *chinchin*

Amino acids composition was quantified after hydrolysis using model 120A PTH amino acid analyzer (HPLC) in the reversed-phase column, after derivatization with 9-fluorenylmethyl chloroformate and a fluorescence detector (AOAC, 2015). The sample was dried at 70 °C to constant weight, defatted, hydrolyzed, evaporated in a rotary evaporator and loaded into the biosystem PTH Amino Acid Analyzer. The concentration of the individual amino acids was calculated in relation to the protein content.

2.4 Evaluation of Sensory quality attributes of *chinchin* samples

A panel of 30 individuals were employed for this study, where a 9-point hedonic scale was used; with 9 representing like extremely and 1 representing dislike extremely.

3.0 Results and Discussion

3.1 Proximate composition of *chinchin* from rice, cowpeas and sweet potato

The proximate composition of the *chinchin* samples is presented in Table 3. The moisture content significantly increased ($p < 0.05$) with the increasing incorporation of CPF and SPF. This increase in moisture content could be attributed to the increased hydrophilic properties of the fiber in CPF and SPF as their levels of incorporation rose. Moisture content is a critical indicator of a product's shelf stability; the lower the moisture content, the better the shelf stability (Ayogu *et al.*, 2016). Bongjo *et al.* (2023) reported similar results for wheat-defatted peanut-orange peel based *chinchin*. These results are similar those reported by (Ayogu *et al.*, 2016) (7.82–9.76%) who evaluated the potential of two local cowpea varieties for wheat-based cookies and higher than reported by Adegunwa *et al.*, (2014) (3.98-5.05%) who produced *chinchin* from a millet-wheat composite flour. The trend in results also agree with other studies (Akindele *et al.*, 2017).

The protein content varied significantly across all samples. This significant increase ($p < 0.05$) in protein content was observed as the incorporation of CPF and SPF increased. These results are higher than those of Adegunwa *et al.* (2014), who produced *chinchin* from millet and wheat. **Adebayo-Oyetero *et al.* (2017)** reported protein contents ranging from 11.1% to 14.9%, and similar ranges were reported by another study with 10.51% to 14.58%. This indicates that the *chinchin* produced in this study is rich in protein. Additionally, the protein content observed in this study was higher than those Badau *et al.* (2013) reported for Garabia (a Nigerian snack produced from different rice varieties), ranging from 10.72 % to 12.86 %.

The substitution with CPF and SPF in the *chinchin* samples increased fat content. This increase in fat content differed significantly among all the samples and could be attributed to the fat content in the CPF as well as due to the recipe. This finding aligns with a study by Dharsenda *et al.* (2015) on the substitution of defatted peanut flour in peanut flour-based cookies. However, the fat content results in this study are lower than those reported by Badau *et al.* (2013) for Garabia-a Nigerian traditional snack who reported 28 to 42.30 % for fat content. The high fat content in the *chinchin* samples was expected, given the significant increase in the OAC of the flours with higher CPF incorporation. Additionally, this high fat content could be attributed to oil absorption during frying and variations in the recipes.

The ash content of the *chinchin* samples increased significantly, an increase in the ash content of the *chinchin* samples was seen as substitution with CPF increased. All samples were significantly different ($p < 0.05$) from each other. The increase in ash content could be attributed to the high ash content of CPF of 3.43 % and SPF of 3.12% (Table 4). Bongjo *et al.* 2023) also reported an increase in the ash content of *chinchin* produced from defatted peanut flour and orange peel flour. The ash content in this study was higher than those reported by several studies on the production of *chinchin* and other snacks (Badau *et al.*, 2013; Deedam *et al.*, 2020; Ugwuanyi *et al.*, 2020). However, the trend in results are

comparable to those reported by (Ajibola *et al.*, 2015) in biscuits incorporated with Moringa flour and cocoa powder. The ash content of a product indicates a rough estimate of its mineral content. (Adelekan *et al.*, 2019). This study therefore indicates that the *chinchin* samples would contribute enormous mineral elements to the body.

There was a significant increase ($p < 0.05$) in fiber content as the substitution with CPF increased. This increase can be attributed to the high fiber content (2.12 %) in CPF, as noted in Table 4, and also reported by Ayogu *et al.* (2016) to contain fibre contents ranging from 4.91 to 5.49 %. The crude fiber content in this study was lower than that reported by Singh and Arivuchudar (2018) for peanut-based cookies. However, the fiber content in this study was comparatively higher than those reported by Adebayo-Oyetoro *et al.* (2017), Deedam *et al.* (2020b, 2020a), and Ndife *et al.* (2020), who produced *chinchin* from different composite flours. The results were also comparable to those by Adegunwa *et al.* (2014) in their wheat-millet *chinchin*.

A significant decrease ($p < 0.05$) was observed in the carbohydrate content with increase in CPF and SPF incorporation. This outcome is consistent with research by Shakpo and Osundahunsi, (2016) and Nwafor *et al.* (2020), who produced *Ipekere Agbado* (cowpea-enriched maize snack) and Yam-cowpea flour, respectively, and reported a decrease in carbohydrate content (65-30 %) and (81-63 %). This study's carbohydrate content is higher than that of (Badau *et al.*, 2013) but greater than that of Nwafor *et al.* (2020)'s study.

The *chinchin* samples' energy contents increased. As the percentage of CPF grew, there was some significant decrease ($p < 0.05$) across the samples as substitution with CPF increased. These findings are in tandem with other authors who produced *chinchin* from different composite flours (Bongjo *et al.*, 2023; Ubbor *et al.*, 2022; Ubbor and Akobundu, 2009).

3.2 Mineral content of *chinchin* from rice, cowpea and sweet potato flour blends

The mineral composition of the *chinchin* produced is presented in Table 4. A significant increase in Calcium contents ($p < 0.05$) was observed across all the samples with increase in the incorporation of CPF and SPF. Calcium is known to play a major role in muscle contraction, building strong bones and teeth, blood clotting, nerve impulse, transmission, regulating heart beat and fluid balance within cells (Grace *et al.*, 2015). It has also been identified to play major role in managing blood pressure, and preventing breast cancer. These results are lower than those reported by Bongjo *et al.* (2023) for defatted peanut-based *chinchin* who revealed calcium contents ranging from 153.23 to 415.69 mg/100g. Yilma and Admassu, (2019) also reported lower calcium contents of 0.22 to 0.38 mg/100g for biscuits from cowpea-wheat flour blends. Akubor *et al.* (2023) in a study to produce noodles from wheat, unripe banana and cowpea flour blends, the calcium content was observed to range between 84.40 to 126.0 mg/100g which is far lower than results obtained in the present study.

The iron content of the samples increased significantly ($p < 0.05$). The increase in iron content was observed as the level of incorporation of CPF and SPF in the flours increased. Iron plays a crucial role in strengthening the immune system. It is a functional component of hemoglobin and other essential compounds involved in respiration, immune function, and cognitive development. Ensuring an adequate iron intake is especially important in the diets of pregnant women, nursing mothers, and infants to prevent anemia (Olayinka and Etejere, 2018). Results in this study are higher than those reported by for cookies incorporated with different varieties of cowpeas ranging from 6.49 to 8.62 mg/100g. This is also consistent with reports for *Garabia* produced from varieties of rice (Badau *et al.*, 2013). Also, authors have reported iron contents ranging from 0.06 to 0.149 mg/100g for a cowpea enriched maize snack (*Ipekere Agbado*).

There was a significant increase ($p < 0.05$) in potassium content as the level of CPF and SPF incorporation increased. Potassium is known to play vital roles in maintaining fluid balance and ensuring the proper functioning of essential organs such as the brain, nerves, heart, and muscles (Asouzu *et al.*, 2020). Additionally, potassium aids in nerve impulse transmission and is a major cation in intracellular fluid (Olayinka and Etejere, 2018). Bongjo *et al.* (2023) witnessed a similar increasing trend with increase in defatted peanut flour and orange peel for the production of *chinchin*. Other reports have reported lower values (6.29-7.8 mg/100g) for cowpea-wheat flour snacks Yilma & Admassu, (2019) and 25.73 to 30.65 mg/100g for wheat-based cookies supplemented with cowpea flours.

Magnesium contents increased with significant difference ($p < 0.05$) across samples. Combining RF, CPF, and SPF flour increased the magnesium content of the snacks, as seen by the higher values obtained for the composite flour *chinchin* samples when compared to CTRL (wheat flour *chinchin*). The value found in this investigation exceeded the 5.60–13.60 mg/100g range previously reported for maize–wheat chin–chin (Dong *et al.*, 2016).

Table 3 Proximate Composition (%) of *chinchin* produced from flour blends of rice, cowpea and sweet potato

| RF: CPF: SPF | Moisture content | Crude protein | Crude fat | Fibre | Ash | CHO | Calories (kcal) |
|---------------------|-------------------------|--------------------------|--------------------------|-------------------------|-------------------------|--------------------------|---------------------------|
| 100:0:0 | 5.23 ^b ±0.01 | 10.12 ^b ±0.02 | 12.55 ^b ±0.03 | 1.87 ^a ±0.02 | 1.73 ^b ±0.01 | 68.51 ^e ±0.03 | 427.43 ^e ±0.22 |
| 90:5:5 | 5.59 ^c ±0.02 | 12.05 ^c ±0.05 | 13.63 ^c ±0.16 | 3.09 ^c ±0.04 | 2.62 ^c ±0.00 | 63.01 ^d ±0.20 | 422.97 ^c ±0.73 |
| 80:15:5 | 5.95 ^d ±0.07 | 14.11 ^d ±0.01 | 15.18 ^d ±0.03 | 3.55 ^d ±0.02 | 3.20 ^d ±0.01 | 58.00 ^c ±0.09 | 425.09 ^d ±0.20 |
| 70:25:5 | 6.24 ^e ±0.02 | 15.20 ^e ±0.03 | 15.82 ^e ±0.04 | 4.81 ^e ±0.02 | 4.18 ^e ±0.01 | 53.74 ^b ±0.05 | 418.20 ^b ±0.32 |
| 60:35:5 | 7.15 ^f ±0.05 | 16.25 ^f ±0.02 | 16.21 ^f ±0.03 | 5.02 ^f ±0.01 | 5.11 ^f ±0.02 | 50.26 ^a ±0.09 | 411.90 ^a ±0.13 |
| CTRL | 5.11 ^a ±0.02 | 9.76 ^a ±0.06 | 12.35 ^a ±0.01 | 2.52 ^b ±0.02 | 1.60 ^a ±0.01 | 68.67 ^e ±0.06 | 424.84 ^d ±0.09 |

Values are presented as means±standard deviation of triplicate determinations.

Means within the same column with different superscripts are significantly different at p<0.05

RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour, CHO-carbohydrate

Comparable ranges of 88.62–178.05 mg/100g were reported by Ndife *et al.* (2020) for *chinchin* prepared from maize, soybean and sweet potato flour blends. Because it maintains a healthy immune system, stabilizes heart rhythm, helps to maintain normal muscle and nerve function, and keeps bones strong, magnesium is necessary for optimal health. Akubor *et al.* (2023) reported lower magnesium contents (35.63 to 53.26 mg/100g) for cowpea-based noodles.

The results for zinc indicate a significant increase ($p < 0.05$) in zinc content with the increased incorporation of CPF and SPF into *chinchin*. This trend aligns with findings by Adeleke *et al.* (2021), who reported an increase in zinc content from 2.37 mg/100g to 3.50 mg/100g in *chinchin* enriched with polyphenol extracted from *Amaranthus viridis*. Additionally, Guyih *et al.* (2020) found zinc content values ranging from 5.75 mg/100g to 7.12 mg/100g for wheat-almond-carrot-based cookies, while Akindede *et al.* (2017) reported zinc values from 11.07 mg/100g to 17.93 mg/100g for *chinchin* enriched with *Ugu* and Indian spinach leaves. However, these results differ from those reported by Singh and Arivuchudar, (2018) for peanut-incorporated cookies. Zinc is an essential micronutrient involved in protein formation, blood formation, wound healing, taste perception, growth, and the maintenance of all tissues. It also plays a vital role in the immune system and is a component of many enzymes (Adenike, 2013).

Phosphorous content of the *chinchin* samples increased significantly ($p < 0.05$) with addition of CPF and SPF. Phosphorus deficiency leads to decreased growth, poor tooth development, and rickets (Ayogu *et al.*, 2016). Some reports have reported phosphorus contents ranging from 13.35 to 18.26 mg/100g in cookies. Another parallel study on noodles produced from cowpea-wheat flour reported phosphorus contents of between 35.75 to 148.75 mg/100g. Ndife *et al.* (2020) got similar results ranging from 78.23 to 139.17 mg/100g for *chinchin* from maize, soybean and sweet potato flour blends.

Table 4 Mineral content (mg/100 g) of *chinchin* from rice, cowpea and sweet potato flour blends

| RF: CPF: SPF | Potassium | Calcium | Magnesium | Phosphorus | Iron | Zinc |
|--------------|---------------------------|---------------------------|--------------------------|---------------------------|--------------------------|--------------------------|
| 100:0:0 | 101.65 ^b ±1.33 | 89.00 ^a ±0.77 | 63.06 ^b ±0.61 | 121.98 ^d ±0.09 | 67.90 ^b ±0.04 | 10.40 ^a ±0.08 |
| 90:5:5 | 95.82 ^a ±0.58 | 99.56 ^b ±0.11 | 59.06 ^a ±0.39 | 100.14 ^a ±0.02 | 65.14 ^a ±0.02 | 13.19 ^b ±0.07 |
| 80:15:5 | 112.28 ^c ±0.61 | 107.09 ^c ±0.33 | 65.67 ^c ±0.37 | 112.21 ^b ±0.24 | 71.31 ^d ±1.15 | 21.23 ^d ±1.11 |
| 70:25:5 | 120.80 ^d ±0.34 | 113.77 ^d ±0.79 | 78.91 ^e ±0.16 | 141.09 ^e ±1.03 | 78.38 ^c ±0.54 | 31.26 ^e ±0.94 |
| 60:35:5 | 131.51 ^e ±2.26 | 120.62 ^e ±0.84 | 93.59 ^f ±0.25 | 162.90 ^f ±0.16 | 82.13 ^f ±0.65 | 34.76 ^f ±0.43 |
| CTRL | 139.21 ^f ±1.33 | 158.57 ^f ±1.34 | 68.02 ^d ±0.16 | 113.27 ^c ±0.94 | 76.78 ^e ±0.42 | 14.74 ^c ±0.06 |

Values are presented as means±standard deviation of triplicate determinations.

Means within the same column with different superscripts are significantly different at $p < 0.05$

RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour

3.3 Vitamin content of *chinchin* from rice, cowpea and sweet potato flour blends

The vitamin content of the *chinchin* made from rice, cowpea and sweet potato flour blends as shown in Table 5, indicates that the inclusion of CPF and SPF flour in rice flour significantly ($p < 0.05$) increased across the samples. The values ranged from 3.14 to 8.12 µg/100g. (Ndife *et al.*, 2020) observed a similar trend in pro-vitamin A contents and recorded 0.3 to 2.1 µg/100g of *chinchin*. These are higher than those reported in this study. These findings are consistent with those of Makanjuola and Adebawale, (2020), who reported an increase in vitamin A content (0.45–5.67 mg/100 g) with the inclusion of cocoyam flour in wheat-cocoyam flour-based biscuits. Vitamin A is crucial for proper and clear vision, and consuming foods rich in this vitamin is essential for improving the body's vitamin A levels. However, all the values for vitamin A in this study were below the recommended daily intake of 300 µg/day (Tadesse *et al.*, 2015).

Thiamine (Vitamin B1) functions as the coenzyme thiamine pyrophosphate (TPP) in the metabolism of carbohydrates and branched-chain amino acids. The inclusion of CPF and SPF flour in the blends resulted in a significant ($p < 0.05$) increase in thiamine content. The increase in thiamine content is likely

due to the inclusion of CPF and SPF. This finding aligns with the report by Dinnah *et al.* (2020), who observed an increase in thiamine content (1.52–2.45 mg/100 g) with the addition of almond seed and carrot flour in blends. Ndife *et al.* (2020) reported higher values of 0.02 to 0.14 mg/100g for *chinchin* from maize-soybean-orange fleshed sweet potato composite flour.

No significant difference ($p>0.05$) observed in the B2 samples. The B vitamins are needed for carbohydrate and protein metabolism, and are essential for growth, well structuring and functioning of the cells (Dinnah *et al.*, 2020). Dinnah *et al.* (2020) reported higher vitamin contents in vitamin B2 (0.65 – 0.92 mg/100 g) for cookies incorporated with carrot flour and almond seed flour. Ndife *et al.* (2020) reported similar results ranging from 0.03 to 0.09 mg/100g.

Vitamin B3 contents significant increase as the level of CPF and SPF inclusion increased. These results are higher than those reported by Badau *et al.* (2013) for *Garabia* produced from different rice varieties and cowpea flour. They reported vitamin B3 contents ranging from 0.05 to 0.23 mg/100g. Ndife *et al.* (2020) also reported higher vitamin B2 contents ranging from 0.41 to 1.31 mg/100g.

UNDER PEER REVIEW

Table 5 Vitamin content of mg/100 g) *chinchin* from rice, cowpea and sweet potato flour blends

| RF: CPF: SPF | Provitamin A | Thiamine | Riboflavin | Niacin |
|--------------|-------------------------|-------------------------|-------------------------|-------------------------|
| 100:0:0 | 3.43 ^b ±0.01 | 0.05 ^d ±0.00 | 0.04 ^a ±0.00 | 0.49 ^b ±0.01 |
| 90:5:5 | 5.04 ^c ±0.04 | 0.03 ^b ±0.00 | 0.04 ^a ±0.00 | 0.60 ^c ±0.02 |
| 80:15:5 | 6.10 ^d ±0.02 | 0.05 ^c ±0.00 | 0.05 ^a ±0.00 | 0.64 ^d ±0.02 |
| 70:25:5 | 7.54 ^e ±0.02 | 0.06 ^e ±0.00 | 0.08 ^a ±0.01 | 0.80 ^e ±0.01 |
| 60:35:5 | 8.12 ^f ±0.02 | 0.07 ^f ±0.00 | 0.09 ^a ±0.00 | 0.90 ^f ±0.03 |
| CTRL | 3.14 ^a ±0.02 | 0.02 ^a ±0.00 | 0.04 ^a ±0.00 | 0.42 ^a ±0.02 |

Values are presented as means±standard deviation of triplicate determinations. Means within the same column with different superscripts are significantly different at $p<0.05$ RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour

3.6 Amino acid composition of *chinchin* from rice, cowpea and sweet potato flour blends

The essential amino acids of the *chinchin* produced are presented in Table 6. Essential amino acids are very important for the body, yet little or none of them is synthesized by the human body. As such, essential amino acids are supplied to the body by the diet (Michael and Shamim, 2023). As variations were made in the formulation by addition of CPF and SPF, the amounts of essential amino acids in all *chinchin* increased significantly ($p<0.05$). The high essential amino acid content of *chinchin* could be as a result of the high protein content of the CPF (Tables 3). The relationship between amino acids and proteins has been established by prominent researchers. Mosse and Huet, (2010) explained that amino acids are the building blocks of proteins, thus high values of proteins in a food indicate high amino acid content and vice versa. Comparing the essential amino acid content of the fortified *chinchin* in this study, the result shows that the *chinchin* made from rice flour/cowpea/sweet potato were richer in amino acids, indicating the availability of more essential amino acids in the *chinchin* from the blends than the CTRL sample. Several studies have also profiled the amino acid content of cowpea (Elharadallou *et al.*, 2015).

The non-essential amino acids of the *chinchin* produced are presented in Table 7. Non-essential amino acids are synthesized in the human body, and the amount obtained from food acts only as supplements. The high non-essential amino acid content of *chinchin* suggests that it is not only essential amino acids but also non-essential amino acids.

The incorporation of CPF and SPF into the formulation caused a significant increase ($p<0.05$) in the non-essential amino acid content of the samples. It is reported that the addition of a raw material with high amount of nutrients to a material that is lower in nutrients brings about a rise in concentration of nutrients in the latter (Chen *et al.*, 2018). The results in this study follow the same trend but were lower than those reported by Ajoke *et al.* (2018) for extruded snacks from pearl millet and germinated pigeon pea. This goes further to support the study which profiled the amino acids in cowpea flour (Elharadallou *et al.*, 2015).

Table 6: Essential amino acid composition (g/100g) of *chinchin* from rice, cowpea and sweet potato flour blends

| RF: CPF: SPF | Lys | Met | Thr | Ile | Leu | Phe | Val | Trp | His | TEAA |
|----------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------------------------|
| 100:0:0 | 2.84 ^a ±0.02 | 0.87 ^b ±0.01 | 1.24 ^a ±0.01 | 0.91 ^a ±0.01 | 3.26 ^a ±0.01 | 2.96 ^a ±0.01 | 1.97 ^a ±0.02 | 3.06 ^a ±0.02 | 2.19 ^a ±0.01 | 19.27 ^a ±0.06 |
| 90:5:5 | 3.56 ^c ±0.01 | 1.24 ^c ±0.00 | 2.31 ^c ±0.01 | 2.67 ^c ±0.02 | 7.03 ^c ±0.05 | 4.14 ^c ±0.01 | 4.02 ^c ±0.02 | 4.03 ^c ±0.01 | 3.63 ^c ±0.01 | 32.61 ^b ±0.04 |
| 80:15:5 | 4.04 ^d ±0.01 | 1.41 ^d ±0.01 | 2.66 ^d ±0.02 | 3.06 ^d ±0.02 | 8.01 ^d ±0.01 | 4.95 ^d ±0.01 | 4.72 ^d ±0.02 | 4.33 ^d ±0.02 | 3.84 ^d ±0.01 | 37.00 ^d ±0.07 |
| 70:25:5 | 4.14 ^e ±0.01 | 1.64 ^e ±0.01 | 2.81 ^e ±0.01 | 3.26 ^e ±0.02 | 8.21 ^e ±0.01 | 5.20 ^e ±0.01 | 5.02 ^e ±0.01 | 4.76 ^e ±0.01 | 3.98 ^e ±0.02 | 38.99 ^e ±0.01 |
| 60:35:5 | 4.85 ^f ±0.01 | 1.86 ^f ±0.01 | 2.88 ^f ±0.00 | 4.03 ^f ±0.01 | 8.78 ^f ±0.02 | 5.47 ^f ±0.02 | 5.14 ^f ±0.00 | 4.83 ^f ±0.01 | 4.21 ^f ±0.01 | 42.04 ^f ±0.01 |
| CTRL | 3.22 ^b ±0.01 | 0.77 ^a ±0.01 | 1.55 ^b ±0.02 | 1.05 ^b ±0.01 | 3.92 ^b ±0.02 | 3.84 ^b ±0.02 | 2.26 ^b ±0.01 | 3.26 ^b ±0.00 | 2.84 ^b ±0.02 | 22.68 ^b ±0.03 |

Values are presented as means±standard deviation of triplicate determinations.

Means within the same column with different superscripts are significantly different at $p < 0.05$

RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour, TEAA-Total Essential AminoAcid

Table 7: Non-essential amino acid composition (g/100g) of *chinchin* from rice, cowpea and sweet potato flour blends

| RF: CPF: SPF | Ser | Cys | Tyr | Ala | Asp | Glu | Gly | Pro | Arg | TNEAA |
|----------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------------------------|-------------------------|-------------------------|-------------------------|--------------------------|
| 100:0:0 | 2.35 ^a ±0.02 | 1.76 ^a ±0.00 | 3.91 ^a ±0.02 | 0.65 ^a ±0.02 | 3.25 ^a ±0.01 | 6.97 ^a ±0.02 | 1.56 ^a ±0.01 | 1.02 ^a ±0.05 | 1.14 ^a ±0.02 | 22.59 ^a ±0.09 |
| 90:5:5 | 3.14 ^c ±0.02 | 2.05 ^c ±0.05 | 5.59 ^c ±0.04 | 1.73 ^c ±0.01 | 5.53 ^c ±0.01 | 9.13 ^c ±0.01 | 3.43 ^c ±0.01 | 1.86 ^c ±0.02 | 2.47 ^b ±0.23 | 34.91 ^c ±0.28 |
| 80:15:5 | 3.37 ^d ±0.04 | 2.20 ^d ±0.04 | 5.84 ^d ±0.02 | 2.19 ^d ±0.04 | 7.36 ^d ±0.01 | 11.05 ^d ±0.08 | 3.65 ^d ±0.01 | 2.28 ^d ±0.01 | 2.33 ^b ±0.00 | 40.25 ^d ±0.06 |
| 70:25:5 | 3.47 ^e ±0.02 | 2.33 ^e ±0.01 | 5.91 ^e ±0.04 | 2.24 ^e ±0.03 | 7.96 ^e ±0.02 | 13.23 ^e ±0.01 | 3.97 ^e ±0.02 | 2.91 ^e ±0.04 | 3.09 ^c ±0.04 | 45.09 ^e ±0.10 |
| 60:35:5 | 3.76 ^f ±0.02 | 2.58 ^f ±0.00 | 6.12 ^f ±0.02 | 2.53 ^f ±0.01 | 8.14 ^f ±0.02 | 15.27 ^f ±0.29 | 4.27 ^f ±0.01 | 3.34 ^f ±0.02 | 3.22 ^c ±0.03 | 49.21 ^f ±0.33 |
| CTRL | 2.46 ^b ±0.01 | 1.98 ^b ±0.02 | 4.84 ^b ±0.02 | 0.76 ^b ±0.01 | 4.34 ^b ±0.02 | 8.11 ^b ±0.01 | 2.08 ^b ±0.02 | 1.30 ^b ±0.01 | 1.27 ^a ±0.01 | 27.12 ^b ±0.03 |

Values are presented as means±standard deviation of triplicate determinations.

Means within the same column with different superscripts are significantly different at $p < 0.05$

RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour, TNEAA-Total Non-Essential Amino Acid

3.7 Sensory attributes of *chinchin* from rice, cowpea and sweet potato flour blends

Sensory attributes such as appearance, aroma, taste, crunchiness and overall acceptability are useful in determining the quality of *chinchin*. The sensory quality attributes as presented in Table 8 showed that the inclusion of CPF and SPF led to a significant difference ($p < 0.05$) between CTRL and the test samples. All samples were generally acceptable but as expected, the CTRL sample was rated significantly higher than the other samples containing rice flour and sweet potato. This therefore affirms the potential of using these flours in food product development. Several studies affirming the use of CPF and SPF for the production of acceptable snacks and other related food products have been reported (Badau *et al.*, 2013; Ayogu, *et al.*, 2016; Ndife *et al.*, 2020; Ubbor *et al.*, 2022).

Table 8: Sensory attributes of *chinchin* from rice, cowpea and sweet potato flour blends

| RF: CPF: SPF | Appearance | Aroma | Taste | Crunchiness | Overall acceptability |
|--------------|--------------------------|--------------------------|-------------------------|-------------------------|--------------------------|
| 100:0:0 | 6.68 ^a ±1.25 | 6.00 ^a ±1.20 | 5.68 ^a ±1.29 | 5.84 ^a ±1.77 | 6.58 ^{ab} ±1.54 |
| 90:5:5 | 7.30 ^{ab} ±0.92 | 6.85 ^{ab} ±1.50 | 6.70 ^b ±1.26 | 5.70±1.03 | 7.20 ^b ±1.11 |
| 80:15:5 | 7.25 ^{ab} ±1.02 | 6.60 ^{ab} ±1.35 | 6.90 ^b ±1.07 | 7.25 ^b ±1.07 | 7.15 ^b ±1.46 |
| 70:25:5 | 6.70 ^a ±1.22 | 6.05 ^a ±1.19 | 5.80 ^a ±1.36 | 6.00 ^a ±1.86 | 6.70 ^{ab} ±1.59 |
| 60:35:5 | 6.60 ^a ±1.43 | 6.00 ^a ±1.45 | 5.50 ^a ±1.40 | 5.70 ^a ±1.53 | 5.95 ^a ±1.79 |
| CTRL | 7.80 ^c ±1.06 | 7.15 ^c ±1.04 | 7.95 ^c ±0.83 | 7.05 ^b ±1.10 | 8.10 ^c ±0.55 |

Values are presented as means±standard deviation of triplicate determinations.

Means within the same column with different superscripts are significantly different at $p < 0.05$

RF-Rice Flour, CPF-Cowpea Flour, SPF-Sweet potato flour, CTRL-Wheat flour

Conclusion

Chinchin produced from rice, cowpea, and sweet potato composite flours offers a nutritionally superior and consumer-acceptable alternative to wheat-based products. These findings support the potential of these locally available crops for use in the commercial production of healthier snacks.

Disclaimer (Artificial intelligence)

Option 2:

Author(s) hereby declare that generative AI technologies such as Claude.ai and Consensus.app have been used during the writing or editing of manuscripts.

Details of the AI usage are given below:

1. Claude 3.5 Sonnet

2. Consensus.app

References

- Abioye, V. F., Olodude, O. A., Atiba, V., & Oyewo, I. O. (2020). Quality evaluation of $chinchin$ produced from composite flours of wheat and germinated finger millet flour. *Agrosearch*, 20(1), 13–22. <https://doi.org/10.4314/agrosh.v20i1.2s>
- Aburime, L. C., Ene-Obong, N., & David-Oku, E. (2021). Functional Properties and Sensory Evaluations of “Chinchin”, Bread and Biscuits Produced From Composite Flours From African Yam Bean, Orange Fleshed Sweet Potatoes, Plantain, Cocoyam, Maize, and Wheat. *European Journal of Food Science and Technology*, 9(1), 18–36.
- Adegunwa, M., Ganiyu, A., Bakare, H., & Adebowale, A. (2014). Quality evaluation of composite millet-wheat *Chinchin*. *Agriculture and Biology Journal of America*, 5(1), 33–39. <https://doi.org/10.5251/abjna.2014.5.1.33.39>
- Adelekan, E. O., Adegunwa, M. O., Adebowale, A. A., Bakare, H. A., & Alamu, E. O. (2019). Quality evaluation of snack produced from black pepper (*Piper nigrum* L.), plantain (*Musa paradisiaca* L.), and tigernut (*Cyperus esculentus* L.) flour blends. *Cogent Food and Agriculture*, 5(1), 1–14. <https://doi.org/10.1080/23311932.2019.1614285>
- Adeleke, S. A., Taiwo, K. A., & Gbadamosi, S. O. (2021). Studies on the Chemical Composition and Storage Properties of *Chinchin* Enriched with Polyphenol Extracted from *Amaranthus viridis* L. *Food Science and Nutrition Technology*, 6(1), 1–17. <https://doi.org/DOI: 10.23880/fsnt-16000244>
- Adenike, O. M. (2013). Studies on the Nutritional quality, anti-nutritional factors, amino acid profile and functional properties of co-fermented wheat/cowpea. *Elixir Food Science*, 54(1), 12360–12364.
- Ahure, D., Guyih, M. D., & Eke, M. O. (2020). Vitamin Content and Storage Studies of Cookies Produced from Wheat, Almond and Carrot Flour Blends. *Asian Food Science Journal*, 15(3), 31–41. <https://doi.org/10.9734/afsj/2020/v15i330154>
- Ajibola, C., Oyerinde, V., & Adeniyani, O. (2015). Physicochemical and Antioxidant Properties of Whole-Wheat Biscuits Incorporated with *Moringa oleifera* Leaves and Cocoa Powder. *Journal of Scientific Research and Reports*, 7(3), 195–206. <https://doi.org/10.9734/jsrr/2015/18070>
- Ajoke, M., Mopelola, O., Esther, O., Olusola, M., & Babatunde, I. (2018). *Physicochemical Properties and Amino Acid Profile of Extruded*. 19(2), 183–190.
- Akindele, O., Gbadamosi, O., Taiwo, K., Oyedele, D., & Adebooye, C. (2017). Proximate, Mineral, Sensory Evaluations and Shelf Stability of *Chinchin* Enriched with Ugu and Indian Spinach Vegetables. *International Journal of Biochemistry Research & Review*, 18(4), 1–14. <https://doi.org/10.9734/ijbcrr/2017/35736>
- Akubor, P. I. (2004). Protein contents, physical and sensory properties of Nigerian snack foods (cake, *chinchin* and puff-puff) prepared from cowpea - Wheat flour blends. *International Journal of Food Science and Technology*, 39(4), 419–424. <https://doi.org/10.1111/j.1365-2621.2004.00771.x>
- Akubor, P. I., Agada, T. O., & Okereke, G. O. (2023). Quality Evaluation of Noodles Produced From Blends of Wheat, Unripe Banana and Cowpea Flours. *Journal of Nutrition Food Science and Technology*, 4(2), 1–13. <https://doi.org/10.47485/2834-7854.1023>
- Alamu, E. O., Maziya-Dixon, B., Olaniyan, B., Pheneas, N., & Chikoye, D. (2020). Evaluation of nutritional properties of cassava-legumes snacks for domestic consumption-consumer acceptance and willingness to pay in Zambia. *AIMS Agriculture and Food*, 5(3), 500–520. <https://doi.org/10.3934/agrfood.2020.3.500>
- Andrey, S. (2023). Protein substances of rice flour and its use in wheat bread technology. *Naukovi Praci Nacional'nogo Universitetu Harčovih Tehnologij*. <https://doi.org/10.24263/2225-2924-2023-29-1-13>

AOAC. (2012). *Official Methods of Analysis. 18th Edition, Association of Official Analytical Chemists, Arlington, 806-84.*

Asouzu, A. I., Oly-Alawuba, N. M., & Umerah, N. N. (2020). Functional Properties and Chemical Composition of Composite Flour Made from Cooking Banana (*Musa Paradisiaca*) and Yellow Maize (*Zea Mays*). *Research Journal of Food and Nutrition*, 4(2)(2), 6–12. https://www.researchgate.net/profile/Nkemjika-Umerah/publication/341769201_Functional_Properties_and_Chemical_Composition_of_Composite_Flour_Made_from_Cooking_Banana_Musa_Paradisiaca_and_Yellow_Maize_Zea_Mays/links/5ed30d98299bf1c67d2cafa9/Functional-Prop

Ayogu, R. N. B., Nnam, N. M., & Mbah, M. (2016). Evaluation of two local cowpea species for nutrient, antinutrient, and phytochemical compositions and organoleptic attributes of their wheat-based cookies. *Food and Nutrition Research*, 60, 1–8. <https://doi.org/10.3402/fnr.v60.29600>

Badau, M. H., Ngozi, C., & Danbaba, N. (2013). Quality of Garabia (a nigerian traditional snack) from four varieties of rice as affected by the addition of Cowpea. *Advance Journal of Food Science and Technology*, 5(3), 249–254. <https://doi.org/10.19026/ajfst.5.3252>

Bongjo, N. B., Ahemen, S. A., Gbertyo, J. A., Guyih, M. D., & Muyong, M. G. (2023). Chemical and organoleptic properties of *Chinchin* produced from flour blends of wheat, defatted peanut and orange peels. *Journal of Nutritional Health & Food Engineering*, 13(1). <https://doi.org/10.15406/jnhfe.2023.07.00367>

Bongjo, N. B., Aondoaver, A. S., Gbertyo, J. A., Fulai, A. M. A., & Nchung, L. (2022). Nutritional and Functional Properties of Wheat-Defatted Peanut-Orange Peel Composite Flour. *European Journal of Nutrition & Food Safety*, 14(12), 74–86. <https://doi.org/10.9734/ejnfs/2022/v14i121283>

Bukuni, S. J., Kwagh-al Ikya, J., Dinnah, A., & Bongjo, N. B. (2022). Chemical and Functional Properties of Composite Flours Made from Fermented Yellow Maize, Bambara Groundnut, and Mango Fruit for 'Ogi' Production. *Asian Food Science Journal*, 21(2), 22–33. <https://doi.org/10.9734/afsj/2022/v21i230405>

Chen, Y., Michalak, M., & Agellon, L. B. (2018). Importance of Nutrients and Nutrient Metabolism on Human Health. *The Yale Journal of Biology and Medicine*, 9(2), 95–103.

Comparative Evaluation of Enriched Biscuits Made From Selected Cereals and Groundnut (*Arachis hypogaea*) Composite Flours. (2020). *Nigerian Journal of Agriculture, Food and Environment*, 16(2), 140–151.

Cristina, M., Rosell, Yaiza, B.-G., & Raquel, G. (2021). Rice flour breads. In *Trends in Wheat and Bread Making* (pp. 405–429). <https://doi.org/10.1016/B978-0-12-821048-2.00014-3>

Deedam, N. J., China, M. A., & Wachukwu, H. I. (2020). Utilization of Soursop (*Annona muricata*) Flour for the Production of *Chinchin*. *Agriculture and Food Sciences Research*, 7(1), 97–104. <https://doi.org/10.20448/journal.512.2020.71.97.104>

Dong, O., T. C., E., & S. U., O. (2016). Evaluation of the Quality of Composite Maize-Wheat *Chinchin* Enriched with *Rhynchophorus phoenicis*. *Journal of Food Research*, 5(4), 26. <https://doi.org/10.5539/jfr.v5n4p26>

Elharadallou, S. B., Khalid, I. I., Gobouri, A. A., & Abdel-Hafez, S. H. (2015). Amino Acid Composition of Cowpea (*Vigna unguiculata* L. Walp) Flour and Its Protein Isolates. *Food and Nutrition Sciences*, 06(09), 790–797. <https://doi.org/10.4236/fns.2015.69082>

Grace, U., Elijah, A., & Nicholas, A. (2015). Vitamin and mineral evaluation of mixed fruit jam from blends of pineapple, orange and sourplum. *Pakistan Journal of Food*, 25(3), 137–143. http://www.psfst.com/_jpd_fstr/4b5c479a7d874a815a94a4fb0cf987ad.pdf

- Guyih, M. D., Dinnah, A., & Ojotu Eke, M. (2020). Production and Quality Evaluation of Cookies from Wheat, Almond Seed and Carrot Flour Blends. *International Journal of Food Science and Biotechnology*, 5(4), 55. <https://doi.org/10.11648/j.ijfsb.20200504.11>
- Igbabul, B. D., Iorliam, B. M., & Umana, E. N. (2015). Physicochemical and Sensory Properties of Cookies Produced From Composite Flours of Wheat, Cocoyam and African Yam Beans. *Journal of Food Research*, 4(2), 150. <https://doi.org/10.5539/jfr.v4n2p150>
- Jan, R., Saxena, D. C., & Singh, S. (2018). Characteristics of wheat cookies supplemented with Chenopodium (*Chenopodium album*) flour: Physico-chemical, in-vitro antioxidant, textural and sensory properties. *International Journal of Advanced Research in Science and Engineering*, 7(4), 1802–1815. http://www.ijarse.com/images/fullpdf/1524848524_JK1652IJARSE.pdf
- Makanjuola, O. M., & Adebawale, O. J. (2020). Vitamins, Functional and Sensory Attributes of Biscuit Produced from Wheat-Cocoyam Composite Flour. *Journal of Scientific and Innovative Research*, 9(2), 77–82.
- Marques, A. C., Lidon, F. C., Coelho, A. R. F., Pessoa, C. C., Daccak, D., Luís, I. C., Simões, M., Scotti-Campos, P., Almeida, A. S., Guerra, M., Leitão, R. G., Bagulho, A., Moreira, J., Pessoa, M. F., Legoinha, P., Ramalho, J. C., Semedo, J. N., Palha, L., Silva, C., ... Reboredo, F. H. (2023). Elemental Composition and Implications on Brown Rice Flour Biofortified with Selenium. *Plants*, 12(8), 1–12. <https://doi.org/10.3390/plants12081611>
- Matsuda, T. (2019). Rice flour: A promising food material for Nutrition and Global Health. *Journal of Nutritional Science and Vitaminology*, 65, S13–S17. <https://doi.org/10.3177/jnsv.65.S13>
- Michael, J. L., & Shamim, S. M. (2023). Biochemistry, Essential Amino Acids. *StatPearls Publishing*.
- Mosse, J., & Huet, J. C. (2010). Amino acid composition and nutritional score for ten cereals and six legumes or oilseeds: causes and ranges of variations according to species and to seed nitrogen content. *Sciences Des Alimentations*, 10, 151–173.
- Ndife, J., Abasiokong, K. S., Nweke, B., Linus-Chibuezeh, A., & Ezeocha, V. C. (2020). Production and Comparative Quality Evaluation of *Chinchin* Snacks From Maize, Soybean and Orange Fleshed Sweet Potato Flour Blends. *Fudma Journal of Sciences*, 4(2), 300–307. <https://doi.org/10.33003/fjs-202-0401-220>
- Nwafor, J. O., Kanu, A. N., Kelechukwu, E. C., Nwohu, N. O., & Ezebuio, V. N. (2020). Physico-Chemical Properties of Water Yam and Cowpea Flour Blends for Production of Snacks. *South Asian Journal of Research in Microbiology*, January, 1–8. <https://doi.org/10.9734/sajrm/2020/v6i330149>
- Nwatum, I. A., Ukeyima, M. T., & Eke, M. O. (2020). Production and Quality Evaluation of Cookies from Wheat, Defatted Peanut and Avocado Composite Flour. *Asian Food Science Journal*, 15(4), 1–12. <https://doi.org/10.9734/afsj/2020/v15i430156>
- Odunlade, T. V., Famuwagun, A. A., Taiwo, K. A., Gbadamosi, S. O., Oyedele, D. J., & Adebooye, O. C. (2017). Chemical Composition and Quality Characteristics of Wheat Bread Supplemented with Leafy Vegetable Powders. *Journal of Food Quality*, 2017, 1–7. <https://doi.org/10.1155/2017/9536716>
- OECD. (2016). Cowpea (*Vigna unguiculata*). In *OECD Consensus Documents* (Vol. 6, Issue December 2015, pp. 211–241). OECD Publishing. <https://doi.org/10.1787/9789264253421-8-en>
- Olayinka, B. U., & Etejere, E. O. (2018). Proximate and chemical compositions of watermelon (*Citrullus lanatus* (Thunb.) Matsum and Nakai cv red and cucumber (*Cucumis sativus* L. cv Pipino). *International Food Research Journal*, 25(3), 1060–1066.
- Oluwamukomi, M. O., Awolu, O. O., & Olapade, K. T. (2021). Nutritional Composition, Antioxidant and Sensory Properties of a Maize-based Snack (Kokoro) Enriched with Defatted Sesame and Moringa

Seed Flour. *Asian Food Science Journal*, 20(10), 100–113.
<https://doi.org/10.9734/afsj/2021/v20i1030366>

- Omoba, O. S., Dada, O. O., & Salawu, S. O. (2015). Antioxidant properties and consumer acceptability of pearl millet – tiger nut biscuits. *Nutrition and Food Science*, 45(6), 818–828.
<https://doi.org/10.1108/NFS-06-2015-0074>
- Ram, S., Narwal, S., Gupta, O. P., Pandey, V., & Singh, G. P. (2020). Anti-nutritional factors and bioavailability: approaches, challenges, and opportunities. In *Wheat and Barley Grain Biofortification*. Elsevier Inc. <https://doi.org/10.1016/b978-0-12-818444-8.00004-3>
- Shakpo, I. O., & Osundahunsi, O. F. (2016). Effect of some processing methods on the proximate, mineral, microbiological and sensory qualities of cowpea enriched maize snack (Ipekere Agbado). *Research Journal of Food Science and Nutrition*, 1(1), 1–9. <https://doi.org/10.31248/rjfsn2016.006>
- Singh, R., & Arivuchudar, R. (2018). Formulation and evaluation of peanut flour incorporated cookies. *International Journal of Food Science and Nutrition*, 3(5), 56–59.
- Sombié, P. A. E. D., Compaoré, M., Coulibaly, A. Y., Ouédraogo, J. T., De La Salle Tignégré, J. B., & Kiendrébéogo, M. (2018). Antioxidant and Phytochemical Studies of 31 Cowpeas (*Vigna unguiculata* (L. Walp.)) genotypes from Burkina Faso. *Foods*, 7(9). <https://doi.org/10.3390/foods7090143>
- Tadesse, T. F., Nigusse, G., & Kurabachew, H. (2015). Nutritional, microbial and sensory properties of flat-bread (kitta) prepared from blends of maize (*Zea mays* L.) and orange-fleshed sweet potato (*Ipomoea batatas* L.) flours. *International Journal of Food Science and Nutrition Engineering*, 5(1), 33–3.
- Tumenova, G. T., & Myktabayeva, M. S. (2023). The Possibility of Using Rice Flour for the Production of Functional Meat Products. *Vestnik of M. Kozybayev North Kazakhstan University*, 4(4 (52.1)), 115–121. <https://doi.org/10.54596/2309-6977-2021-4-115-121>
- Ubbor, S. C., & Akobundu, E. N. T. (2009). Quality Characteristics of Cookies from Composite Flours of Watermelon Seed, Cassava and Wheat. *Pakistan Journal of Nutrition*, 8(7), 1097–1102.
<https://doi.org/10.3923/pjn.2009.1097.1102>
- Ubbor, S. C., Arukwe, D. C., Ezeocha, V. C., Nwoso, O. N., Iguh, B. N., & Nwibo, O. G. (2022). Production and Quality Evaluation of Ready To Eat Extruded Snacks From Flour Blends of Acha-Cowpea and Sweet Potato Starch. *Fudma Journal of Sciences*, 6(4), 245–153. <https://doi.org/10.33003/fjs-2022-0604-1071>
- Ugwuanyi, R. G., Eze, J. I., & Okoye, E. C. (2020). Effect of Soybean, Sorghum and African Breadfruit Flours on the Proximate Composition and Sensory Properties of *Chinchin*. *European Journal of Nutrition & Food Safety*, 12(1), 85–98. <https://doi.org/10.9734/ejnfs/2020/v12i130187>
- Vagadia, B. H., Vanga, S. K., & Raghavan, V. (2017). Inactivation methods of soybean trypsin inhibitor – A review. *Trends in Food Science and Technology*, 64, 115–125.
<https://doi.org/10.1016/j.tifs.2017.02.003>
- Yilma, M., & Admassu, S. (2019). Product Development and Quality Evaluation of Biscuit and Ready-to-Eat Snack from Cowpea-wheat Flour Blends. *Advances in Food Technology and Nutrition Sciences – Open Journal*, 5(3), 92–106. <https://doi.org/10.17140/aftnsoj-5-161>