

## Unveiling the Rice hybrid parental lines - A study of genetic variability for yield attributing traits (*Oryza sativa*. L)

### Abstract:

Sixty-five rice hybrid lines along with five checks were evaluated in augmented block design (ABD) at ICAR-IIRR Hyderabad, to determine the genetic variability, heritability and genetic advance for ten traits (which season, year?). Analysis of variance showed that significant variation was present in the hybrid parental lines studied. The phenotypic coefficient of variation was found to be higher than the genotypic coefficient of variation for all the traits. High GCV (abbreviate) and PCV levels were recorded for number of productive tillers and grain yield per plant. Heritability was high for all the traits studied. High heritability coupled with high genetic advance as percent of mean was recorded for majority of the traits suggesting importance should be given to these characters during selection. (give the values for variability parameters; eg: how much heritability percentage)

**Keywords:** Rice, genetic variability, heritability, genetic advance as percent of mean

### Introduction:

Rice is the most significant grain and staple food for over 100 countries, earning it the title "Global Grain". Despite being the second largest rice producer worldwide, India faces challenges with low productivity. Rice contains 80-90 grams of carbohydrates and 7-8 grams of protein per 100 grams of uncooked grains (Juliano [11], Kennedy, Burlingame, & Nguyen [10]). Rice has 0.3-0.5 grams of fiber (Slavin [16]). The increasing demand for rice, coupled with limited cultivable land and low domestic output, may result in future shortages. By 2025, an estimated 785 million tons of paddy will be required, a 70% increase from current levels, to meet this demand. The FAO (2020) reports that productivity of rice in India is 3.21 t/ha, significantly lower than China (6.35 t/ha) and the global average (4.15 t/ha). This low productivity is mainly due to the lack of high-yielding rice varieties, making it crucial for India to focus on enhancing rice productivity to sustain its population.

The existence of genetic variation is paramount for the efficacy of any agricultural enhancement initiative. A comprehensive understanding of genetic parameters such as variability, heritability, and genetic advance is imperative for the augmentation of crop breeding endeavorsendeavours. Genetic diversity among traits is indispensable for the

identification of ~~favorable~~favourable genotypes and the establishment of a robust genetic framework for selective breeding practices. Investigations into heritability, genetic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), and genetic advance yield critical insights into the traits being examined. This accumulated knowledge facilitates an understanding of gene action and the identification of the most appropriate breeding methodologies to adopt. In light of these considerations, the current study was undertaken to assess genetic variability within rice hybrid parental lines.

(Describe the importance of the use of hybrid lines in this study)

### **Materials and methods:**

The present study was conducted at ICAR- Indian Institute of Rice Research, Rajendranagar, Hyderabad during *Kharif*, 2023. Experiment was comprised of 65 hybrid parental lines and five checks (ISM, Tetep, CO 101A 51, BPT 5204 and TN-1) evaluated in Augmented block design. The experimental field was divided into five blocks. Recommended agronomic practices were followed during crop growth period. The parental lines were evaluated for the following traits: Days to 50 % flowering, plant height, panicle length, number of productive tillers, filled grains, 1000 seed weight, kernel length, kernel breadth, kernel length/ breadth and grain yield per plant.

The analysis of variance for the design of the experiment was carried out according to the procedure outlined by Panse and Sukhatme [13], Ferderer [5]. The phenotypic and genotypic coefficients of variation were estimated according to the method suggested by Burton and de Vane [3]. "Broad sense heritability ( $h^2$ ) expressed as the percentage of the ratio of the genotypic variance ( $\sigma^2_g$ ) to the phenotypic variance ( $\sigma^2_p$ ) and was estimated on genotype mean basis" as described by Allard [2] and "Genetic advance in absolute unit (GA) and percent of the mean (GAM), assuming selection of superior 5% of the genotypes was estimated in accordance with the methods" illustrated by Johnson *et al.* [9]

### **Results:**

#### **Analysis of variance:**

In the current study, the examination of variance concerning yield and its associated attributes indicated that significant discrepancies existed among the parental lines. The outcomes of the variance analysis for the augmented block design pertaining to all ten attributes are displayed in Table 1. The variability observed in the blocks was statistically significant for the traits assessed *viz.*, days to 50% flowering, plant height, filled grains, 1000 seed weight, grain yield per plant. The difference among the checks was found to be highly significant for all the traits under study.

#### **Mean performance:**

A wider range of mean variability was observed for the traits *viz.*, filled grains (78-242), plant height (69.05-145.0), days to 50% flowering (82-114), number of productive tillers (4-17) and 1000 seed weight (10.13-24.15). The results showed that these traits should be used as selection prospects to improve performance through a breeding procedure like selection and hybridization. [\(write units\)](#)

#### **Genetic variability:**

The degree of variability for any specific trait holds significant relevance for the enhancement of a population through selective breeding. The estimations of genotypic variation, phenotypic variation, heritability, genetic gain, genotypic coefficient of variation (GCV), and phenotypic coefficient of variation (PCV) across various traits are delineated in Table 2. The extent of environmental influence on any trait signifies the magnitude of the disparity between the genotypic and phenotypic coefficients of variation. A substantial difference denotes a greater environmental impact, while a minimal difference indicates a predominance of genetic influence. Phenotypic coefficient of variation (PCV) was higher than genotypic coefficient of variation (GCV) in all the studied characters which indicates the presence of environmental influence on the phenotypic expression of the character to some degree. [\(add references as required in this para.\)](#)

The difference between PCV and GCV was less for the characters, days to 50% flowering, panicle length, plant height, thousand seed weight. Phenotypic coefficient of variation ranged from 5.87 (days to 50% flowering) to 21.82 (Grain yield / plant), while GCV ranges from 6.28 (days to 50% flowering) to 23.88 (Grain yield /plant). (Fig.1) [\(unit\)](#)

**Table 1. Analysis of variance for yield components in sixty five hybrid parental lines of rice**

S.No	Traits	Block (df = 4)	Treatments (df = 64)	Treatment: Check (df = 4)	Treatment: Test vs. Check (df = 1)	Residuals (df = 16)
1	<b>DFP</b>	159.57 **	81.84 **	940.66 **	48.74 **	4.83
2	<b>PH</b>	312.51 **	347.91 **	4097.44 **	8.14 ns	18.14
3	<b>PL</b>	3.48 ns	12.33 **	39.45 **	123.65 **	2.09
4	<b>NPTS</b>	10.91 ns	6.81 ns	51.14 **	5.97 ns	5.14
5	<b>FG</b>	2018.28 **	1432.94 **	8236.26 **	25756.52 **	139.06
6	<b>TSW</b>	14.16 **	7.18 **	22.31 **	2.07 ns	1.23
7	<b>KL</b>	0.03 ns	0.20 **	2.68 **	2.48 **	0.05
8	<b>KB</b>	0.02 ns	0.06 ns	0.41 **	1.75 **	0.03
9	<b>L*B</b>	0.05 ns	0.05 **	0.48 **	0.83 **	0.02
10	<b>GYP</b>	22.15 **	31.53 **	99.34 **	246.40 **	4.14

<sup>ns</sup>  $P > 0.05$ ; \*  $P \leq 0.05$ ; \*\*  $P \leq 0.01$

Formatted: Font: Italic

DFF- days to 50% flowering, PH- Plant height, PL- panicle length, NPTS- number of productive tillers, FG- filled grains, TSW- 1000 seed weight, KL- kernel length, KB- kernel breadth, L\*B-kernel length/breadth, GYP-grain yield per plant

UNDER PEER REVIEW

**Table 2. Genetic parameters for yield components in sixty-five hybrid parental lines of rice**

Trait	Mean	GCV (%)	PCV (%)	$h^2$ (%)	GAM (%) at 5%
<b>DFP</b>	98.79	5.87	6.28	87.43	11.32
<b>PH</b>	96.75	11.25	12.08	86.72	21.61
<b>PL</b>	23.41	11.22	12.8	76.72	20.27
<b>NPTS</b>	9.51	20.12	22.28	74.67	44.32
<b>FG</b>	163.75	14.85	16.5	80.96	27.57
<b>TSW</b>	16.01	15.19	16.69	82.74	28.49
<b>KL</b>	5.46	8.22	10.32	79.20	19.27
<b>KB</b>	2.29	9.16	11.56	86.32	22.51
<b>L*B</b>	2.4	14.28	16.62	89.68	28.30
<b>GYP</b>	20.93	21.82	23.88	83.43	41.11

#### Abbreviations

High GCV and PCV values were recorded for grain yield per plant (21.82 and 23.88) and number of productive tillers (20.12 and 22.28). Similar results were reported by Hemalatha et al. [8], Gayathridevi et al. [6], Pradhan et al. [14], Gupta et al. [7], Yashwanth et al. [19] for grain yield per plant.

Moderate GCV and PCV values were recorded for plant height (11.25 and 12.08), panicle length (11.22 and 12.80), filled grains (14.85 and 16.50), 1000 seed weight (15.19 and 16.69), kernel length / breadth (14.28 and 16.62). This suggests moderate variability for these traits, which can be exploited for improvement through selection in advanced generations.

Low PCV and GCV values recorded for days to 50% flowering (5.87 and 6.28). Vanaja and Babu et al. [18], Uma Devi et al. [17], Mythili et al. [12], Rashid et al. [15] observed low values for days to 50% flowering. Low GCV values were noticed for kernel length (8.22) and kernel breadth (9.16) while PCV values are moderate (10.32 and 11.56). Lower GCV and PCV suggest a narrow genetic base for these traits. Enhancing these traits

may require hybridization or induced mutagenesis to broaden the genetic base, followed by pedigree selection in later generation

UNDER PEER REVIEW


### Heritability:

Broad-sense heritability is valuable for selecting elite genotypes from homozygous lines, while narrow-sense heritability is more useful for segregating populations. Heritability estimates for different traits ranged from 74.67% (number of productive tillers) to 89.68% (kernel length / breadth). ([what is the h2 used in this study?](#))


Heritability estimates were high for traits such as days to 50% flowering (87.43), plant height (86.72), panicle length (76.72), number of productive tillers (74.67), filled grains (80.96), 1000 seed weight (82.74), kernel length (79.20), kernel breadth (86.32), kernel length/breadth (89.68), grain yield per plant (83.43). (Fig. 2) ([unit](#))

### Genetic advance:

Genetic advance as percent of mean was high for plant height (21.61), panicle length (20.27), number of productive tillers (44.32), filled grains (27.57), 1000 seed weight (28.49 gm), kernel breadth (22.51mm), kernel length / breadth (28.30 mm), grain yield per plant (41.11 gm) and moderate for days to 50 % flowering (11.32) and kernel length (19.27 mm).



**Fig 1. Genotypic and phenotypic coefficients of variation for yield components in rice hybrid parental lines**



**Fig 2. Heritability and genetic advance as per cent of mean for yield components in rice hybrid parental lines**

High genetic advance coupled with high heritability was recorded for plant height, panicle length, number of productive tillers, filled grains, 1000 seed weight, kernel breadth, kernel length/ breadth and grain yield per plant. Similar results for plant height were reported by Akinwale et al. [1], Gayathridevi et al. [6], Pradhan et al. [14] for grain yield per plant. Demeke et al. [4] reported similar results for plant height, number of productive tillers, filled grains, 1000 seed weight and grain yield per plant. Generally, traits with high heritability and high genetic advance are controlled by additive gene action and can be improved through simple or progeny selection methods.

High heritability coupled with moderate genetic advance was recorded for days to 50% flowering and kernel length. High heritability with moderate or low genetic advance can be improved by intermating superior genotypes of segregating populations developed from combination breeding.

### **Conclusion:**

Greater genetic variability enhances the likelihood of selecting superior genotypes. High estimates of genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) on number of productive tillers and grain yield per plant gives us an idea

about the magnitude of variability present in these characters in a genetic population. High heritability along with high genetic advance as percent of mean for all the traits except days to 50% flowering and kernel length indicated additive gene action predicting genetic gain under selection than heritability estimates alone. Therefore, plant breeding efforts should focus on manipulating existing genetic diversity in the desired direction through appropriate crop improvement programs to develop rice hybrid lines with enhanced grain yield.

### References:

1. Akinwale MG, Gregorio G, Nwilene F, Akinyele BO, Ogunbayo SA and Odiyi AC. Heritability and correlation coefficient analysis for yield and its components in rice (*Oryza sativa* L.). *African Journal of plant science*.2011; 5(3): 207-212.
2. Allard RW. Principles of Plant Breeding. John Wiley & Sons, New York, 1960.
3. Burton GW, de Vane EH. Estimating heritability in Tall Fescue (*Festuca arundinacea*) from replicated clonal material. *Agronomy Journal*. 1953; 45: 481-487.
4. Demeke B, Dejene T and Abebe D. Genetic variability, heritability, and genetic advance of morphological, yield related and quality traits in upland rice (*Oryza sativa* L.) genotypes at pawe, northwestern Ethiopia. *Cogent Food & Agriculture*. 2023; 9 (1): 2157099.
5. Federer WT. (1956). Augmented (or Hoonuiaku) designs. *Hawaiian Planters' Record*. 1956; 55: 191-208.
6. Gayathridevi G, Shanthi P, Suresh R, Manonmani S, Geetha S, Sathyabama K, Geetha P. Genetic variability, association and multivariate analysis for yield and yield parameters in rice (*Oryza sativa* L.) landraces. *Electronic Journal of Plant Breeding*. 2023;14(3):991-9.
7. Gupta S, Upadhyay S, Koli GK, Rathi SR, Bisen P, Loitongbam B, Singh PK, Sinha B. Trait association and path analysis studies of yield attributing traits in rice (*Oryza sativa* L.) germplasm. *International Journal of Bio-resource and Stress Management*. 2020;11(6): 508-17.

8. Hemalatha M, Geetha S, Saraswathi R, Raveendran M, Hemalatha G. Diversity and Multi-variate analysis of basmati and non-basmati rice Genotypes over two seasons using organo-leptic traits. *International Journal of Plant & Soil Science*. 2023 Jul 25;35(18):923-41.
9. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans. *Agronomy Journal*. 1955; 47: 314-318.
10. Juliano BO. Rice chemistry and technology (2nd ed.). *American Association of Cereal Chemists*. 1985.
11. Kennedy G, Burlingame B and Nguyen VN. Nutritional contribution of rice and impact of biotechnology and biodiversity in rice-consuming countries. Proceedings of the 20th Session of the International Rice Commission. FAO. 2002.
12. Mythili V, Pradhan B, Dash M and Bastia DN. Genetic variability studies in low land elite rice genotypes for yield and cellulose content. *Journal of Pharmacognosy and Phytochemistry*. 2018; 7(5): 1265-1270.
13. Panse VG, Sukhatme PV. Statistical methods for Agricultural workers. ICAR, New Delhi, 1967
14. Pradhan P, Das B, Kumar D, Thapa B, Chakraborty S, Roy SK, Hijam L, Kundu A. Genetic variability and diversity analysis of rice for yield characteristics across submergence stress and normal condition; 2024.
15. Rashid MM, Nuruzzaman M, Hassan L and Begum SN. Genetic variability analysis for various yield attributing traits in rice genotypes. *Journal of Bangladesh Agricultural University*. 2017; 15 (1): 15–19.
16. Slavin JL. Whole grains and human health. *Nutrition Research Reviews*. 2004; 17(1): 99-110.
17. Uma Devi M, Veerabhadram P, Manonmani S, Shanmugasundram P. Physico chemical and cooking characteristics of rice genotypes. *Electronic Journal of Plant Breeding*. 2010; 1(2):114–123.
18. Vanaja T, Babu LC. Variability in grain quality attributes of high yielding rice varieties (*Oryza sativa* L.) of diverse origin. *Journal of Tropical Agriculture*. 2006; 44: 61-3.

19. Yashwanth JV, Shanthi P, Shekar MR, Reddy VLN and Kumari PL. Estimates of genetic variability, heritability and genetic advance for yield and its component traits in advanced breeding lines of rice (*Oryza sativa* L.). *The Pharma Innovation Journal*. 2021; 10(8): 1422-1425.

UNDER PEER REVIEW