

## Review Article

# Cosmetic, Nutrition, and Drug value of *Carica*

## *papaya* - A short review

### ABSTRACT

Papaya is an evergreen plant which offers different food, nutrition and therapeutic values for human beings. The fruit is delicious and nutritious. The different parts of the plant such as leaves, seeds, peel, blooms, bark, root, and squash have beneficial activities. It is used for the cure of a variety of diseases including digestive disorders, constipation, expulsion of worms, antibacterial, corns, sinuses, skin inflammation, antifertility, dengue fever, warts, glandular tumors, blood pressure, diabetic, anti-HIV, invigorate regenerative organs. It is also used as nutraceutical agent. The present work emphasizes on major characteristics of chemical constituents, nutritional components, health and medical benefits of papaya.

**Keywords:** Papaya, food, nutrition and therapeutic values, antifungal activity, anti-HIV activity, anti-fertility activity, cosmetic value.

## 1. Introduction

Papaya is grown in all areas of the globe. Papaya is considered as the one of the most profitable agricultural produces and it contains protein, starch, vitamins, beta-carotene and minerals. The papaya plant is a little, scarcely expanded plant, the stem developing from 5-10 m tall, leaves are extensive lobed with usually seven projections and about 0.5-0.7m in width. Common components of papaya have smooth skin, green, while ripe papaya becomes yellow (**Figure 1**). The papaya grown in Philippine is usually pear shape around 0.1 to 0.4 m long when become fully mature. Ripe papaya when served feels slippery, the skin has yellowish color, and when opened it has orange to yellowish tissue. Its seeds are black and located in the vacant inner core<sup>1,2</sup>.

## 2. Methodology

Google Scholar, Pub Med and Research Gate etc. have been consulted to find out the needed literatures to have relevant research papers as we have done in our previous works<sup>3-7</sup>. The keywords, "*C. papaya*", "*Food and nutritional value of C. papaya*", "*Drug value of C. papaya*", "*Anticancer activities*", "*Pharmacology of C. papaya*", "*Cosmetic value of C. papaya*" etc. have are used to get expected research papers and accumulate necessary information on different aspects of the papaya and chemical constituents to be used in this review work.

## 3. Findings and Discussion

### 3.1. Plant Taxonomy

*Carica papaya* belongs to the family Caricaceae<sup>8</sup>. Its botanical identification is given below -  
Kingdom: Plantae, Division: Magnoliophyta, Class: Magnoliopsida, Order: Brassicales, Family: Caricaceae, Genus: *Carica*, Species: *papaya* Linn.

As a tropical fruit, its position is fourth in the global trade and commerce. It is extensively cultivated in different parts of the world such as Bangladesh, India, Thailand, Philippines, southern Mexico and Central America. It is called pepe (in Bangla), papita (in Hindi), pawpaw tree (in English), and Erandkarkati (in Sanskrit). Papaya is being used from ancient time as a nutritional and medicinal plant. Papaya is a polygamous species; hard to recognize whether it is female, male, or hermaphrodite. Its fruit differs in size and shape; it widens as globe and has a large central cavity. The seeds of the ripe papaya are black and that of green papaya is white. The plant grows vertically that grows about 5-10 m tall, is a perennial herb with spirally arrayed leaves located to the top of the trunk. The trunk is hollow, brown, or deep purple, and leaf scars

roughen the base. The color of the flowers yellowish-white color, these are small, and numerous in number.



**Figure 1:** The *C. papaya* plant with different parts (Photograph was taken by MSA from Dhaka University Residential area at Fuller Road).

### 3.2. Chemical constituents of *C. papaya*

*C. papaya* contains a lot of chemical components such as antioxidant nutrients (e.g., vitamin C, flavonoids and Carotene), vitamins B (e.g., pantothenic acid), minerals (e.g., potassium and magnesium), and fibers. Papaya leaves contain alkaloids, carpaine, vitamins C and vitamin E, pseudocarpain, caproside, choline and dehydrocarpaine. Fruits possess macromolecules such as carbohydrates, protein, fats, vitamins, minerals, volatile compounds, alkaloids, and glycosides. Its juice contains stearic acid, linolenic acid, myristic acid, palmitic acid, N-butyric acid and n-hexanoic. *C. Papaya* plant contains specialized cells called lactifers and hence the plant is known as lactiferous plant. Latex oozes out from the lactiferous cells and distributes this to all most all tissues of the plant. It has been found that papain is the most vital proteolytic protein (enzyme) extracted from the unripe fruit of the *C. papaya* plant<sup>9</sup>. The outcomes of the test results exhibited that latex components can differ in leaves fruit, and roots. Lactiferous cells fall in number when the fruit ripens<sup>10</sup>. Another report showed a lipase (an enzyme) which was unstable for the necessary lipolysis function in papaya latex<sup>11</sup>. Vitamin A, vitamin E and vitamin C are excellent antioxidants found in *C. papaya* and the minerals K and Mg; the vitamin B includes - B<sub>5</sub> and folate are also found in papaya. It is rich in fibers. The root is abundant in myrosin and

caprosideenzymes. The seeds of papaya is enriched in benzyl isothiocyanate, carpain, fatty acid, protein, fibers, myrosin enzymes, and carcaine<sup>12</sup>. The chemical constituents are shown in **Table 1**.

**Table 1:** Phytoconstituents found in *C. papaya*.

Parts of Plant	Chemical constituents present
Flower	Phalobatanine, Flavonoids
Fruits	Ascorbic acid, Fibers, Sugar, Protein, minerals, phosphorus, iron, Vit C, thiamine, niacin, riboflavin, citric acid, malic acid, carotene, amino acid, benzyl isothiocyanate, cis and trans 2.6-dimethyl-3.6epoxy-7-octen 2-ol.
Leaves	Papain, cystatin, tocopherol, carpain, chymopapain, dihydrocarpaine I & II, carposide, Vitamin C and E.
Seed	Benzyl isothiocyanate, fatty acid, protein, fibers, carpain, myrosin enzyme and carcaine
Root	Myrosin, Caproside.

### 3.3. Food and nutritional features of *C. papaya*

Papaya is an excellent hub of food and nutrition as it contains many of the elements of basic food needed for human being. It's calory value is low butenrich in necessary minerals and vitamins. The comparative low calory content helps the obese people to reduce their weight. The nutritional values<sup>13,14</sup> of *C. papaya* has been shown in **Table2**.


**Table 2:** Nutritional elements and their values in *C. papaya* in 100g of papaya fruit both green and ripe.

Nutrients	Green papaya	Ripe papaya
Proteins	600mg	700mg
Fat	100mg	200mg
Mineral	500mg	500mg
Carbohydrate	720mg	570mg
Fiber	800mg	900mg
Energy	32Kcal	27Kcal
Total carotene	2740µg	-

Beta- carotene	888µg	-
----------------	-------	---

### 3.4. Pharmacological properties of *C. papaya*

A schematic diagram (Figure 2) summarizes the pharmacological properties of *C. papaya*. A brief description is also given in the following sections.



**Figure 2:** Pharmacological activities of *C. papaya* at a glance.

#### 3.4.1. Effect of *C. papaya* on blood-related issues

Different parts of *C. papaya* show activity against blood related diseases. One such study with seed and leaf explored the phytoconstituents and hematological activity of *C. papaya*<sup>15</sup>. Another study showed that *C. papaya* improved the formation of certain blood parameters in albino rats<sup>16</sup>.

#### 3.4.2. Effect of *C. papaya* on digestive disorder

*C. papaya* showed beneficial activity against different ailments of digestive disorder. The astringent ashes of the unripe peel and of the leaves are taken in dysentery and diarrhea. The roots are administered in digestive disorders, dysentery and other ailments. The *C. papaya* preparation also showed beneficial effect in the symptoms of acid reflux<sup>17</sup>.

#### 3.4.3. Hyperglycemia and ethanolic extract of *C. papaya*

The phytoconstituents of *C. papaya* showed antihyperglycemic activity. *C. papaya* was extracted by ethanol and the obtained extract was administered to streptozotocin-induced diabetic rats and

blood sugar was measured. It was seen that *C. papaya* exhibited hypoglycemic activity in diabetic rats. This experiment concluded that the phytoconstituents present in *C. papaya* could offer anti-diabetic action<sup>18,19</sup>.

#### **3.4.4. Effect of *C. papaya* on dengue fever**

*C. papaya* leaves have beneficial effects on recovering from dengue fever which is caused by the bite of female anophelid mosquitoes during day time. A group of dengue fever patients were given 25 mL of leaf extract of *C. papaya* for two times a day and the patients recovered from dengue fever as evidenced by rising of platelet counts from 46.0% to 78.3%<sup>21</sup>.

Cyclophosphamide-induced thrombocytopenic rats showed an increase in thrombocyte after administration of 400 mg/kg and 800 mg/kg of *C. papaya* leaf extract for fifteen days<sup>22</sup>. It was thus concluded that *C. papaya* leaves have a positive impact in curing dengue fever.

#### **3.4.5. Antifungal activity**

Fungicidal actions of *C. papaya* have been studied by Chavez-Quintal *et al.* and they have shown that the extracts exhibited blocking of fungal activity against three phytopathogenic growths -*Rhizopus stolonifer*, *Fusarium* spp., and *Colletotrichum gloeosporioides*<sup>23</sup>. Another report showed the fungicidal activity of the methanolic extract of the seeds and the extract imparted fungicidal actions against *Aspergillus flavus*, *Candida albicans*, and *Penicillium citrinum*.

#### **3.4.6. Safety and side effects of papaya on pregnancy**

The activity of ripe papaya preparation (500 ml/l water) and latex of papaya was studied on pregnant rats (Sprague Dawley species). It was found that ripe papaya has no threatening effect but the latex induced shrinking of the muscles of the uterus. It was inferred that normal use of ripe papaya has little impact on pregnancy. The unripe or semi-ripe papaya might not be safe in pregnancy<sup>24</sup>.

#### **3.4.7. Antihelminthic and antiamebic activity of *C. papaya* seeds**

The beneficial effect of powdered and dried *C. papaya* seed was examined against human intestinal parasitosis<sup>25</sup>. The dried powder of *C. papaya* was given to the patients divided into two groups - blended with nectar and nectar alone as control. It was seen that 76.7% of the patients who have been administered nectar mixed with powder of seeds of papaya eliminated the parasites within seven days, on the other hand parasites were eliminated from only 16.7% of the patients who have been administered nectar alone. This study indicated that shade-dried *C.*

*papaya* seeds are able to treat human intestinal parasites and there are no major critical adverse effects.

#### **3.4.8. *C. papaya* and its effects on cancer and immune system**

Few studies reported the anti-cancer and immune adjuvant property of leaf extract of *C. papaya*. The studies were done on tumor cell lines<sup>26,27,33</sup>. A compound 'papain' found in Papaya which is considered to have activity against cancer growth. It also contains 'lycopene' which gives free radical scavenging activity. Papaya contains another compound 'Isothiocyanate' which has strong action against lung, pancreas and prostate cancer. These compounds prevent DNA re-arrangement process and healing of diseased cell.

#### **3.4.9. Antibacterial and wound healing effect of *C. papaya***

A study on *C. papaya* demonstrated the antibacterial property of seed extract against *Streptococcus faecalis*, *Bacillus cereus*, *Escherichia coli*, *Proteus vulgaris*, *Staphylococcus aureus*, and *Shigella flexneri*<sup>28</sup>. This study showed that *C. papaya* has antibacterial effects which could be beneficial in treating skin ulcers. Another study showed antibacterial action of the papaya latex against *Agrobacterium* sp., *Bacillus subtilis*, *E. coli*, and *Rhizobium* sp<sup>29</sup>.

#### **3.4.10. Antifertility activity**

A study by Poharkaret *al.* focused light on the impacts of *C. papaya* on grown-up and pregnant rodent which were administered different parts of the natural product<sup>30</sup>. The animals were allowed to take the papaya mixed-food normally and the results exhibited that the green papaya fruit interrupted with the estrous cycle and fetal expulsion. This effect disappeared when ripe fruit was given to the animals. Chloroform fraction of *C. papaya* seeds showed azoospermia in monkey. Moreover, papaya indicated the counter implantation and abortifacient action.

#### **3.4.11. Hepatoprotective effect**

Hepatotoxicity was induced by CCl<sub>4</sub> in rats by Sadequeet *al.* and they applied dried natural products of papaya which showed hepatoprotective action. This action supported the action of Vitamin-E<sup>31</sup>. The results indicated that *C. papaya* and Vitamin E showed excellent hepatoprotective action against CCl<sub>4</sub>-induced hepatotoxicity. However, *C. papaya* showed more subtle alterations in alkaline phosphatase level than Vitamin E. In another study, the ethanol and aqueous extracts of *C. papaya* exhibited impressive hepatoprotective activity against CCl<sub>4</sub> induced hepatotoxicity<sup>32</sup>.

#### **3.4.12. *C. papaya* male flower and its effects as Antioxidant**

The antioxidant functions of n-hexane fraction of male flower part was studied by Sianiparet *al*<sup>34</sup>. The antioxidant property was tested by using the  $\alpha,\alpha$ -diphenyl- $\beta$ -picrylhydrazyl method. Phytochemical screening of the n-hexane part contains triterpenoids and steroids.

#### **3.4.14. Antimicrobial activity of *C. papaya***

A study investigated the phytoconstituents and antimicrobial activities of various parts of *C. papaya*<sup>35</sup>. The antimicrobial activities were tested against the Gram-positive and Gram-negative bacterial strains and growth by observing the zone of inhibition. The strains of Gram-positive bacteria were *S. aureus*, *B. cereus*, and *Micrococcus luteus*, and that of Gram-negative bacteria were *E. coli*, and *Klebsiella pneumoniae*. The fungus applied in the test were *Aspergillus niger*, *A. flavus*, *C. albicans*, *Candida tropicalis*, *Cryptococcus neoformans* and *Candida kefyr*. Sumathi<sup>36</sup> studied the *in vitro* antimicrobial action of *C. papaya* against pathogens. Among the different parts of *C. papaya*, leaf extract showed greater antimicrobial action than stem and root. Cold water, hot water and ethanol were used as solvent to extract the phytochemicals from different parts of *C. papaya*.

#### **3.4.15. Activity of *C. papaya* against HIV**

Phytochemical screening and the evaluation of papaya extract against HIV-1 was done Rashed *et al.* (2013)<sup>37</sup>. The methanol and water extract of *C. papaya* were used by applying the syncytia advancement test to find out activity of papaya against HIV-1. The result indicated that *C. papaya* have limited action against HIV-1.

### **4. Use of *C. papaya* in Cosmetics**

Papaya can be used in cosmetics for its skin-brightening, exfoliating, and moisturizing properties<sup>35</sup>. Some of the uses are – (i) **Face mask** - Mash papaya and mix with honey to create a face mask that can help remove tan and lighten skin, (ii) **Body scrub** - Blend papaya flesh with papaya seeds and olive oil to create a body scrub that exfoliates, (iii) **Facial** - Mash papaya with turmeric to create a facial that can give your skin a natural glow, (iv) **Hand soap** - Some hand soaps contain papaya to soften and cleanse hands.

Papaya is rich in vitamins and enzymes, including papain, which can help improve the health and appearance of the skin, for example – (i) **Exfoliation** - Papain removes dead skin cells and promotes cell renewal, (ii) **Skin tone** - Papaya can help reduce the appearance of blemishes, acne scars, and uneven skin tone, (iii) **Antioxidants**– Papaya's vitamin C content helps protect skin against free radicals, environmental pollutants, and UV damage, (iv) **Hydration**– Papaya's

vitamin E provides deep hydration and nourishment. But before using papaya for the skin or hair, a doctor or dermatologist should be consulted.

## 5. Conclusion

Fruits part of papaya plant is mainly consumed as the food component all over the globe for its food, nutritional, cosmetic and medicinal properties. Different parts of papaya plant such as leaves, seeds, roots, blossoms, ripe and unripe fruit juices show these properties. Its pharmacological activities indicate that it might be good pharmaceutical product if processed properly. Therefore, critical measure of work on the uses of its constituents, appropriate institutionalization, clinical trials are expected to explore the healing utility to fight diverse ailments by applying *C. papaya*.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## References

1. Oliver-Bever B. Medicinal Plants in Tropical West Africa. Cambridge: Cambridge University Press; 1986. p. 161.

2. Yogiraj V, Goyal PK, Chauhan CS, Goyal A, Vyas B. *Carica papaya* Linn: An overview. Int J Herb Med., 2014, 2:1-8.
3. Md Rafat Tahsin, Tanzia Islam Tithi, Sabiha Rahman Mim, Ehfazul Haque, Arifa Sultana, Nasiba Binte Bahar, Raju Ahmed, Jakir Ahmed Chowdhury, Abu Asad Chowdhury, Shaila Kabir, Fahima Aktar, Md Sahab Uddin, Md Shah Amran, In Vivo and In Silico Assessment of Diabetes Ameliorating Potentiality and Safety Profile of *Gynura procumbens* Leaves, Evidence-Based Complementary and Alternative Medicine, 2022 (1), 9095504.
4. Nusrat Jahan Shawon, Kakoli Sutradhar, Shopnil Akash, Jakir Ahmed Chowdhury, Abu Asad Chowdhury, Shaila Kabir, Md Shah Amran, In vivo and In silico Analysis of Antihypertensive Activities of *Ficus religiosa* Fruit Extract, Dhaka University Journal of Pharmaceutical Sciences, 2024, 23(1), 23-36.
5. M.S. Hossain, M.R.I. Khan, A.S.M. Anisuzzaman, M. Ahmed, M.S. Amran and A. Islam, Antidiabetic and Glycogenesis Effects of Different Fractions of Ethanolic Extract of leaves of *Mangifera indica* (Linn.) in Normal and Alloxan Induced Diabetic Rats, *Journal of Medical Sciences*, 2010, 10 (4), 80-86.
6. S Shermin, F Aktar, M Ahsan, CM Hasan. Antioxidant and Cytotoxic Activity of *Limonia acidissima* L. Dhaka University Journal of Pharmaceutical Sciences, 2012, 11 (1), 75-77
7. Marufa Islam, AH Rupak, Najibah Nasrin, Md Mustafiz Chowdhury, Priyanka Sen, Ahammad Ullah Foysal, Md Jasim Uddin, Jannatul Ferdous, Md Rafat Tahsin, Fahima Aktar, Shaila Kabir, Abu Asad Chowdhury, Md Shah Amran, Tahmina Akter, Jakir Ahmed Chowdhury. An evaluation of potential hepato-protective properties of *Hylocereus undatus* fruit in experimental rat model. *Biomedical Journal of Scientific & Technical Research*, 2022, 43 (2), 34405-16
8. Pigli RK, Runja C. Medicinal plants used in dengue treatment. Int J Chem Nat Sci., 2014, 2:70-6.
9. Hettige S. Salutary effects of *Carica papaya* leaf extract in dengue fever patients—a pilot study. *Sri Lankan Fam Physician* 2008, 29:17-9.
10. Yunita F, Hanani E, Kristianto J. The effect of *Carica papaya* L. Leaves extract capsules on platelets count and hematocrit level in dengue fever patient. Int J Med Aromat Plants 2012, 2:573-8.

11. Manohar PR. Papaya, dengue fever and ayurveda. *AncSci Life*, 2013, 32:131-3.
12. Senthilvel P, Lavanya P, Kumar KM, Swetha R, Anitha P, Bag S, *et al.* Flavonoid from *Carica papaya* inhibits NS2B-NS3 protease and prevents dengue 2 viral assembly. *Bioinformation* 2013, 9:889-95.
13. Sarala N, Paknikar S. Papaya extract to treat dengue: A novel therapeutic option? *Ann Med Health Sci Res.*, 2014, 4:320-4.
14. Dharmarathna SL, Wickramasinghe S, Waduge RN, Rajapakse RP, Kularatne SA. Does *Carica papaya* leaf-extract increase the platelet count? An experimental study in a murine model. *Asian Pac J Trop Biomed* 2013, 3:720-4.
15. Kasture PN, Nagabhushan KH, Kumar A. A multi-centric, double blind, placebo-controlled, randomized, prospective study to evaluate the efficacy and safety of *Carica papaya* leaf extract, as empirical therapy for thrombocytopenia associated with dengue fever. *J Assoc Physicians India* 2016, 64:15-20.
16. Ikpeme EV, Ekaluo UB, Kooffreh ME, Udensi O. Phytochemistry and hematological potential of ethanol seed leaf and pulp extracts of *Carica papaya* (Linn.). *Pak J Biol Sci.*, 2011, 14:408-11.
17. Jaiswal P, Kumar P, Singh VK, Singh DK. *Carica papaya* Linn: A potential source for various health problems. *J Pharm Res.*, 2010, 3:998-1003.
19. Sasidharan S, Sumathi V, Jegathambigai NR, Latha LY. Antihyperglycaemic effects of ethanol extracts of *Carica papaya* and *Pandanus amaryfollius* leaf in streptozotocin-induced diabetic mice. *Nat Prod Res.*, 2011, 25:1982-7.
20. Maniyar Y, Bhixavatimath P. Antihyperglycemic and hypolipidemic activities of aqueous extract of *Carica papaya* linn. Leaves in alloxan-induced diabetic rats. *J Ayurveda Integr Med.*, 2012, 3:70-4.
21. Ahmad N, Fazal H, Ayaz M, Abbasi BH, Mohammad I, Fazal L, *et al.* Dengue fever treatment with *Carica papaya* leaves extracts. *Asian Pac J Trop Biomed.*, 2011, 1:330-3.
22. Patil S, Shetty S, Bhide R, Narayanan S. Evaluation of platelet augmentation activity of *Carica papaya* leaf aqueous extract in rats. *J PharmacogPhytochem.*, 2013, 1:227-32.
23. Chávez-Quintal P, González-Flores T, Rodríguez-Buenfil I, Gallegos-Tintoré S. Antifungal activity in ethanolic extracts of *Carica papaya* L. Cv. Maradol leaves and seeds. *Indian J Microbiol.*, 2011, 51:54-60.

24. Adebiyi A, Adaikan PG, Prasad RN. Papaya (*Carica papaya*) consumption is unsafe in pregnancy: Fact or fable? Scientific evaluation of a common belief in some parts of Asia using a rat model. *Br J Nutr.*, 2002, 88:199-203.
25. Okeniyi JA, Ogunlesi TA, Oyelami OA, Adeyemi LA. Effectiveness of dried *Carica papaya* seeds against human intestinal parasitosis: A pilot study. *J Med Food*, 2007, 10:194-6.
26. Otsuki N, Dang NH, Kumagai E, Kondo A, Iwata S, Morimoto C, *et al.* Aqueous extract of *Carica papaya* leaves exhibits anti-tumor activity and immunomodulatory effects. *J Ethnopharmacol.*, 2010, 127:760-7.
27. Fauziya S, Krishnamurthy R. Papaya source material for anticancer. *CIB Tech J Pharm Sci.*, 2013, 2:25-34.
28. Dawkins G, Hewitt H, Wint Y, Obiefuna PC, Wint B. Antibacterial effects of *Carica papaya* fruit on common wound organisms. *West Indian Med J.*, 2003, 52:290-2.
29. Islam A, Al-Mamun MA, Parvin S, Meh S. Evaluation of antibacterial activity of Latex of *Caricaceae*. *Asian J Pharm Cli Res.*, 2015, 8:308-11.
30. Poharkar RD, Saraswat RK, Kotkar S. Survey of plants having antifertility activity from Western Ghat area of Maharashtra state. *J Herb Med Toxicol.*, 2010, 4:71-5.
31. Sadeque MZ, Begum ZA, Umar BU, Ferdous AH, Sultana S, Uddin MK. Comparative efficacy of dried fruits of *Carica papaya* Linn. And Vitamin E on preventing hepatotoxicity in rats. *Faridpur Med Coll J.*, 2012, 7:29-32.
32. Raj Kapoor B, Jayakar B, Kavimani S. Effect of *Carica papaya* on hepatotoxicity. *Biol Pharm Bull.*, 2002, 25:1645-6.
33. Praveena P, Jethinlalkhosh JP, Doss VA. Pharmacological evaluation of antineoplastic activity of hydroethanolic extract of unripe fruit of *Carica papaya* Linn using animal model. *Asian J Pharm Clin Res.*, 2017, 10:179-81.
34. Sianipar MP, Suwarso E, Rosidah R. Antioxidant and anticancer activities of hexane fraction from *Carica papaya* L. Male flower. *Asian J Pharm Clin Res.*, 2018, 11:81-3.
35. Baskaran C, Bai VR, Velu S, Kumaran K. The efficacy of *Carica papaya* leaf extract on some bacterial and fungal strain by well diffusion method. *Asian Pac J Trop Dis.*, 2012, 2012:347-9.
36. Sumathi R. Phytochemical analysis and *in vitro* antimicrobial activity of aqueous and solvent extracts of *Carica papaya* against clinical pathogens. *Int J Adv Res Biol Sci.*, 2014, 1:73-7.

37. <https://www.clinikally.com/blogs/news/unlocking-the-power-of-papaya-enzyme-in-skincare-regimens>.(Accessed on 20-10-2024).

UNDER PEER REVIEW