

FOREST ECOSYSTEM MANAGEMENT AND CLIMATE CHANGE

ABSTRACT: Climate change is one of the world's greatest challenges. ~~W~~hich creates effects on our forest ecosystem. Sensitivity of forest ecosystem and climate significance affects the growth rate and structure of the forest, and it also ~~have a direct impacts affectson~~ species ~~compositiondistribution~~, growth of trees, their mortality and reproduction of trees is also affected by climate change. Increase and decrease in temperature leads to change in time phenology due to which earlier burst of bud, leafing and flowering in trees is affected, this review is all about how climate change affects forest ecosystem. All direct and indirect alterations and shifts in the ecosystems are affected by change in climate. Rise in temperature will affect respiration process due to which photosynthesis rate will also change. Disturbance in climate change. Affects carbon cycle, forest structure, composition of species and alteration in forest ecosystem. Disturbance includes; invasion in species, forest fire, insects and disease outbreaks. Disease outbreaks destroy the structure, composition and function of forest ecosystems. Forest fire also disturbs our forest ecosystem by disturbing wildlife, habitat, increase and decrease in nutrient cycling and mortality of individual. Trees frequency, size, seasonality, intensity due to fire multiplication of pests and pathogen is faster. They expand their range and invade into new regions under warmer, wetter, or carbon dioxide enriched conditions. Nutrient availability is affected by an increase in the temperature through change of organic matter decomposition and mineralization of soil nutrients, mainly nitrogen.

Keywords: Carbon sequestration, Climate Change, carbon, Forest Ecosystem, Nitrogen.

INTRODUCTION


A functional unit or a system comprising soil, tree and other living things is termed as forest ecosystem. About 4.1 billion ha of global land surface is covered by forest (Hisano et al. 2018). Forest ecosystems provide many eco-services like timber and non-timber products and play a key role in contribution to the world economy (Hui et al. 2017). Other than this, the forest provides many other services which include habitat for wildlife, clean water, carbon storage & climate action (Lindquist et al. 2012). The collection and reserving of carbon in this carbon cycle is termed carbon sequestration ~~reference~~. A major pool of carbon is forest biomass and soil carbon sequestration ~~reference~~. About 359 million tons is the total carbon sequestered in forest vegetation (Hobbset al. 2016). Carbon cycling and curbing climate change is majorly affected by forest. The world's biggest threat is climate change. Due to climate changes many activities that are harmful for forest ecosystems like fossil fuel burning, deforestation carbon dioxide level has been increased from 280 ppm to 379 ppm (Penuelas et al. 2013). Human activities contribute mostly to forest loss, fragmentation, and degradation (FAO 2015) ~~this and this effect~~ led to loss of forest ecosystem (Var der pals et al. 2016) ~~h~~ However, forest ecosystem and climate are interrelated. The impact of climate change on forest ecosystems primarily focuses on their roles in carbon emissions and sequestration. (Mori et al. 2017). Increases in disturbance severity could dramatically impact forest ecosystem structures, species composition, and their capacity to function as carbon sinks. (Cardinale. 2012). ~~And the increase in disturbance is due to Climate change. It Climate change can have a direct~~ly impact on nutrient cycling ~~throughby~~ affecting ~~sing on~~ temperature and precipitation. A gradual rise in temperature will ~~affect impact the regeneration and~~ growth ~~and establishment~~ of certain species. ~~Climate change majorly affects forests and their services flow~~. Global climate changes consequences that include long term droughts limited by the availability of nitrogen; a fundamental element essential for plant growth. Nitrogen often limits forest growth under most conditions. However, excessive nitrogen can negatively affect soil and trees, and can also impact forest composition, growing season length, and the water cycle. Increased temperatures can hinder the regeneration of tree species by reducing soil moisture and increasing competition during the seedling stage. Climatic variability and change lead to the degradation of forest resources, resulting in higher carbon dioxide emissions and adversely affecting the forest's ability to provide its ecosystem services. Whereas high CO₂ emission results in increasing temperature, affecting nutrient availability in the soil through stimulation of organic matter decomposition and mineralization of soil nutrients. We can perform some conservation practices like afforestation (It is defined as and imbalances in water supplies; decreasing carbon sequestering capabilities,

soil erosion, loss of soil fertility, desertification, reduction in food production capacity (Isbell et al. 2015). Decrease in precipitation and higher temperature will lead to increase in arid land which is a major threat to biodiversity and survival of certain forests ecosystem (Zuidema et al. 2013) The revival of these harmful impact is costly or irreversible for ecosystems and biodiversity (Peters et al. 2013). Ecosystem is the interaction between living organism and their physical environment within a particular area is defined as ecosystem and whereas on other hand, Biodiversity is defined as the variety and variation of living form under an ecosystem therefore biodiversity and ecosystem are interrelated. Biodiversity and forest ecosystem conservation is increasing concern and demand worldwide to cope with the declining biodiversity and forest ecosystem. (Tian et al. 2012; Gebeyehu et al. 2019). Conservation of biodiversity and forest ecosystems will also help in managing climate changes. It also changes the frequency or intensity of natural disturbances in certain forests, especially those caused by fire and insects. These changes will reduce the ability to mitigate harmful effects, and will reduce human wellbeing, particularly for the poorest populations of low-income countries and arid climate regions (Dukes et al. 2014). The productivity and health of forest ecosystems are closely connected to nutrient availability, and climate change like precipitation pattern can alter nutrient dynamics by modifying the rate of litter decomposition. The ability of ecosystems to absorb carbon is creating a new forest by planting more and more trees) because trees and soil nutrient availability are interrelated to each other. The biochemical cycling between the soil and plants is one of the main factors of forest ecosystem functioning. Climate variation affects readiness of nutrients for plant uptake and limits the output and persistence of Forest ecosystems. Nutrient readiness affects forest ecosystem and net energy production (Juneja et al. 2014). Nitrogen can be sourced from biomass broken down by microbes (e.g.: fallen leaves release nitrogen to the soil). Boosting the nitrogen requisition is essential for plant growth, development and builds protein. However, boosting carbon dioxide tier in the atmosphere will lead to global warming, further leading to climate changes. The yield of forest ecosystem appears to be governed by quantity and readiness of nutrients. Temperature boost has direct physical reaction effects on rate photosynthesis, macronutrient dosage in soils and leaves of trees, shrubs and herbs. boost in temperature can lead to reductions in nutrient readiness (Tian et al. 2013). Climate change poses significant threats to forests and the vital services they provide. Expected impacts include prolonged droughts and disruptions in water availability, reduced carbon sequestration capabilities, soil erosion, decreased soil fertility, desertification, diminished food production potential, and increased risks from certain species and ecosystems.

FOREST ECOSYSTEM

Forests are the largest terrestrial ecosystems on Earth, covering over 4.1 billion hectares (Pohjanmies et al. 2013). They offer habitats for most of the planet's species and supply crucial ecosystem resources and services to humans, including food, fiber, timber, clean water, carbon storage, and climate regulation (an et al. 2022).

Figure 1 FAO. 2020. *Global Forest Resources Assessment 2020*. Rome. <https://www.fao.org/forest-resources-assessment/fra-2020/maps/en/>



FOREST ECOSYSTEM SERVICES:

Environmental services are the direct and indirect participation that ecosystems make to human well-being. These benefits encompass supplying, managing, cultural, and supporting services, all of which are essential for

sustaining other ecosystem functions.(May et al. 2015) Forests are widely recognized as key providers of ecosystem services (Peñuelaset al. 2017). National classifications often identify up to 100 distinct environmental services, including food, timber, and fuel production; water conservation and regulation; nutrient retention; carbon sequestration; biodiversity protection; climate regulation; ecotourism; and the preservation of spiritual and traditional values. Biodiversity- ecosystem functioning (BEF) (Chitsazet al .2012) show that biodiversity is related to functionality of ecosystem(Example – primary production , decomposition , nutrient cyclingEtc.) and also provide ecosystem services (e.g. – food production , climate regulation ,pollinationand other) .Experimental and theoretical work has developed BEF theory with the contribution from experimental manipulation of plant assemblages in grassland ecosystem over the last several decades (Mitrică et al. 2015) .

NUTRIENT CYCLE AND CLIMATE CHANGES:

Climate crisis is becoming a great warning to the world. It is leading to a boost in global surface temperature at an uncontrolled rate. Increase in temperature, CO2 level will affect rainfall patterns, nitrogen deposition will affect the tree growth and forest dynamics and majorly the accumulation of carbon in forest ecosystem (Damkjaeret al.2013).It is crucial for predicting ecosystem carbon dynamic in forest ecosystem. Carbon sequestration can be defined as the process of storing carbon in a carbon pool. Carbon sequestration majorly contributes to limiting climate change. This process helps in controlling climate changes by reducing the amount of carbon dioxide in the atmosphere. Forest ecosystems can either act as carbon sinks, sequestering more carbon, or as carbon sources, releasing more carbon, depending on the net carbon exchange.

The effects of climate crisis on forests have shown both positive and negative impacts. On the positive side, elevated CO2 levels can lead to increased forest growth and improved water use efficiency. Conversely, negative effects include reduced growth, which can arise from a combination of factors such as higher temperatures, altered precipitation patterns, and increased frequency of extreme weather events.




Figure 2 Countries with required inclusion of soil and water management, stakeholder inputs and high conservation value forest designation in forest management plans.(Macdicken et al. 2015)

IMPACT OF NITROGEN IMPEACHMENT ON CARBON SEQUESTRATION BY FOREST ECOSYSTEM:




Figure 3. Carbon sequestration in forest ecosystems, including major carbon pools and fluxes (Hui et al. 2015)

Over the past 25 years assessments of global carbon (C) budget show that more than 50 % of the human activities CO₂ emission is stored in oceans and land. For oceans and terrestrial ecosystems, the most recent global estimate of C sinks is $2.6 \pm 0.5 \text{ Pg C yr}^{-1}$ and $2.6 \pm 0.8 \text{ Pg C yr}^{-1}$ respectively (Corinne Le Quéré et al. 2013). The storage of human-caused CO₂ emissions mainly happens in forest ecosystems. (Pan et al. 2013). For the prediction of long-term henceforward global forest sink, it is important to have insight of interaction between environment of the processes that determines the forest C balance. Other than site disruption and forest management many field studies, and models have been approached that suggest increasing atmospheric CO₂ and climatic warming. They are the major driver of continental and global carbon sinks. However, the significant disruption of the nitrogen (N) cycle since the early nineteenth century has led to increased atmospheric nitrogen deposition in forests. There is growing evidence that this has also significantly boosted forest carbon sequestration. (Büntgen et al. 2014). While most of the forest ecosystems are nitrogen deficient, increased nitrogen deposits will increase net primary production (NPP) thus stimulating carbon sequestration in trees and it will lead to declining biodiversity.

Increased productivity can also enhance carbon sequestration in the soil through greater soil carbon inputs from litterfall (de Vries et al. 2011) and reduced decomposition of organic matter. There is substantial evidence that nitrogen availability, and more broadly nutrient availability, is crucial in how forest ecosystems respond to higher CO₂ concentrations, elevated temperatures, and altered water availability (Lu et al. 2014)

DIFFERENT FACTORS EFFECTING CARBON SEQUESTRATION

- 1) Consequences of CO₂
- 2) Consequences of temperature
- 3) Consequences of precipitation
- 4) Consequences of O₃

Consequences OF CO₂


The effect of increasing atmospheric CO₂ levels on carbon sequestration in forest biosphere has been researched for decades. Natural ecosystem and socio-economic systems are impacted by any global change in climate. Since the industrial revolution, many human activities like burning fossil fuels have led to a boost in the concentration of greenhouse gases in the atmosphere causing temperature rise which is defined as global warming. (Fernández-Martínez et al. 2020). Warming has seriously altered justifiable development, and climate crisis has been listed in the top 10 ecological problems faced by human beings (Huanget al. 2019). In early years,

many researchers used closed chambers, glasshouses and growth chambers to grow tree seedlings or small trees under controlled condition. As glasshouse and growth chambers are artificial mean of growing tree seedlingsit cannot really copy the real-world environment so these practices of growing under controlled condition are criticized by many other researchers because these gave different results as under natural conditions. Glasshouses have an open top chamber which provides natural light and precipitation which may maintain air and soil temperature inside the chambers. Recently, free air Co2 enrichment (FACE) technology made it feasible for trees to artificially provide exposure to higher Co2 levels. This technique involves the treatment of entire forest ecosystems with CO2 enriched air in large scale,unrestricted plots, eliminating the need of enclosed structure (glasshouse,polyhousesetc.) (Drake et al. 2011)

CONSEQUENCES OF TEMPERATURE

9,000 soil profile reveals that carbon sequestration is affected by high annual mean temperature, importantly it is common in coarse - textured soils, this soil has less capacity to stabilize organic matter as compared to fine textured soils with better stabilization properties. This trend is constant across both cool and warm regions and affecting factors like plant yield, rain,aridity,cation exchange capacity and pH. The outputs are represented by an accepted earth system model (ESM).Warming will promote real soil losses, but ESMs may not be able to predict the losses accurately. (Hartley et al. 2021). Low soil temperature in cold and wet regionsencourages theemergence of thick organic horizons and peats, resulting in storage of carbon i.e.~500 Pg of C (Z.C.Yu2012).

FIGURE 4.Underground carbon budgets for a warm-temperate forest exposed to elevated atmospheric CO2 at the Duke free-air CO2 enrichment (FACE) site. Ambient CO2 is shown in white, while elevated CO2 is shown in black. Ovals reflect pools and have units of g C m⁻²; squares within arrows reflect fluxes of C and have units of g C m⁻² year⁻¹(Drake et al. 2011)



soils are mixture of organic and mineral material, which is further referred as mineral soil, this soil contains two third of the global soil C stock, the role climatic variables in controlling C storage have been frequently questioned (Doetterl et al. 2015). In the inadequacy of anaerobic environment, decomposers will affect long term persistence of soil organic matter (SOM) in mineral soil (Dungait et al. 2012). The study of soil profile is significant because soil properties are important for C stock judging, role of temperature in determining C storage may be hyperbolize, which limits the potential of release of C from soils as the climate warms. (Dominguez et al. 2019). Whereas physiochemical stabilization mechanisms are considered temperature sensitive. For example, sorption (stabilization) and desorption (destabilization)are argued to be both more and less sensitive reactions, with the relative temperature sensitivities varying with harmony of organic matter surfaces (Wiesmeier et al. 2011). Evidence shows that the decomposition of older and more protected SOM is more sensitive to temperature than rapidly increasing SOM, showing that destabilization reactions are highly temperature reactive (Conant et al. 2020). Thus, it is controversial to identify that at which extent of temperature C storage controls in mineral soils, and it is not known whether the C stored in mineral soils with large capacities stabilizes more carbon or less vulnerable to climate warming than the C stored in soils with more limited stabilization capacities.

CONSEQUENCES OF PRECIPITATION


15 % of the universal ground cover, in which 14.4% of global population is supported represents transitional section between arid and semi humid regions (Jia et al. 2024). Low soil water storage capacity, sparse vegetation cover, and smashable ecosystems, these regions exhibit sensitivity to climate change and human activities (Gong et al. 2024). Over the past century, Semi-arid regions have undergone noticeable warming which accounts 44% of the universal average annual surface warming trend (Wanget al. 2014). The increase in universal warming, serious drought poses important threats to the carbon sequestration capacity and biodiversity in terrestrial ecosystems within semi-arid areas (Huanget al.2012), this projection indicates the threat will increase warming in future (Pal et al. 2023) Semi-arid regions, as current scientific hotspots in global climate change (Knowles et al. 2020), have significant carbon stocks from SOC due to a range of reasons. Therefore, to propose feasible strategies and protect this fragile ecological environment accordingly requires understanding of the characteristics of semi-arid regions and how they will respond under climate change. Abstract Global warming is thought to intensify hydrological processes within terrestrial ecosystems. This resulted in a broad increase in total global PRE and heightened occurrence rates of extreme PRE, especially over semi-arid regions (Ahlström et al. 2015). PRE limits the physiological processes of vegetation (Ameer et al., 2023) and grassland-dominated semi-arid ecosystems show a high sensitivity in photosynthesis carbon uptake to varying PRE strength (Chenet al. 2021). The longevity and amount of PRE affect land surface ecological processes, such as plant physiology development or soil moisture condition which indirectly modify the carbon flux strength and direction (Jayara et al., 2023). Vegetation growth can be promoted by an increase in PRE in water limited terrestrial ecosystems. This is only possible when there is a high connection between PRE and vegetation (Zhang et al. 2024). Many studies demonstrated PRE as enhancer of carbon absorption in ecosystem (Zhan et al. 2023), Furthermore, in most of semi-arid region's groundwater is deep and relatively disengage from soil moisture. Therefore, resulting in boost in facile soil water content (SWC) which primarily depends on vertical drainage during PRE events (ao et al. 2017). Near surface and root zone) (Liu et al. 2024). Enhancement in photosynthesis, boost in carbon absorption in vegetation, carry in microbial activity and organic matter decomposition in the soil is done by adequate SWC levels (Liu et al. 2019), soil respiration generates carbon dioxide pulses which are commonly noticed in rewetting of seasonally dry soil after PRE in semi-arid regions (Wanget al. 2023). Therefore, controlling ecosystem processes and land atmosphere interactions is controlled by PRE.

CONSEQUENCES OF O₃

Ozone can have many effects on forest ecosystems. Increased human activities increases tropospheric ozone abundance. It is widespread like a pollutant in regional air which leads to decrease in rate of photosynthesis and decrease in growth of plant (Prajapati et al., 2012). The probability of O₃ becoming more abundant in future is very high resulting in limiting terrestrial carbon sequestration across broader portions of earth. The impact of O₃ on the forest carbon cycling is estimated by using coupled climate biogeochemical cycling model. They are parameterized by using the physiological responses observed during seedling and saplings. The few ecosystem-scale forest O₃ experiments that have been conducted (Ainsworth et al., 2012) have revealed that small plants responses cannot always be extrapolated to larger scale. It shows that when level of CO₂ increases physiological response of O₃ can be modified by environmental and biological interaction. As if the understanding of these interactions is comparatively poor for O₃ and ecosystem scale. Therefore, there is a critical need for research on the impact of O₃ on forest carbon cycling remains constant. As for our knowledge of the interactive effects of CO₂ and O₃ on the forest, carbon cycling is poorly developed. The Aspen face experiment was framed to understand how ecosystem works. In developing forests there is a competition among species and genotypes to interact with CO₂ and O₃ to influence carbon cycling. (Diaz-de-ouijano et al, 2012) The focal species for this experiment was trembling aspen, trembling aspen is north America's most widespread tree species and O₃ exposure is a common component of north America's forest. Aspen was planted in mixed species (aspen- birch, aspen-maple) or either in mixed genotype (five clones varying in sensitivity to CO₂ and O₃) communities, representing common forest type in the north central United States

Figure 5. Successional responses to O₃ by individual species within the simulated forest. Sensitive species: *Liriodendron tulipifera* (a) and *Acer rubrum* (b). Intermediate species: *Acer saccharum* (c) and *Quercus velutina*

(d). Tolerant species: *Quercus alba* (e) and *Fagus grandifolia* (f). Dark and red line denote without and with O₃ stress, respectively. (Wang et al. 2016)



In this experiment, these all forests were exposed to factorial combination of increased CO₂ and O₃ for 11 years and advanced from open grow seedling < 0.25 m in height to closed canopy stands that were >8 m tall. Our first objective of doing this aspen, FACE experiment was to quantify the effects of increased CO₂ and O₃ on ecosystem C content. At the first phase of experiment in 1997 we presumed that ecosystem C content can be extended by increased CO₂ and lowered by elevated O₃. According to our limited knowledge, we further presumed that CO₂ and O₃ would show no such interaction such that two gases would have on ecosystem C content.

CONCLUSION

In this review global average C-N feedback of forest ecosystem are shown, where gradual gathering of tree wood, with long revenue times, lies between 5 to 25 Kg N⁻¹ which is totally dependent on forest type. N deposition leads to lower growth responses, but inauspicious impacts occur below 20-30 kg N ha⁻¹ yr⁻¹. This will boost the nitrogen loads associated with other environmental problems, including higher nitrogen leaching and decline in biodiversity. The values later vary from 10 and 20 Kg N ha⁻¹ for broadleaved forests and for coniferous forest it is 5-15 Kg N ha⁻¹. (Vries et al. 2014). Physical and chemical features of the geosphere are influenced by forest carbon cycle thus it is considered as large and important. Ecological models till now predict carbon cycle at big scale had only contemplate the outcome of nitrogen but it does not take others into account such as phosphorus or pH of the soil, which is related to the availability of nutrients. New ideas and modelling are done to evaluate the importance of forest biosphere-climate interaction for future climate change. Rather than model projection, attention to understanding processes.

Prediction of better constraints for the estimation of upcoming changes in forest biosphere and climate will be done by upgraded correlation between facts and models. (Hui et al. 2015)

REFERENCES:

- 1) Hisano, M., Searle, E. B., & Chen, H. Y. (2018). Biodiversity as a solution to mitigate climate change impacts on the functioning of forest ecosystems. *Biological Reviews*, 93(1), 439-456. [10.1111/brv.12351](https://doi.org/10.1111/brv.12351)
- 2) Hui, D., Deng, Q., Tian, H., & Luo, Y. (2017). Climate change and carbon sequestration in forest ecosystems. *Handbook of climate change mitigation and adaptation*, 555, 594.

[10.1007/978-3-319-14409-2_13](https://doi.org/10.1007/978-3-319-14409-2_13)

- 3) Lindquist, E. J., D'Annunzio, R., Gerrand, A., MacDicken, K., Achard, F., Beuchle, R., ... & Stibig, H. J. (2012). *Global forest land-use change 1990-2005* (No. 169, pp. xi+-40).
- 4) Hobbs, T. J., Neumann, C. R., Meyer, W. S., Moon, T., & Bryan, B. A. (2016). Models of reforestation productivity and carbon sequestration for land use and climate change adaptation planning in South Australia. *Journal of environmental management*, 181, 279-288. <https://doi.org/10.1016/j.jenvman.2016.06.049>
- 5) Peñuelas, J., Sardans, J., Estiarte, M., Ogaya, R., Carnicer, J., Coll, M., ... & Jump, A. S. (2013). Evidence of current impact of climate change on life: a walk from genes to the biosphere. *Global change biology*, 19(8), 2303-2338. [10.1111/gcb.12143](https://doi.org/10.1111/gcb.12143)
- 6) FAO, F. (2015). *Global forest resources assessment 2015 Desk reference*. Food and agriculture organization of the United Nations, Rome.
- 7) van Der Plas, F., Manning, P., Soliveres, S., Allan, E., Scherer-Lorenzen, M., Verheyen, K., ... & Fischer, M. (2016). Biotic homogenization can decrease landscape-scale forest multifunctionality. *Proceedings of the National Academy of Sciences*, 113(13), 3557-3562. <https://doi.org/10.1073/pnas.1517903113>
- 8) Mori, A. S., Lertzman, K. P., & Gustafsson, L. (2017). Biodiversity and ecosystem services in forest ecosystems: a research agenda for applied forest ecology. *Journal of Applied Ecology*, 54(1), 12-27. [10.1111/1365-2664.12669](https://doi.org/10.1111/1365-2664.12669)
- 9) Cardinale. (2012). Contextualizing context for synthetic biology—identifying causes of failure of synthetic biological systems. *Biotechnology journal*, 7(7), 856-866. <https://doi.org/10.1002/biot.201200085>
- 10) Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., ... & Eisenhauer, N. (2015). Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature*, 526(7574), 574-577. [10.1038/nature15374](https://doi.org/10.1038/nature15374)
- 11) Zuidema, P. A., Baker, P. J., Groenendijk, P., Schippers, P., van der Sleen, P., Vlam, M., & Sterck, F. (2013). Tropical forests and global change: filling knowledge gaps. *Trends in plant science*, 18(8), 413-419. <https://doi.org/10.1016/j.tplants.2013.05.006>
- 12) Peters, E. B., Wythers, K. R., Zhang, S., Bradford, J. B., & Reich, P. B. (2013). Potential climate change impacts temperate forest ecosystem processes. *Canadian Journal of Forest Research*, 43(10), 939-950. <https://doi.org/10.1139/cjfr-2013-0013>
- 13) Tian, H., Chen, G., Zhang, C., Liu, M., Sun, G., Chappelka, A., ... & Vance, E. (2012). Century-scale responses of ecosystem carbon storage and flux to multiple environmental changes in the southern United States. *Ecosystems*, 15, 674-694.: [10.1007/s10021-012-9539-x](https://doi.org/10.1007/s10021-012-9539-x)
- 14) Gebeyehu, M. N., & Hirpo, F. H. (2019). Review of the effect of climate change on forest ecosystem. *International Journal of Environmental Sciences & Natural Resources*, 17(4), 126-129. [10.19080/IJESNR.2019.17.555968](https://doi.org/10.19080/IJESNR.2019.17.555968)
- 15) Dukes, J. S., Classen, A. T., Wan, S., & Langley, J. A. (2014). Using results from global change experiments to inform land model development and calibration. *New Phytologist*, 204(4), 744-746. <https://doi.org/10.1111/nph.13083>
- 16) Juneja, A., Ceballos, R. M., & Murthy, G. S. (2013). Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: a review. *Energies*, 6(9), 4607-4638. <https://doi.org/10.3390/en6094607>
- 17) Tian, N., Poudyal, N. C., Hodges, D. G., Young, T. M., & Hoyt, K. P. (2015). Understanding the factors influencing nonindustrial private forest landowner interest in supplying ecosystem services in Cumberland Plateau, Tennessee. *Forests*, 6(11), 3985-4000.: [10.3390/f6113985](https://doi.org/10.3390/f6113985)
- 18) Pohjanmies, T., Triviño, M., Le Tortorec, E., Mazziotta, A., Snäll, T., & Mönkkönen, M. (2017). Impacts of forestry on boreal forests: An ecosystem services perspective. *Ambio*, 46, 743-755. <https://doi.org/10.1007/s13280-017-0919-5>
- 19) Pan, Q., Wen, Z., Wu, T., Zheng, T., Yang, Y., Li, R., & Zheng, H. (2022). Trade-offs and synergies of forest ecosystem services from the perspective of plant functional traits: A systematic review. *Ecosystem Services*, 58, 101484. <https://doi.org/10.1016/j.ecoser.2022.101484>
- 20) Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., ... & Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest ecology and management*, 259(4), 660-684. <https://doi.org/10.1016/j.foreco.2009.09.001>
- 21) FAO. 2020. *Global Forest Resources Assessment 2020*. Rome. <https://www.fao.org/forest-resources-assessment/fra-2020/maps/en/>
- 22) . May, N., Guenther, E., & Haller, P. (2017). Environmental indicators for the evaluation of wood products in consideration of site-dependent aspects: A review and integrated approach. *Sustainability*, 9(10), 1897. <https://doi.org/10.3390/su9101897>
- 23) Peñuelas, J., Sardans, J., Filella, I., Estiarte, M., Llusà, J., Ogaya, R., ... & Terradas, J. (2017). Impacts of global change on Mediterranean forests and their services. *Forests*, 8(12), 463. [10.3390/F8120463](https://doi.org/10.3390/F8120463)

- 24) Chitsaz, N., & Azarnivand, A. (2017). Water scarcity management in arid regions based on an extended multiple criteria technique. *Water Resources Management*, 31, 233-250. DOI:[10.1007/s11269-016-1521-5](https://doi.org/10.1007/s11269-016-1521-5)
- 25) Mitrică, B., Mitrică, E., Enciu, P., & Mocanu, I. (2017). An approach for forecasting of public water scarcity at the end of the 21st century, in the Timiș Plain of Romania. *Technological Forecasting and Social Change*, 118, 258-269. [10.1016/j.techfore.2017.02.026](https://doi.org/10.1016/j.techfore.2017.02.026)
- 26) MacDicken, K. G., Sola, P., Hall, J. E., Sabogal, C., Tadoum, M., & de Wasseige, C. (2015). Global progress toward sustainable forest management. *Forest Ecology and Management*, 352, 47-56. <https://doi.org/10.1016/j.foreco.2015.02.005>
- 27) Damkjaer, S., & Taylor, R. (2017). The measurement of water scarcity: Defining a meaningful indicator. *Ambio*, 46(5), 513-531. [10.1007/s13280-017-0912-z](https://doi.org/10.1007/s13280-017-0912-z)
- 28) Hui, D., Deng, Q., Tian, H., & Luo, Y. (2017). Climate change and carbon sequestration in forest ecosystems. *Handbook of climate change mitigation and adaptation*, 555, 594. [10.1007/978-1-4614-6431-0_13-2](https://doi.org/10.1007/978-1-4614-6431-0_13-2)
- 29) Corinne Le Quéré, R. M., Andrew, R. M., Peters, G., Ciais, P., Friedlingstein, P., Jones, S., & Armeth, A. (2015). *Global carbon budget 2014*. Earth System Science Data, 2018, 10: 2141-2194. <https://doi.org/10.5194/essd-10-405-2018>
- 30) Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., ... & Hayes, D. (2011). A large and persistent carbon sink in the world's forests. *science*, 333(6045), 988-993. [10.1126/science.1201609](https://doi.org/10.1126/science.1201609)
- 31) Büntgen, U., Tegel, W., Kaplan, J. O., Schaub, M., Hagedorn, F., Bürgi, M., ... & Liebhold, A. (2014). Placing unprecedented recent fir growth in a European- wide and Holocene- long context. *Frontiers in Ecology and the Environment*, 12(2), 100-106. <https://doi.org/10.1890/130089>
- 32) de Vries, W., Du, E., & Butterbach-Bahl, K. (2014). Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems. *Current Opinion in Environmental Sustainability*, 9, 90-104. <https://doi.org/10.1016/j.cosust.2014.09.001>
- 33) Lu, M., Zhou, X., Luo, Y., Yang, Y., Fang, C., Chen, J., & Li, B. (2011). Minor stimulation of soil carbon storage by nitrogen addition: a meta-analysis. *Agriculture, ecosystems & environment*, 140(1-2), 234-244. <https://doi.org/10.1016/j.agee.2010.12.010>
- 34) Fernández-Martínez, M., Vicca, S., Janssens, I. A., Sardans, J., Luysaert, S., Campioli, M., ... & Peñuelas, J. (2014). Nutrient availability as the key regulator of global forest carbon balance. *Nature Climate Change*, 4(6), 471-476. [10.1038/nclimate2177](https://doi.org/10.1038/nclimate2177)
- 35) Huang, L., Zhou, M., Lv, J., & Chen, K. (2020). Trends in global research in forest carbon sequestration: A bibliometric analysis. *Journal of Cleaner Production*, 252, 119908. <https://doi.org/10.1016/j.jclepro.2019.119908>
- 36) Drake, J. E., Gallet- Budynek, A., Hofmockel, K. S., Bernhardt, E. S., Billings, S. A., Jackson, R. B., ... & Finzi, A. C. (2011). Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long- term enhancement of forest productivity under elevated CO₂. *Ecology letters*, 14(4), 349-357. [10.1111/j.1461-0248.2011.01593.x](https://doi.org/10.1111/j.1461-0248.2011.01593.x)
- 37) Hartley, I. P., Hill, T. C., Chadburn, S. E., & Hugelius, G. (2021). Temperature effects on carbon storage are controlled by soil stabilisation capacities. *Nature Communications*, 12(1), 6713. <https://doi.org/10.1038/s41467-021-27101-1>
- 38) Drake, J. E., Gallet- Budynek, A., Hofmockel, K. S., Bernhardt, E. S., Billings, S. A., Jackson, R. B., ... & Finzi, A. C. (2011). Increases in the flux of carbon belowground stimulate nitrogen uptake and sustain the long- term enhancement of forest productivity under elevated CO₂. *Ecology letters*, 14(4), 349-357. <https://doi.org/10.1111/j.1461-0248.2011.01593.x>
- 39) Yu, Z. C. (2012). Northern peatland carbon stocks and dynamics: a review. *Biogeosciences*, 9(10), 4071-4085. <https://doi.org/10.5194/bg-9-4071-2012>
- 40) Doetterl, S., Stevens, A., Six, J., Merckx, R., Van Oost, K., Casanova Pinto, M., ... & Boeckx, P. (2015). Soil carbon storage is controlled by interactions between geochemistry and climate. *Nature Geoscience*, 8(10), 780-783. [10.1038/NCEO2516](https://doi.org/10.1038/NCEO2516)
- 41) Dungait, J. A., Hopkins, D. W., Gregory, A. S., & Whitmore, A. P. (2012). Soil organic matter turnover is governed by accessibility not recalcitrance. *Global Change Biology*, 18(6), 1781-1796. <https://doi.org/10.1111/j.1365-2486.2012.02665.x>
- 42) González-Domínguez, B., Niklaus, P. A., Studer, M. S., Hagedorn, F., Wacker, L., Haghypour, N., ... & Abiven, S. (2019). Temperature and moisture are minor drivers of regional-scale soil organic carbon dynamics. *Scientific reports*, 9(1), 6422. <https://doi.org/10.1038/s41598-019-42629-5>
- 43) Wiesmeier, M., Hübner, R., Spörlein, P., Geuß, U., Hangen, E., Reischl, A., ... & Kögel- Knabner, I. (2014). Carbon sequestration potential of soils in southeast Germany derived from stable soil organic carbon saturation. *Global change biology*, 20(2), 653-665. <https://doi.org/10.1111/gcb.12384>
- 44) Conant, R. T., Ryan, M. G., Ågren, G. I., Birge, H. E., Davidson, E. A., Eliasson, P. E., ... & Bradford, M. A. (2011). Temperature and soil organic matter decomposition rates—synthesis of current knowledge and a way forward. *Global change biology*, 17(11), 3392-3404. <https://doi.org/10.1111/j.1365-2486.2011.02496.x>
- 45) Jia, Y., Kuzyakov, Y., Wang, G., Tan, W., Zhu, B., & Feng, X. (2020). Temperature sensitivity of decomposition of soil organic matter fractions increases with their turnover time. *Land Degradation & Development*, 31(5), 632-645. <https://doi.org/10.1002/ldr.3477>

- 46) Gong, H., Wang, G., Fan, C., Zhuo, X., Sha, L., Kuang, Z., ... & Cheng, T. (2024). Temporal accumulation and lag effects of precipitation on carbon fluxes in terrestrial ecosystems across semi-arid regions in China. *Agricultural and Forest Meteorology*, 356, 110189. <https://doi.org/10.1016/j.agrformet.2024.110189> [Get rights and content](#)
- 47) Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T., & Liu, Z. Y. (2014). The global monsoon across timescales: coherent variability of regional monsoons. *Climate of the Past*, 10(6), 2007-2052. <https://doi.org/10.5194/cp-10-2007-2014>
- 48) Huang, L., Liu, J., Shao, Q., & Xu, X. (2012). Carbon sequestration by forestation across China: Past, present, and future. *Renewable and Sustainable Energy Reviews*, 16(2), 1291-1299. <https://doi.org/10.1016/j.rser.2011.10.004>
- 49) Pal, S. C., Chatterjee, U., Chakraborty, R., Roy, P., Chowdhuri, I., Saha, A., ... & Islam, M. K. (2023). Anthropogenic drivers induced desertification under changing climate: Issues, policy interventions, and the way forward. *Progress in Disaster Science*, 20, 100303. <https://doi.org/10.1016/j.pdisas.2023.100303>
- 50) Knowles, J. F., Scott, R. L., Minor, R. L., & Barron-Gafford, G. A. (2020). Ecosystem carbon and water cycling from a sky island montane forest. *Agricultural and Forest Meteorology*, 281, 107835. <https://doi.org/10.1016/j.agrformet.2019.107835>
- 51) Ahlström, A., Raupach, M. R., Schurgers, G., Smith, B., Arneeth, A., Jung, M., ... & Zeng, N. (2015). The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink. *Science*, 348(6237), 895-899. [10.1126/science.1266688](https://doi.org/10.1126/science.1266688)
- 52) Ameer, I., Kubar, K. A., Ali, Q., Ali, S., Khan, T., Shahzad, K., ... & Talpur, K. H. (2023). Land degradation resistance potential of a dry, semiarid region in relation to soil organic carbon stocks, carbon management index, and soil aggregate stability. *Land Degradation & Development*, 34(3), 624-636. <https://doi.org/10.1002/ldr.4480>
- 53) Chen, Y., Feng, X., Tian, H., Wu, X., Gao, Z., Feng, Y., ... & Fu, B. (2021). Accelerated increase in vegetation carbon sequestration in China after 2010: A turning point resulting from climate and human interaction. *Global Change Biology*, 27(22), 5848-5864. <https://doi.org/10.1111/gcb.15854>
- 54) Jayara, A. S., Pandey, S., & Bhatt, M. K. (2023). Carbon sequestration as influenced by cropping practices: A review. *Agricultural Reviews*, 44(4), 477-484. [10.18805/ag.R-2253](https://doi.org/10.18805/ag.R-2253)
- 55) Zhang, Y., He, B., Chen, R., Zhang, H., Fan, C., Yin, J., & Li, Y. (2024). The potential of optical and SAR time-series data for the improvement of aboveground biomass carbon estimation in Southwestern China's evergreen coniferous forests. *GIScience & Remote Sensing*, 61(1), 2345438. <https://doi.org/10.1080/15481603.2024.2345438>
- 56) Zhan, T., Zhao, H., Zhang, J., Cheng, C., & Zhang, Z. (2023). Differential effects of grazing intensity on carbon sequestration in arid versus humid grasslands across China. *Science of The Total Environment*, 881, 163221. <https://doi.org/10.1016/j.scitotenv.2023.163221>
- 57) Gao, Y., Tian, J., Pang, Y., & Liu, J. (2017). Soil inorganic carbon sequestration following afforestation is probably induced by pedogenic carbonate formation in Northwest China. *Frontiers in plant science*, 8, 1282. <https://doi.org/10.3389/fpls.2017.01282>
- 58) Liu, H., Sun, Z., Cui, Y., Dong, Y., He, P., An, S., & Zhang, X. (2024). Storage, pattern and driving factors of soil organic carbon in the desert rangelands of northern Xinjiang, north-west China. *Frontiers of Earth Science*, 1-13. <https://doi.org/10.1007/s11707-022-0978-1> RESEARCH ARTICLE
- 59) Liu, P., Zha, T., Jia, X., Black, T. A., Jassal, R. S., Ma, J., ... & Wu, Y. (2019). Different effects of spring and summer droughts on ecosystem carbon and water exchanges in a semiarid shrubland ecosystem in Northwest China. *Ecosystems*, 22, 1869-1885. <https://doi.org/10.1007/s10021-019-00379-5>
- 60) Wang, W., Wang, B. Z., Zhou, R., Ullah, A., Zhao, Z. Y., Wang, P. Y., ... & Xiong, Y. C. (2022). Biocrusts as a nature-based strategy (NbS) improve soil carbon and nitrogen stocks and maize productivity in semiarid environment. *Agricultural Water Management*, 270, 107742. <https://doi.org/10.1016/j.agwat.2022.107742>
- 61) Prajapati, S. K., Choudhary, S., Kumar, V., Dayal, P., Srivastava, R., Gairola, A., & Borate, R. B. (2023). Carbon Sequestration: A Key Strategy for Climate Change Mitigation towards a Sustainable Future. *Emerg. Trnd. Clim. Chng.*, 2(2), 1-14. <http://dx.doi.org/10.18782/2583-4770.128>
- 62) Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., & Emberson, L. D. (2012). The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annual review of plant biology*, 63(1), 637-661. [10.1146/annurev-arplant-042110-103829](https://doi.org/10.1146/annurev-arplant-042110-103829)
- 63) Díaz-de-Quijano, M., Schaub, M., Bassin, S., Volk, M., & Peñuelas, J. (2012). Ozone visible symptoms and reduced root biomass in the subalpine species *Pinus uncinata* after two years of free-air ozone fumigation. *Environmental Pollution*, 169, 250-257. [10.1016/j.envpol.2012.02.011](https://doi.org/10.1016/j.envpol.2012.02.011)
- 64) Wang, B., Shugart, H. H., Shuman, J. K., & Lerdau, M. T. (2016). Forests and ozone: Productivity, carbon storage and feedbacks. *Scientific Reports*, 6(1), 22133. [10.1038/srep22133](https://doi.org/10.1038/srep22133)
- 65) de Vries, W., Du, E., & Butterbach-Bahl, K. (2014). Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems. *Current Opinion in Environmental Sustainability*, 9, 90-104. <https://doi.org/10.1016/j.cosust.2014.09.001>
- 66) Hui, D., Deng, Q., Tian, H., & Luo, Y. (2017). Climate change and carbon sequestration in forest ecosystems. *Handbook of climate change mitigation and adaptation*, 555, 594.: [10.1007/978-1-4614-6431-0_13-2](https://doi.org/10.1007/978-1-4614-6431-0_13-2)